

06/27/00



30852 U.S. PRO

06-28-00

Box 56A

BGI-122CP

Page 1 of 2

Customer No. 000959

Case Docket No. BGI-122CP

Assistant Commissioner for Patents
BOX PATENT APPLICATION
Washington, D.C. 20231

"Express Mail" Mailing Label Number	EL 373 207 242 US
Date of Deposit	June 27, 2000
<p>I hereby certify that this transmittal letter and the papers referred to as being enclosed therein are being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.110 on the date indicated above and is addressed to the Assistant Commissioner for Patents, Box Patent Application, Washington, DC 20231.</p>	
<p><i>Larry Taylor</i> Signature</p>	
<p>Larry Taylor Please Print Name of Person Signing</p>	

Sir:

Transmitted herewith for filing is the patent application of:

Inventors: Markus Pompejus *et al.*

For: "*Corynebacterium Glutamicum* Genes Encoding Phosphoenolpyruvate: Sugar Phosphotransferase System Proteins"

Enclosed are:

- ☒ 59 pages of specification, 5 pages of claims and 1 page of abstract;
- ☒ 1 page of Table 1;
- ☒ 16 pages of Table 2;
- ☒ 6 pages of Table 3;
- ☒ 1 page of Table 4;
- ☒ 7 pages of Appendix A;
- ☒ 3 pages of Appendix B;
- ☒ 47 pages of Sequence Listing;
- ☒ Diskette Containing Sequence Listing;
- ☒ Transmittal Letter for Diskette Containing Sequence Listing;
- ☒ An *unexecuted* Declaration, Petition and Power of Attorney; and
- ☒ A pre-paid acknowledgment postcard.

The filing fee has been calculated as shown below:

	(Col. 1)	(Col. 2)
FOR:	NO. FILED	NO. EXTRA
BASIC FEE	////////////////////	
TOTAL CLAIMS	52 - 20	= 32
INDEP. CLAIMS	15 - 3	= 12
<input checked="" type="checkbox"/> MULTIPLE DEPENDENT CLAIMS PRESENTED		

* If the difference in Col. 2 is less than zero,
enter "0" in Col. 2.

SMALL ENTITY	
RATE	FEE
////////	\$
x 9=	\$
x 39	\$
+130	\$
TOTAL	0

OTHER THAN SMALL ENTITY	
RATE	FEE
////////	\$ 690
x 18=	\$ 576
x 78	\$ 936
+260	\$ 260
TOTAL	\$2462.00

05604231 062700

☒ FILING FEES ARE NOT BEING PAID AT THIS TIME.

☐ The Commissioner is hereby authorized to charge payment of the following fees associated with this communication or credit any overpayment to Deposit Account No.

☐ Any additional filing fees required under 37 C.F.R. 1.16.

☐ Any patent application processing fees under 37 C.F.R. 1.17.

☐ The Commissioner is hereby authorized to charge payment of the following fees during the pendency of this application or credit any overpayment to Deposit Account No. _____

A duplicate copy of this sheet is enclosed.

☐ Any patent application processing fees under 37 C.F.R. 1.17.

☐ The issue fee set in 37 C.F.R. 1.18 at or before mailing of the Notice of Allowance, pursuant to 37 C.F.R 1.31 I(b).

☐ Any filing fees under 37 C.F.R. 1.16 for presentation of extra claims.

☐ A check in the amount of \$ _____ to cover the recording of assignment documents is also enclosed.

☒ Address all future communications (May only be completed by applicant, or attorney or agent of record) to **Elizabeth A. Hanley, Esq.** at **Customer Number: 000959** whose address is:

Lahive & Cockfield, LLP
28 State Street
Boston, Massachusetts 02109

Date: **June 27, 2000**

LAHIVE & COCKFIELD, LLP

28 State Street
Boston, MA 02109
(617) 227-7400 (Tel.)
(617) 742-4214 (Fax)

Attorneys at Law

By:

Elizabeth A. Hanley, Esq.
Registration No. 33,505
Attorney for Applicants

**CORYNEBACTERIUM GLUTAMICUM GENES ENCODING
PHOSPHOENOLPYRUVATE: SUGAR PHOSPHOTRANSFERASE SYSTEM
PROTEINS**

5 Related Applications

This application claims priority to U.S. Provisional Patent Application No.: 60/142,691, filed on July 1, 1999, and also to U.S. Provisional Patent Application No.: 60/150,310, filed on August 23, 1999, incorporated herein in their entirety by this reference. This application also claims priority to German Patent Application No.: 10 19942095.5, filed on September 3, 1999, and also to German Patent Application No.: 19942097.1, filed on September 3, 1999, incorporated herein in their entirety by this reference.

Background of the Invention

15 Certain products and by-products of naturally-occurring metabolic processes in cells have utility in a wide array of industries, including the food, feed, cosmetics, and pharmaceutical industries. These molecules, collectively termed 'fine chemicals', include organic acids, both proteinogenic and non-proteinogenic amino acids, nucleotides and nucleosides, lipids and fatty acids, diols, carbohydrates, aromatic
20 compounds, vitamins and cofactors, and enzymes. Their production is most conveniently performed through large-scale culture of bacteria developed to produce and secrete large quantities of a particular desired molecule. One particularly useful organism for this purpose is *Corynebacterium glutamicum*, a gram positive, nonpathogenic bacterium. Through strain selection, a number of mutant strains have
25 been developed which produce an array of desirable compounds. However, selection of strains improved for the production of a particular molecule is a time-consuming and difficult process.

Summary of the Invention

30 The invention provides novel bacterial nucleic acid molecules which have a variety of uses. These uses include the identification of microorganisms which can be used to produce fine chemicals, the modulation of fine chemical production in *C. glutamicum* or related bacteria, the typing or identification of *C. glutamicum* or related bacteria, as reference points for mapping the *C. glutamicum* genome, and as markers for
35 transformation. These novel nucleic acid molecules encode proteins, referred to herein as phosphoenolpyruvate:sugar phosphotransferase system (PTS) proteins.

09604231.062700

5 *C. glutamicum* is a gram positive, aerobic bacterium which is commonly used in industry for the large-scale production of a variety of fine chemicals, and also for the degradation of hydrocarbons (such as in petroleum spills) and for the oxidation of terpenoids. The PTS nucleic acid molecules of the invention, therefore, can be used to identify microorganisms which can be used to produce fine chemicals, e.g., by fermentation processes. Modulation of the expression of the PTS nucleic acids of the invention, or modification of the sequence of the PTS nucleic acid molecules of the invention, can be used to modulate the production of one or more fine chemicals from a microorganism (e.g., to improve the yield or production of one or more fine chemicals from a *Corynebacterium* or *Brevibacterium* species).

10 The PTS nucleic acids of the invention may also be used to identify an organism as being *Corynebacterium glutamicum* or a close relative thereof, or to identify the presence of *C. glutamicum* or a relative thereof in a mixed population of microorganisms. The invention provides the nucleic acid sequences of a number of *C. glutamicum* genes; by probing the extracted genomic DNA of a culture of a unique or mixed population of microorganisms under stringent conditions with a probe spanning a region of a *C. glutamicum* gene which is unique to this organism, one can ascertain whether this organism is present. Although *Corynebacterium glutamicum* itself is nonpathogenic, it is related to species pathogenic in humans, such as *Corynebacterium diphtheriae* (the causative agent of diphtheria); the detection of such organisms is of significant clinical relevance.

15 The PTS nucleic acid molecules of the invention may also serve as reference points for mapping of the *C. glutamicum* genome, or of genomes of related organisms. Similarly, these molecules, or variants or portions thereof, may serve as markers for genetically engineered *Corynebacterium* or *Brevibacterium* species.

20 The PTS proteins encoded by the novel nucleic acid molecules of the invention are capable of, for example, transporting high-energy carbon-containing molecules such as glucose into *C. glutamicum*, or of participating in intracellular signal transduction in this microorganism. Given the availability of cloning vectors for use in *Corynebacterium glutamicum*, such as those disclosed in Sinskey *et al.*, U.S. Patent No. 4,649,119, and techniques for genetic manipulation of *C. glutamicum* and the related *Brevibacterium* species (e.g., *lactofermentum*) (Yoshihama *et al.*, *J. Bacteriol.* 162: 591-597 (1985); Katsumata *et al.*, *J. Bacteriol.* 159: 306-311 (1984); and Santamaria *et al.*, *J. Gen. Microbiol.* 130: 2237-2246 (1984)), the nucleic acid molecules of the invention may be utilized in the genetic engineering of this organism to make it a better or more efficient producer of one or more fine chemicals.

09604231.062700

The PTS molecules of the invention may be modified such that the yield, production, and/or efficiency of production of one or more fine chemicals is improved. For example, by modifying a PTS protein involved in the uptake of glucose such that it is optimized in activity, the quantity of glucose uptake or the rate at which glucose is translocated into the cell may be increased. The breakdown of glucose and other sugars within the cell provides energy that may be used to drive energetically unfavorable biochemical reactions, such as those involved in the biosynthesis of fine chemicals. This breakdown also provides intermediate and precursor molecules necessary for the biosynthesis of certain fine chemicals, such as amino acids, vitamins and cofactors. By increasing the amount of intracellular high-energy carbon molecules through modification of the PTS molecules of the invention, one may therefore increase both the energy available to perform metabolic pathways necessary for the production of one or more fine chemicals, and also the intracellular pools of metabolites necessary for such production.

Further, the PTS molecules of the invention may be involved in one or more intracellular signal transduction pathways which may affect the yields and/or rate of production of one or more fine chemical from *C. glutamicum*. For example, proteins necessary for the import of one or more sugars from the extracellular medium (e.g., HPr, Enzyme I, or a member of an Enzyme II complex) are frequently posttranslationally modified upon the presence of a sufficient quantity of the sugar in the cell, such that they are no longer able to import that sugar. While this quantity of sugar at which the transport system is shut off may be sufficient to sustain the normal functioning of the cell, it may be limiting for the overproduction of the desired fine chemical. Thus, it may be desirable to modify the PTS proteins of the invention such that they are no longer responsive to such negative regulation, thereby permitting greater intracellular concentrations of one or more sugars to be achieved, and, by extension, more efficient production or greater yields of one or more fine chemicals from organisms containing such mutant PTS proteins.

This invention provides novel nucleic acid molecules which encode proteins, referred to herein as phosphoenolpyruvate:sugar phosphotransferase system (PTS) proteins, which are capable of, for example, participating in the import of high-energy carbon molecules (e.g., glucose, fructose, or sucrose) into *C. glutamicum*, and/or of participating in one or more *C. glutamicum* intracellular signal transduction pathways. Nucleic acid molecules encoding a PTS protein are referred to herein as PTS nucleic acid molecules. In a preferred embodiment, the PTS protein participates in the import of high-energy carbon molecules (e.g., glucose, fructose, or sucrose) into *C. glutamicum*, and also may participate in one or more *C. glutamicum* intracellular signal transduction

the amino acid sequences of Appendix B and is able to participate in the import of high-energy carbon molecules (e.g., glucose, fructose, or sucrose) into *C. glutamicum*, and/or to participate in one or more *C. glutamicum* intracellular signal transduction pathways, or possesses one or more of the activities set forth in Table 1, and which also includes

5 heterologous nucleic acid sequences encoding a heterologous polypeptide or regulatory regions.

In another embodiment, the isolated nucleic acid molecule is at least 15 nucleotides in length and hybridizes under stringent conditions to a nucleic acid molecule comprising a nucleotide sequence of Appendix A. Preferably, the isolated

10 nucleic acid molecule corresponds to a naturally-occurring nucleic acid molecule. More preferably, the isolated nucleic acid encodes a naturally-occurring *C. glutamicum* PTS protein, or a biologically active portion thereof.

Another aspect of the invention pertains to vectors, e.g., recombinant expression vectors, containing the nucleic acid molecules of the invention, and host cells into which

15 such vectors have been introduced. In one embodiment, such a host cell is used to produce a PTS protein by culturing the host cell in a suitable medium. The PTS protein can be then isolated from the medium or the host cell.

Yet another aspect of the invention pertains to a genetically altered microorganism in which a PTS gene has been introduced or altered. In one

20 embodiment, the genome of the microorganism has been altered by the introduction of a nucleic acid molecule of the invention encoding wild-type or mutated PTS sequence as a transgene. In another embodiment, an endogenous PTS gene within the genome of the microorganism has been altered, e.g., functionally disrupted, by homologous recombination with an altered PTS gene. In another embodiment, an endogenous or

25 introduced PTS gene in a microorganism has been altered by one or more point mutations, deletions, or inversions, but still encodes a functional PTS protein. In still another embodiment, one or more of the regulatory regions (e.g., a promoter, repressor, or inducer) of a PTS gene in a microorganism has been altered (e.g., by deletion, truncation, inversion, or point mutation) such that the expression of the PTS gene is

30 modulated. In a preferred embodiment, the microorganism belongs to the genus *Corynebacterium* or *Brevibacterium*, with *Corynebacterium glutamicum* being particularly preferred. In a preferred embodiment, the microorganism is also utilized for the production of a desired compound, such as an amino acid, with lysine being particularly preferred.

35 In another aspect, the invention provides a method of identifying the presence or activity of *Corynebacterium diphtheriae* in a subject. This method includes detection of one or more of the nucleic acid or amino acid sequences of the invention (e.g., the

09604231.000700

sequences set forth in Appendix A or Appendix B) in a subject, thereby detecting the presence or activity of *Corynebacterium diphtheriae* in the subject.

Still another aspect of the invention pertains to an isolated PTS protein or a portion, e.g., a biologically active portion, thereof. In a preferred embodiment, the isolated PTS protein or portion thereof can participate in the import of high-energy

- 5 carbon molecules (e.g., glucose, fructose, or sucrose) into *C. glutamicum*, and also may participate in one or more *C. glutamicum* intracellular signal transduction pathways. In another preferred embodiment, the isolated PTS protein or portion thereof is sufficiently homologous to an amino acid sequence of Appendix B such that the protein or portion
- 10 thereof maintains the ability to participate in the import of high-energy carbon molecules (e.g., glucose, fructose, or sucrose) into *C. glutamicum*, and /or to participate in one or more *C. glutamicum* intracellular signal transduction pathways.

- The invention also provides an isolated preparation of a PTS protein. In preferred embodiments, the PTS protein comprises an amino acid sequence of Appendix
- 15 B. In another preferred embodiment, the invention pertains to an isolated full length protein which is substantially homologous to an entire amino acid sequence of Appendix B (encoded by an open reading frame set forth in Appendix A). In yet another embodiment, the protein is at least about 50%, preferably at least about 60%, and more preferably at least about 70%, 80%, or 90%, and most preferably at least about 95%,
- 20 96%, 97%, 98%, or 99% or more homologous to an entire amino acid sequence of Appendix B. In other embodiments, the isolated PTS protein comprises an amino acid sequence which is at least about 50% or more homologous to one of the amino acid sequences of Appendix B and is able to participate in the import of high-energy carbon molecules (e.g., glucose, fructose, or sucrose) into *C. glutamicum*, and/or to participate
- 25 in one or more *C. glutamicum* intracellular signal transduction pathways, or has one or more of the activities set forth in Table 1.

- Alternatively, the isolated PTS protein can comprise an amino acid sequence which is encoded by a nucleotide sequence which hybridizes, e.g., hybridizes under stringent conditions, or is at least about 50%, preferably at least about 60%, more
- 30 preferably at least about 70%, 80%, or 90%, and even more preferably at least about 95%, 96%, 97%, 98%, or 99% or more homologous, to a nucleotide sequence of Appendix B. It is also preferred that the preferred forms of PTS proteins also have one or more of the PTS bioactivities described herein.

- The PTS polypeptide, or a biologically active portion thereof, can be operatively
- 35 linked to a non-PTS polypeptide to form a fusion protein. In preferred embodiments, this fusion protein has an activity which differs from that of the PTS protein alone. In other preferred embodiments, this fusion protein results in increased yields, production,

03604231.002700

and/or efficiency of production of a desired fine chemical from *C. glutamicum*. In particularly preferred embodiments, integration of this fusion protein into a host cell modulates the production of a desired compound from the cell.

- In another aspect, the invention provides methods for screening molecules which
- 5 modulate the activity of a PTS protein, either by interacting with the protein itself or a substrate or binding partner of the PTS protein, or by modulating the transcription or translation of a PTS nucleic acid molecule of the invention.

- Another aspect of the invention pertains to a method for producing a fine chemical. This method involves the culturing of a cell containing a vector directing the expression of a PTS nucleic acid molecule of the invention, such that a fine chemical is produced. In a preferred embodiment, this method further includes the step of obtaining
- 10 a cell containing such a vector, in which a cell is transfected with a vector directing the expression of a PTS nucleic acid. In another preferred embodiment, this method further includes the step of recovering the fine chemical from the culture. In a particularly
- 15 preferred embodiment, the cell is from the genus *Corynebacterium* or *Brevibacterium*, or is selected from those strains set forth in Table 3.

- Another aspect of the invention pertains to methods for modulating production of a molecule from a microorganism. Such methods include contacting the cell with an agent which modulates PTS protein activity or PTS nucleic acid expression such that a
- 20 cell associated activity is altered relative to this same activity in the absence of the agent. In a preferred embodiment, the cell is modulated for the uptake of one or more sugars, such that the yields or rate of production of a desired fine chemical by this microorganism is improved. The agent which modulates PTS protein activity can be an agent which stimulates PTS protein activity or PTS nucleic acid expression. Examples
- 25 of agents which stimulate PTS protein activity or PTS nucleic acid expression include small molecules, active PTS proteins, and nucleic acids encoding PTS proteins that have been introduced into the cell. Examples of agents which inhibit PTS activity or expression include small molecules, and antisense PTS nucleic acid molecules.

- Another aspect of the invention pertains to methods for modulating yields of a
- 30 desired compound from a cell, involving the introduction of a wild-type or mutant PTS gene into a cell, either maintained on a separate plasmid or integrated into the genome of the host cell. If integrated into the genome, such integration can random, or it can take place by homologous recombination such that the native gene is replaced by the introduced copy, causing the production of the desired compound from the cell to be
- 35 modulated. In a preferred embodiment, said yields are increased. In another preferred embodiment, said chemical is a fine chemical. In a particularly preferred embodiment,

00004231 002700

said fine chemical is an amino acid. In especially preferred embodiments, said amino acid is L-lysine.

Detailed Description of the Invention

The present invention provides PTS nucleic acid and protein molecules which are involved in the uptake of high-energy carbon molecules (e.g., sucrose, fructose, or glucose) into *C. glutamicum*, and may also participate in intracellular signal transduction pathways in this microorganism. The molecules of the invention may be utilized in the modulation of production of fine chemicals from microorganisms. Such modulation may be due to increased intracellular levels of high-energy molecules needed to produce, e.g., ATP, GTP and other molecules utilized to drive energetically unfavorable biochemical reactions in the cell, such as the biosynthesis of a fine chemical. This modulation of fine chemical production may also be due to the fact that the breakdown products of many sugars serve as intermediates or precursors for other biosynthetic pathways, including those of certain fine chemicals. Further, PTS proteins are known to participate in certain intracellular signal transduction pathways which may have regulatory activity for one or more fine chemical metabolic pathways; by manipulating these PTS proteins, one may thereby activate a fine chemical biosynthetic pathways or repress a fine chemical degradation pathway. Aspects of the invention are further explicated below.

I. Fine Chemicals

The term 'fine chemical' is art-recognized and includes molecules produced by an organism which have applications in various industries, such as, but not limited to, the pharmaceutical, agriculture, and cosmetics industries. Such compounds include organic acids, such as tartaric acid, itaconic acid, and diaminopimelic acid, both proteinogenic and non-proteinogenic amino acids, purine and pyrimidine bases, nucleosides, and nucleotides (as described *e.g.* in Kuninaka, A. (1996) Nucleotides and related compounds, p. 561-612, in Biotechnology vol. 6, Rehm *et al.*, eds. VCH: Weinheim, and references contained therein), lipids, both saturated and unsaturated fatty acids (*e.g.*, arachidonic acid), diols (*e.g.*, propane diol, and butane diol), carbohydrates (*e.g.*, hyaluronic acid and trehalose), aromatic compounds (*e.g.*, aromatic amines, vanillin, and indigo), vitamins and cofactors (as described in Ullmann's Encyclopedia of Industrial Chemistry, vol. A27, "Vitamins", p. 443-613 (1996) VCH: Weinheim and references therein; and Ong, A.S., Niki, E. & Packer, L. (1995) "Nutrition, Lipids, Health, and Disease" Proceedings of the UNESCO/Confederation of Scientific and Technological Associations in Malaysia, and the Society for Free Radical Research –

Asia, held Sept. 1-3, 1994 at Penang, Malaysia, AOCS Press, (1995)), enzymes, polyketides (Cane *et al.* (1998) *Science* 282: 63-68), and all other chemicals described in Gutcho (1983) *Chemicals by Fermentation*, Noyes Data Corporation, ISBN:

0818805086 and references therein. The metabolism and uses of certain of these fine chemicals are further explicated below.

A. Amino Acid Metabolism and Uses

Amino acids comprise the basic structural units of all proteins, and as such are essential for normal cellular functioning in all organisms. The term "amino acid" is art-recognized. The proteinogenic amino acids, of which there are 20 species, serve as structural units for proteins, in which they are linked by peptide bonds, while the nonproteinogenic amino acids (hundreds of which are known) are not normally found in proteins (see Ulmann's Encyclopedia of Industrial Chemistry, vol. A2, p. 57-97 VCH: Weinheim (1985)). Amino acids may be in the D- or L- optical configuration, though L-amino acids are generally the only type found in naturally-occurring proteins. Biosynthetic and degradative pathways of each of the 20 proteinogenic amino acids have been well characterized in both prokaryotic and eukaryotic cells (see, for example, Stryer, L. *Biochemistry*, 3rd edition, pages 578-590 (1988)). The 'essential' amino acids (histidine, isoleucine, leucine, lysine, methionine, phenylalanine, threonine, tryptophan, and valine), so named because they are generally a nutritional requirement due to the complexity of their biosyntheses, are readily converted by simple biosynthetic pathways to the remaining 11 'nonessential' amino acids (alanine, arginine, asparagine, aspartate, cysteine, glutamate, glutamine, glycine, proline, serine, and tyrosine). Higher animals do retain the ability to synthesize some of these amino acids, but the essential amino acids must be supplied from the diet in order for normal protein synthesis to occur.

Aside from their function in protein biosynthesis, these amino acids are interesting chemicals in their own right, and many have been found to have various applications in the food, feed, chemical, cosmetics, agriculture, and pharmaceutical industries. Lysine is an important amino acid in the nutrition not only of humans, but also of monogastric animals such as poultry and swine. Glutamate is most commonly used as a flavor additive (mono-sodium glutamate, MSG) and is widely used throughout the food industry, as are aspartate, phenylalanine, glycine, and cysteine. Glycine, L-methionine and tryptophan are all utilized in the pharmaceutical industry. Glutamine, valine, leucine, isoleucine, histidine, arginine, proline, serine and alanine are of use in both the pharmaceutical and cosmetics industries. Threonine, tryptophan, and D/L-methionine are common feed additives. (Leuchtenberger, W. (1996) *Amino acids - technical production and use*, p. 466-502 in Rehm *et al.* (eds.) *Biotechnology* vol. 6,

chapter 14a, VCH: Weinheim). Additionally, these amino acids have been found to be useful as precursors for the synthesis of synthetic amino acids and proteins, such as N-acetylcysteine, S-carboxymethyl-L-cysteine, (S)-5-hydroxytryptophan, and others described in Ulmann's Encyclopedia of Industrial Chemistry, vol. A2, p. 57-97, VCH:

5 Weinheim, 1985.

The biosynthesis of these natural amino acids in organisms capable of producing them, such as bacteria, has been well characterized (for review of bacterial amino acid biosynthesis and regulation thereof, see Umbarger, H.E.(1978) *Ann. Rev. Biochem.* 47: 533-606). Glutamate is synthesized by the reductive amination of α -ketoglutarate, an intermediate in the citric acid cycle. Glutamine, proline, and arginine are each subsequently produced from glutamate. The biosynthesis of serine is a three-step process beginning with 3-phosphoglycerate (an intermediate in glycolysis), and resulting in this amino acid after oxidation, transamination, and hydrolysis steps. Both cysteine and glycine are produced from serine; the former by the condensation of

10 ketoglutarate, an intermediate in the citric acid cycle. Glutamine, proline, and arginine are each subsequently produced from glutamate. The biosynthesis of serine is a three-step process beginning with 3-phosphoglycerate (an intermediate in glycolysis), and resulting in this amino acid after oxidation, transamination, and hydrolysis steps. Both cysteine and glycine are produced from serine; the former by the condensation of

15 homocysteine with serine, and the latter by the transferal of the side-chain β -carbon atom to tetrahydrofolate, in a reaction catalyzed by serine transhydroxymethylase. Phenylalanine, and tyrosine are synthesized from the glycolytic and pentose phosphate pathway precursors erythrose 4-phosphate and phosphoenolpyruvate in a 9-step biosynthetic pathway that differ only at the final two steps after synthesis of prephenate.

20 Tryptophan is also produced from these two initial molecules, but its synthesis is an 11-step pathway. Tyrosine may also be synthesized from phenylalanine, in a reaction catalyzed by phenylalanine hydroxylase. Alanine, valine, and leucine are all biosynthetic products of pyruvate, the final product of glycolysis. Aspartate is formed from oxaloacetate, an intermediate of the citric acid cycle. Asparagine, methionine,

25 threonine, and lysine are each produced by the conversion of aspartate. Isoleucine is formed from threonine. A complex 9-step pathway results in the production of histidine from 5-phosphoribosyl-1-pyrophosphate, an activated sugar.

Amino acids in excess of the protein synthesis needs of the cell cannot be stored, and are instead degraded to provide intermediates for the major metabolic pathways of

30 the cell (for review see Stryer, L. Biochemistry 3rd ed. Ch. 21 "Amino Acid Degradation and the Urea Cycle" p. 495-516 (1988)). Although the cell is able to convert unwanted amino acids into useful metabolic intermediates, amino acid production is costly in terms of energy, precursor molecules, and the enzymes necessary to synthesize them. Thus it is not surprising that amino acid biosynthesis is regulated by feedback inhibition,

35 in which the presence of a particular amino acid serves to slow or entirely stop its own production (for overview of feedback mechanisms in amino acid biosynthetic pathways, see Stryer, L. Biochemistry, 3rd ed. Ch. 24: "Biosynthesis of Amino Acids and Heme" p.

575-600 (1988)). Thus, the output of any particular amino acid is limited by the amount of that amino acid present in the cell.

B. Vitamin, Cofactor, and Nutraceutical Metabolism and Uses

- 5 Vitamins, cofactors, and nutraceuticals comprise another group of molecules which the higher animals have lost the ability to synthesize and so must ingest, although they are readily synthesized by other organisms, such as bacteria. These molecules are either bioactive substances themselves, or are precursors of biologically active substances which may serve as electron carriers or intermediates in a variety of
- 10 metabolic pathways. Aside from their nutritive value, these compounds also have significant industrial value as coloring agents, antioxidants, and catalysts or other processing aids. (For an overview of the structure, activity, and industrial applications of these compounds, see, for example, Ullman's Encyclopedia of Industrial Chemistry, "Vitamins" vol. A27, p. 443-613, VCH: Weinheim, 1996.) The term "vitamin" is art-
- 15 recognized, and includes nutrients which are required by an organism for normal functioning, but which that organism cannot synthesize by itself. The group of vitamins may encompass cofactors and nutraceutical compounds. The language "cofactor" includes nonproteinaceous compounds required for a normal enzymatic activity to occur. Such compounds may be organic or inorganic; the cofactor molecules of the
- 20 invention are preferably organic. The term "nutraceutical" includes dietary supplements having health benefits in plants and animals, particularly humans. Examples of such molecules are vitamins, antioxidants, and also certain lipids (e.g., polyunsaturated fatty acids).

- The biosynthesis of these molecules in organisms capable of producing them,
- 25 such as bacteria, has been largely characterized (Ullman's Encyclopedia of Industrial Chemistry, "Vitamins" vol. A27, p. 443-613, VCH: Weinheim, 1996; Michal, G. (1999) Biochemical Pathways: An Atlas of Biochemistry and Molecular Biology, John Wiley & Sons; Ong, A.S., Niki, E. & Packer, L. (1995) "Nutrition, Lipids, Health, and Disease" Proceedings of the UNESCO/Confederation of Scientific and Technological
- 30 Associations in Malaysia, and the Society for Free Radical Research - Asia, held Sept. 1-3, 1994 at Penang, Malaysia, AOCS Press: Champaign, IL X, 374 S).

- Thiamin (vitamin B₁) is produced by the chemical coupling of pyrimidine and thiazole moieties. Riboflavin (vitamin B₂) is synthesized from guanosine-5'-triphosphate (GTP) and ribose-5'-phosphate. Riboflavin, in turn, is utilized for the synthesis of flavin
- 35 mononucleotide (FMN) and flavin adenine dinucleotide (FAD). The family of compounds collectively termed 'vitamin B₆' (e.g., pyridoxine, pyridoxamine, pyridoxa-5'-phosphate, and the commercially used pyridoxin hydrochloride) are all derivatives of

the common structural unit, 5-hydroxy-6-methylpyridine. Pantothenate (pantothenic acid, (R)-(+)-N-(2,4-dihydroxy-3,3-dimethyl-1-oxobutyl)- β -alanine) can be produced either by chemical synthesis or by fermentation. The final steps in pantothenate biosynthesis consist of the ATP-driven condensation of β -alanine and pantoic acid. The enzymes responsible for the biosynthesis steps for the conversion to pantoic acid, to β -alanine and for the condensation to pantothenic acid are known. The metabolically active form of pantothenate is Coenzyme A, for which the biosynthesis proceeds in 5 enzymatic steps. Pantothenate, pyridoxal-5'-phosphate, cysteine and ATP are the precursors of Coenzyme A. These enzymes not only catalyze the formation of pantothenate, but also the production of (R)-pantoic acid, (R)-pantolacton, (R)-panthothenol (provitamin B₅), pantetheine (and its derivatives) and coenzyme A.

Biotin biosynthesis from the precursor molecule pimeloyl-CoA in microorganisms has been studied in detail and several of the genes involved have been identified. Many of the corresponding proteins have been found to also be involved in Fe-cluster synthesis and are members of the nifS class of proteins. Lipic acid is derived from octanoic acid, and serves as a coenzyme in energy metabolism, where it becomes part of the pyruvate dehydrogenase complex and the α -ketoglutarate dehydrogenase complex. The folates are a group of substances which are all derivatives of folic acid, which in turn is derived from L-glutamic acid, p-amino-benzoic acid and 6-methylpterin. The biosynthesis of folic acid and its derivatives, starting from the metabolism intermediates guanosine-5'-triphosphate (GTP), L-glutamic acid and p-amino-benzoic acid has been studied in detail in certain microorganisms.

Corrinoids (such as the cobalamines and particularly vitamin B₁₂) and porphyrines belong to a group of chemicals characterized by a tetrapyrrole ring system. The biosynthesis of vitamin B₁₂ is sufficiently complex that it has not yet been completely characterized, but many of the enzymes and substrates involved are now known. Nicotinic acid (nicotinate), and nicotinamide are pyridine derivatives which are also termed 'niacin'. Niacin is the precursor of the important coenzymes NAD (nicotinamide adenine dinucleotide) and NADP (nicotinamide adenine dinucleotide phosphate) and their reduced forms.

The large-scale production of these compounds has largely relied on cell-free chemical syntheses, though some of these chemicals have also been produced by large-scale culture of microorganisms, such as riboflavin, Vitamin B₆, pantothenate, and biotin. Only Vitamin B₁₂ is produced solely by fermentation, due to the complexity of its synthesis. *In vitro* methodologies require significant inputs of materials and time, often at great cost.

C. Purine, Pyrimidine, Nucleoside and Nucleotide Metabolism and Uses

Purine and pyrimidine metabolism genes and their corresponding proteins are important targets for the therapy of tumor diseases and viral infections. The language "purine" or "pyrimidine" includes the nitrogenous bases which are constituents of nucleic acids, co-enzymes, and nucleotides. The term "nucleotide" includes the basic structural units of nucleic acid molecules, which are comprised of a nitrogenous base, a pentose sugar (in the case of RNA, the sugar is ribose; in the case of DNA, the sugar is D-deoxyribose), and phosphoric acid. The language "nucleoside" includes molecules which serve as precursors to nucleotides, but which are lacking the phosphoric acid moiety that nucleotides possess. By inhibiting the biosynthesis of these molecules, or their mobilization to form nucleic acid molecules, it is possible to inhibit RNA and DNA synthesis; by inhibiting this activity in a fashion targeted to cancerous cells, the ability of tumor cells to divide and replicate may be inhibited. Additionally, there are nucleotides which do not form nucleic acid molecules, but rather serve as energy stores (i.e., AMP) or as coenzymes (i.e., FAD and NAD).

Several publications have described the use of these chemicals for these medical indications, by influencing purine and/or pyrimidine metabolism (e.g. Christopherson, R.I. and Lyons, S.D. (1990) "Potent inhibitors of *de novo* pyrimidine and purine biosynthesis as chemotherapeutic agents." *Med. Res. Reviews* 10: 505-548). Studies of enzymes involved in purine and pyrimidine metabolism have been focused on the development of new drugs which can be used, for example, as immunosuppressants or anti-proliferants (Smith, J.L., (1995) "Enzymes in nucleotide synthesis." *Curr. Opin. Struct. Biol.* 5: 752-757; (1995) *Biochem Soc. Transact.* 23: 877-902). However, purine and pyrimidine bases, nucleosides and nucleotides have other utilities: as intermediates in the biosynthesis of several fine chemicals (e.g., thiamine, S-adenosyl-methionine, folates, or riboflavin), as energy carriers for the cell (e.g., ATP or GTP), and for chemicals themselves, commonly used as flavor enhancers (e.g., IMP or GMP) or for several medicinal applications (see, for example, Kuninaka, A. (1996) Nucleotides and Related Compounds in Biotechnology vol. 6, Rehm *et al.*, eds. VCH: Weinheim, p. 561-612). Also, enzymes involved in purine, pyrimidine, nucleoside, or nucleotide metabolism are increasingly serving as targets against which chemicals for crop protection, including fungicides, herbicides and insecticides, are developed.

The metabolism of these compounds in bacteria has been characterized (for reviews see, for example, Zalkin, H. and Dixon, J.E. (1992) "*de novo* purine nucleotide biosynthesis", in: *Progress in Nucleic Acid Research and Molecular Biology*, vol. 42, Academic Press., p. 259-287; and Michal, G. (1999) "Nucleotides and Nucleosides", Chapter 8 in: *Biochemical Pathways: An Atlas of Biochemistry and Molecular Biology*,

Wiley: New York). Purine metabolism has been the subject of intensive research, and is essential to the normal functioning of the cell. Impaired purine metabolism in higher animals can cause severe disease, such as gout. Purine nucleotides are synthesized from ribose-5-phosphate, in a series of steps through the intermediate compound inosine-5'-phosphate (IMP), resulting in the production of guanosine-5'-monophosphate (GMP) or adenosine-5'-monophosphate (AMP), from which the triphosphate forms utilized as nucleotides are readily formed. These compounds are also utilized as energy stores, so their degradation provides energy for many different biochemical processes in the cell. Pyrimidine biosynthesis proceeds by the formation of uridine-5'-monophosphate (UMP) from ribose-5-phosphate. UMP, in turn, is converted to cytidine-5'-triphosphate (CTP). The deoxy- forms of all of these nucleotides are produced in a one step reduction reaction from the diphosphate ribose form of the nucleotide to the diphosphate deoxyribose form of the nucleotide. Upon phosphorylation, these molecules are able to participate in DNA synthesis.

D. Trehalose Metabolism and Uses

Trehalose consists of two glucose molecules, bound in α -, α -1,1 linkage. It is commonly used in the food industry as a sweetener, an additive for dried or frozen foods, and in beverages. However, it also has applications in the pharmaceutical, cosmetics and biotechnology industries (see, for example, Nishimoto *et al.*, (1998) U.S. Patent No. 5,759,610; Singer, M.A. and Lindquist, S. (1998) *Trends Biotech.* 16: 460-467; Paiva, C.L.A. and Panek, A.D. (1996) *Biotech. Ann. Rev.* 2: 293-314; and Shiosaka, M. (1997) *J. Japan* 172: 97-102). Trehalose is produced by enzymes from many microorganisms and is naturally released into the surrounding medium, from which it can be collected using methods known in the art.

II. The Phosphoenolpyruvate: Sugar Phosphotransferase System

The ability of cells to grow and divide rapidly in culture is to a great degree dependent on the extent to which the cells are able to take up and utilize high energy molecules, such as glucose and other sugars. Different transporter proteins exist to transport different carbon sources into the cell. There are transport proteins for sugars, such as glucose, fructose, mannose, galactose, ribose, sorbose, ribulose, lactose, maltose, sucrose, or raffinose, and also transport proteins for starch or cellulose degradation products. Other transport systems serve to import alcohols (*e.g.*, methanol or ethanol), alkanes, fatty acids and organic acids like acetic acid or lactic acid. In bacteria, sugars may be transported into the cell across the cellular membrane by a variety of mechanisms. Aside from the symport of sugars with protons, one of the most

commonly utilized processes for sugar uptake is the bacterial phosphoenolpyruvate: sugar phosphotransferase system (PTS). This system not only catalyzes the translocation (with concomitant phosphorylation) of sugars and hexitols, but it also regulates cellular metabolism in response to the availability of carbohydrates. Such PTS

- 5 systems are ubiquitous in bacteria but do not occur in archaeobacteria or eukaryotes.

- Functionally, the PTS system consists of two cytoplasmic proteins, Enzyme I and HPr, and a variable number of sugar-specific integral and peripheral membrane transport complexes (each termed 'Enzyme II' with a sugar-specific subscript, e.g., 'Enzyme II^{Glc}' for the Enzyme II complex which binds glucose). Enzymes II specific
- 10 for mono-, di-, or oligosaccharides, like glucose, fructose, mannose, galactose, ribose, sorbose, ribulose, lactose, maltose, sucrose, raffinose, and others are known. Enzyme I transfers phosphoryl groups from phosphoenolpyruvate (PEP) to the phosphoryl carrier protein, HPr. HPr then transfers the phosphoryl groups to the different Enzyme II transport complexes. While the amino acid sequences of Enzyme I and HPr are quite
- 15 similar in all bacteria, the sequences for PTS transporters can be grouped into structurally unrelated families. Further, the number and homology between these genes vary from bacteria to bacteria. The *E. coli* genome encodes 38 different PTS proteins, 33 of which are subunits belonging to 22 different transporters. The *M. genitalium* genome contains one gene each for Enzyme I and HPr, and only two genes for PTS
- 20 transporters. The genomes of *T. palladium* and *C. trachomatis* contain genes for Enzyme I- and HPr-like proteins but no PTS transporters.

- All PTS transporters consist of three functional units, IIA, IIB, and IIC, which occur either as protein subunits in a complex (e.g., IIA^{Glc}IICB^{Glc}) or as domains of a single polypeptide chain (e.g., IICBA^{GlcNAc}). IIA and IIB sequentially transfer
- 25 phosphoryl groups from HPr to the transported sugars. IIC contains the sugar binding site, and spans the inner membrane six or eight times. Sugar translocation is coupled to the transient phosphorylation of the IIB domain. Enzyme I, HPr, and IIA are phosphorylated at histidine residues, while IIB subunits are phosphorylated at either cysteine or histidine residues, depending on the particular transporter involved.
- 30 Phosphorylation of the sugar being imported has the advantage of blocking the diffusion of the sugar back through the cellular membrane to the extracellular medium, since the charged phosphate group cannot readily traverse the hydrophobic core of the membrane.

- Some PTS proteins play a role in intracellular signal transduction in addition to their function in the active transport of sugars. These subunits regulate their targets
- 35 either allosterically, or by phosphorylation. Their regulatory activity varies with the degree of their phosphorylation (i.e., the ratio of the non-phosphorylated to the phosphorylated form), which in turn varies with the ratio of sugar-dependent

09604231.0007.00

- dephosphorylation and phosphoenolpyruvate-dependent rephosphorylation. Examples of such intracellular regulation by PTS proteins in *E. coli* include the inhibition of glycerol kinase by dephosphorylated IIA^{Glc}, and the activation of adenylate cyclase by the phosphorylated version of this protein. Also, the HPr and the IIB domains of some
- 5 transporters in these microorganisms regulate gene expression by reversible phosphorylation of transcription antiterminators. In gram-positive bacteria, the activity of HPr is modulated by HPr-specific serine kinases and phosphatases. For example, HPr phosphorylated at serine-46 functions as a co-repressor of the transcriptional repressor CcpA. Lastly, it has been found that unphosphorylated Enzyme I inhibits the sensor
- 10 kinase CheA of the bacterial chemotaxis machinery, providing a direct link between the sugar binding and transport systems of the bacterium and those systems governing movement of the bacterium (Sonenshein, A. L., *et al.*, eds. *Bacillus subtilis* and other gram-positive bacteria. ASM: Washington, D.C.; Neidhardt, F.C., *et al.*, eds. (1996) *Escherichia coli* and *Salmonella*. ASM Press: Washington, D.C.; Lengeler *et al.*, (1999).
- 15 Biology of Prokaryotes. Section II, pp. 68-87, Thieme Verlag: Stuttgart).

III. Elements and Methods of the Invention

- The present invention is based, at least in part, on the discovery of novel molecules, referred to herein as PTS nucleic acid and protein molecules, which
- 20 participate in the uptake of high-energy carbon molecules (*e.g.*, glucose, sucrose, and fructose) into *C. glutamicum*, and may also participate in one or more intracellular signal transduction pathways in these microorganisms. In one embodiment, the PTS molecules function to import high-energy carbon molecules into the cell, where the energy produced by their degradation may be utilized to power less energetically favorable
- 25 biochemical reactions, and their degradation products may serve as intermediates and precursors for a number of other metabolic pathways. In another embodiment, the PTS molecules may participate in one or more intracellular signal transduction pathways, wherein the presence of a modified form of a PTS molecule (*e.g.*, a phosphorylated PTS protein) may participate in a signal transduction cascade which regulates one or more
- 30 cellular processes. In a preferred embodiment, the activity of the PTS molecules of the present invention has an impact on the production of a desired fine chemical by this organism. In a particularly preferred embodiment, the PTS molecules of the invention are modulated in activity, such that the yield, production or efficiency of production of one or more fine chemicals from *C. glutamicum* is also modulated.

- 35 The language, "PTS protein" or "PTS polypeptide" includes proteins which participate in the uptake of one or more high-energy carbon compounds (*e.g.*, mono-, di, or oligosaccharides, such as fructose, mannose, sucrose, glucose, raffinose, galactose,

03604231, 0562700

- ribose, lactose, maltose, and ribulose) from the extracellular medium to the interior of the cell. Such PTS proteins may also participate in one or more intracellular signal transduction pathways, such as, but not limited to, those governing the uptake of different sugars into the cell. Examples of PTS proteins include those encoded by the
- 5 PTS genes set forth in Table 1 and Appendix A. For general references pertaining to the PTS system, see: Stryer, L. (1988) *Biochemistry*. Chapter 37: "Membrane Transport", W.H. Freeman: New York, p. 959-961; Darnell, J. *et al.* (1990) *Molecular Cell Biology* Scientific American Books: New York, p. 552-553, and Michal, G., ed. (1999) *Biochemical Pathways: An Atlas of Biochemistry and Molecular Biology*, Chapter 15
- 10 "Special Bacterial Metabolism". The terms "PTS gene" or "PTS nucleic acid sequence" include nucleic acid sequences encoding a PTS protein, which consist of a coding region and also corresponding untranslated 5' and 3' sequence regions. Examples of PTS genes include those set forth in Table 1. The terms "production" or "productivity" are art-
- 15 recognized and include the concentration of the fermentation product (for example, the desired fine chemical) formed within a given time and a given fermentation volume (e.g., kg product per hour per liter). The term "efficiency of production" includes the time required for a particular level of production to be achieved (for example, how long it takes for the cell to attain a particular rate of output of a fine chemical). The term "yield" or "product/carbon yield" is art-recognized and includes the efficiency of the
- 20 conversion of the carbon source into the product (i.e., fine chemical). This is generally written as, for example, kg product per kg carbon source. By increasing the yield or production of the compound, the quantity of recovered molecules, or of useful recovered molecules of that compound in a given amount of culture over a given amount of time is increased. The terms "biosynthesis" or a "biosynthetic pathway" are art-recognized and
- 25 include the synthesis of a compound, preferably an organic compound, by a cell from intermediate compounds in what may be a multistep and highly regulated process. The terms "degradation" or a "degradation pathway" are art-recognized and include the breakdown of a compound, preferably an organic compound, by a cell to degradation products (generally speaking, smaller or less complex molecules) in what may be a
- 30 multistep and highly regulated process. The language "metabolism" is art-recognized and includes the totality of the biochemical reactions that take place in an organism. The metabolism of a particular compound, then, (e.g., the metabolism of an amino acid such as glycine) comprises the overall biosynthetic, modification, and degradation pathways in the cell related to this compound. The language "transport" or "import" is
- 35 art-recognized and includes the facilitated movement of one or more molecules across a cellular membrane through which the molecule would otherwise be unable to pass.

03604231.062700

In another embodiment, the PTS molecules of the invention are capable of modulating the production of a desired molecule, such as a fine chemical, in a microorganism such as *C. glutamicum*. Using recombinant genetic techniques, one or more of the PTS proteins of the invention may be manipulated such that its function is modulated. For example, a protein involved in the PTS-mediated import of glucose may be altered such that it is optimized in activity, and the PTS system for the importation of glucose may thus be able to translocate increased amounts of glucose into the cell. Since glucose molecules are utilized not only for energy to drive energetically unfavorable biochemical reactions, such as fine chemical biosyntheses, but also as precursors and intermediates in a number of fine chemical biosynthetic pathways (e.g., serine is synthesized from 3-phosphoglycerate). In each case, the overall yield or rate of production of one of these desired fine chemicals may be increased, either by increasing the energy available for such production to occur, or by increasing the availability of compounds necessary for such production to take place.

Further, many PTS proteins are known to play key roles in intracellular signal transduction pathways which regulate cellular metabolism and sugar uptake in keeping with the availability of carbon sources. For example, it is known that an increased intracellular level of fructose 1,6-bisphosphate (a compound produced during glycolysis) results in the phosphorylation of a serine residue on HPr which prevents this protein from serving as a phosphoryl donor in any PTS sugar transport process, thereby blocking further sugar uptake. By mutagenizing HPr such that this serine residue cannot be phosphorylated, one may constitutively activate HPr and thereby increase sugar transport into the cell, which in turn will ensure greater intracellular energy stores and intermediate/precursor molecules for the biosynthesis of one or more desired fine chemicals.

The isolated nucleic acid sequences of the invention are contained within the genome of a *Corynebacterium glutamicum* strain available through the American Type Culture Collection, given designation ATCC 13032. The nucleotide sequence of the isolated *C. glutamicum* PTS DNAs and the predicted amino acid sequences of the *C. glutamicum* PTS proteins are shown in Appendices A and B, respectively. Computational analyses were performed which classified and/or identified these nucleotide sequences as sequences which encode metabolic pathway proteins.

The present invention also pertains to proteins which have an amino acid sequence which is substantially homologous to an amino acid sequence of Appendix B. As used herein, a protein which has an amino acid sequence which is substantially homologous to a selected amino acid sequence is least about 50% homologous to the selected amino acid sequence, e.g., the entire selected amino acid sequence. A protein

which has an amino acid sequence which is substantially homologous to a selected amino acid sequence can also be least about 50-60%, preferably at least about 60-70%, and more preferably at least about 70-80%, 80-90%, or 90-95%, and most preferably at least about 96%, 97%, 98%, 99% or more homologous to the selected amino acid

5 sequence.

The PTS protein or a biologically active portion or fragment thereof of the invention can participate in the transport of high-energy carbon-containing molecules such as glucose into *C. glutamicum*, or can participate in intracellular signal transduction in this microorganism, or may have one or more of the activities set forth in Table 1.

10 Various aspects of the invention are described in further detail in the following subsections:

A. Isolated Nucleic Acid Molecules

One aspect of the invention pertains to isolated nucleic acid molecules that
15 encode PTS polypeptides or biologically active portions thereof, as well as nucleic acid fragments sufficient for use as hybridization probes or primers for the identification or amplification of PTS-encoding nucleic acid (e.g., PTS DNA). As used herein, the term "nucleic acid molecule" is intended to include DNA molecules (e.g., cDNA or genomic DNA) and RNA molecules (e.g., mRNA) and analogs of the DNA or RNA generated
20 using nucleotide analogs. This term also encompasses untranslated sequence located at both the 3' and 5' ends of the coding region of the gene: at least about 100 nucleotides of sequence upstream from the 5' end of the coding region and at least about 20 nucleotides of sequence downstream from the 3' end of the coding region of the gene. The nucleic acid molecule can be single-stranded or double-stranded, but preferably is
25 double-stranded DNA. An "isolated" nucleic acid molecule is one which is separated from other nucleic acid molecules which are present in the natural source of the nucleic acid. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank the nucleic acid (i.e., sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in
30 various embodiments, the isolated PTS nucleic acid molecule can contain less than about 5 kb, 4kb, 3kb, 2kb, 1 kb, 0.5 kb or 0.1 kb of nucleotide sequences which naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived (e.g., a *C. glutamicum* cell). Moreover, an "isolated" nucleic acid molecule, such as a DNA molecule, can be substantially free of other cellular material,
35 or culture medium when produced by recombinant techniques, or chemical precursors or other chemicals when chemically synthesized.

09604231.062700

09604231.062700

A nucleic acid molecule of the present invention, *e.g.*, a nucleic acid molecule having a nucleotide sequence of Appendix A, or a portion thereof, can be isolated using standard molecular biology techniques and the sequence information provided herein. For example, a *C. glutamicum* PTS DNA can be isolated from a *C. glutamicum* library

5 using all or portion of one of the sequences of Appendix A as a hybridization probe and standard hybridization techniques (*e.g.*, as described in Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual*. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989).

10 Moreover, a nucleic acid molecule encompassing all or a portion of one of the sequences of Appendix A can be isolated by the polymerase chain reaction using oligonucleotide primers designed based upon this sequence (*e.g.*, a nucleic acid molecule encompassing all or a portion of one of the sequences of Appendix A can be isolated by the polymerase chain reaction using oligonucleotide primers designed based upon this same sequence of Appendix A). For example, mRNA can be isolated from normal endothelial cells (*e.g.*,

15 by the guanidinium-thiocyanate extraction procedure of Chirgwin *et al.* (1979) *Biochemistry* 18: 5294-5299) and DNA can be prepared using reverse transcriptase (*e.g.*, Moloney MLV reverse transcriptase, available from Gibco/BRL, Bethesda, MD; or AMV reverse transcriptase, available from Seikagaku America, Inc., St. Petersburg, FL). Synthetic oligonucleotide primers for polymerase chain reaction amplification can be

20 designed based upon one of the nucleotide sequences shown in Appendix A. A nucleic acid of the invention can be amplified using cDNA or, alternatively, genomic DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques. The nucleic acid so amplified can be cloned into an appropriate vector and characterized by DNA sequence analysis. Furthermore,

25 oligonucleotides corresponding to a PTS nucleotide sequence can be prepared by standard synthetic techniques, *e.g.*, using an automated DNA synthesizer.

In a preferred embodiment, an isolated nucleic acid molecule of the invention comprises one of the nucleotide sequences shown in Appendix A. The sequences of Appendix A correspond to the *Corynebacterium glutamicum* PTS DNAs of the

30 invention. This DNA comprises sequences encoding PTS proteins (*i.e.*, the "coding region", indicated in each sequence in Appendix A), as well as 5' untranslated sequences and 3' untranslated sequences, also indicated in Appendix A. Alternatively, the nucleic acid molecule can comprise only the coding region of any of the sequences in Appendix A.

35 For the purposes of this application, it will be understood that each of the sequences set forth in Appendix A has an identifying RXA, RXN, RXS, or RXC number having the designation "RXA", "RXN", "RXS", or "RXC" followed by 5 digits (*i.e.*,

09504231.062700

RXA01503, RXN01299, RXS00315, or RXC00953). Each of these sequences comprises up to three parts: a 5' upstream region, a coding region, and a downstream region. Each of these three regions is identified by the same RXA, RXN, RXS, or RXC designation to eliminate confusion. The recitation "one of the sequences in Appendix A", then, refers to any of the sequences in Appendix A, which may be distinguished by their differing RXA, RXN, RXS, or RXC designations. The coding region of each of these sequences is translated into a corresponding amino acid sequence, which is set forth in Appendix B. The sequences of Appendix B are identified by the same RXA, RXN, RXS, or RXC designations as Appendix A, such that they can be readily correlated. For example, the amino acid sequences in Appendix B designated RXA01503, RXN01299, RXS00315, and RXC00953 are translations of the coding regions of the nucleotide sequence of nucleic acid molecules RXA01503, RXN01299, RXS00315, and RXC00953, respectively, in Appendix A. Each of the RXA, RXN, RXS, and RXC nucleotide and amino acid sequences of the invention has also been assigned a SEQ ID NO, as indicated in Table 1. For example, as set forth in Table 1, the nucleotide sequence of RXN01299 is SEQ ID NO: 7, and the corresponding amino acid sequence is SEQ ID NO:8.

Several of the genes of the invention are "F-designated genes". An F-designated gene includes those genes set forth in Table 1 which have an "F" in front of the RXA, RXN, RXS, or RXC designation. For example, SEQ ID NO:3, designated, as indicated on Table 1, as "F RXA00315", is an F-designated gene, as are SEQ ID NOs: 9, 11, and 13 (designated on Table 1 as "F RXA01299", "F RXA01883", and "F RXA01889", respectively).

In one embodiment, the nucleic acid molecules of the present invention are not intended to include *C. glutamicum* those compiled in Table 2. In the case of the dapD gene, a sequence for this gene was published in Wehrmann, A., *et al.* (1998) *J. Bacteriol.* 180(12): 3159-3165. However, the sequence obtained by the inventors of the present application is significantly longer than the published version. It is believed that the published version relied on an incorrect start codon, and thus represents only a fragment of the actual coding region.

In another preferred embodiment, an isolated nucleic acid molecule of the invention comprises a nucleic acid molecule which is a complement of one of the nucleotide sequences shown in Appendix A, or a portion thereof. A nucleic acid molecule which is complementary to one of the nucleotide sequences shown in Appendix A is one which is sufficiently complementary to one of the nucleotide sequences shown in Appendix A such that it can hybridize to one of the nucleotide sequences shown in Appendix A, thereby forming a stable duplex.

In still another preferred embodiment, an isolated nucleic acid molecule of the invention comprises a nucleotide sequence which is at least about 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, or 60%, preferably at least about 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, or 70%, more preferably at least about 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, or 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, or 90%, or 91%, 92%, 93%, 94%, and even more preferably at least about 95%, 96%, 97%, 98%, 99% or more homologous to a nucleotide sequence shown in Appendix A, or a portion thereof. Ranges and identity values intermediate to the above-recited ranges, (e.g., 70-90% identical or 80-95% identical) are also intended to be encompassed by the present invention. For example, ranges of identity values using a combination of any of the above values recited as upper and/or lower limits are intended to be included. In an additional preferred embodiment, an isolated nucleic acid molecule of the invention comprises a nucleotide sequence which hybridizes, e.g., hybridizes under stringent conditions, to one of the nucleotide sequences shown in Appendix A, or a portion thereof.

Moreover, the nucleic acid molecule of the invention can comprise only a portion of the coding region of one of the sequences in Appendix A, for example a fragment which can be used as a probe or primer or a fragment encoding a biologically active portion of a PTS protein. The nucleotide sequences determined from the cloning of the PTS genes from *C. glutamicum* allows for the generation of probes and primers designed for use in identifying and/or cloning PTS homologues in other cell types and organisms, as well as PTS homologues from other *Corynebacteria* or related species. The probe/primer typically comprises substantially purified oligonucleotide. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 12, preferably about 25, more preferably about 40, 50 or 75 consecutive nucleotides of a sense strand of one of the sequences set forth in Appendix A, an anti-sense sequence of one of the sequences set forth in Appendix A, or naturally occurring mutants thereof. Primers based on a nucleotide sequence of Appendix A can be used in PCR reactions to clone PTS homologues. Probes based on the PTS nucleotide sequences can be used to detect transcripts or genomic sequences encoding the same or homologous proteins. In preferred embodiments, the probe further comprises a label group attached thereto, e.g. the label group can be a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used as a part of a diagnostic test kit for identifying cells which misexpress a PTS protein, such as by measuring a level of a PTS-encoding nucleic acid in a sample of cells e.g., detecting PTS mRNA levels or determining whether a genomic PTS gene has been mutated or deleted.

06604231.062700

In one embodiment, the nucleic acid molecule of the invention encodes a protein or portion thereof which includes an amino acid sequence which is sufficiently homologous to an amino acid sequence of Appendix B such that the protein or portion thereof maintains the ability to participate in the transport of high-energy carbon molecules (such as glucose) into *C. glutamicum*, and may also participate in one or more intracellular signal transduction pathways. As used herein, the language "sufficiently homologous" refers to proteins or portions thereof which have amino acid sequences which include a minimum number of identical or equivalent (*e.g.*, an amino acid residue which has a similar side chain as an amino acid residue in one of the sequences of Appendix B) amino acid residues to an amino acid sequence of Appendix B such that the protein or portion thereof is capable of transporting high-energy carbon-containing molecules such as glucose into *C. glutamicum*, and may also participate in intracellular signal transduction in this microorganism. Protein members of such metabolic pathways, as described herein, function to transport high-energy carbon-containing molecules such as glucose into *C. glutamicum*, and may also participate in intracellular signal transduction in this microorganism. Examples of such activities are also described herein. Thus, "the function of a PTS protein" contributes to the overall functioning and/or regulation of one or more phosphoenolpyruvate-based sugar transport pathway, and /or contributes, either directly or indirectly, to the yield, production, and/or efficiency of production of one or more fine chemicals. Examples of PTS protein activities are set forth in Table 1.

In another embodiment, the protein is at least about 50-60%, preferably at least about 60-70%, and more preferably at least about 70-80%, 80-90%, 90-95%, and most preferably at least about 96%, 97%, 98%, 99% or more homologous to an entire amino acid sequence of Appendix B.

Portions of proteins encoded by the PTS nucleic acid molecules of the invention are preferably biologically active portions of one of the PTS proteins. As used herein, the term "biologically active portion of a PTS protein" is intended to include a portion, *e.g.*, a domain/motif, of a PTS protein that is capable of transporting high-energy carbon-containing molecules such as glucose into *C. glutamicum*, or of participating in intracellular signal transduction in this microorganism, or has an activity as set forth in Table 1. To determine whether a PTS protein or a biologically active portion thereof can participate in the transportation of high-energy carbon-containing molecules such as glucose into *C. glutamicum*, or can participate in intracellular signal transduction in this microorganism, an assay of enzymatic activity may be performed. Such assay methods are well known to those of ordinary skill in the art, as detailed in Example 8 of the Exemplification.

Additional nucleic acid fragments encoding biologically active portions of a PTS protein can be prepared by isolating a portion of one of the sequences in Appendix B, expressing the encoded portion of the PTS protein or peptide (*e.g.*, by recombinant expression *in vitro*) and assessing the activity of the encoded portion of the PTS protein or peptide.

The invention further encompasses nucleic acid molecules that differ from one of the nucleotide sequences shown in Appendix A (and portions thereof) due to degeneracy of the genetic code and thus encode the same PTS protein as that encoded by the nucleotide sequences shown in Appendix A. In another embodiment, an isolated nucleic acid molecule of the invention has a nucleotide sequence encoding a protein having an amino acid sequence shown in Appendix B. In a still further embodiment, the nucleic acid molecule of the invention encodes a full length *C. glutamicum* protein which is substantially homologous to an amino acid sequence of Appendix B (encoded by an open reading frame shown in Appendix A).

It will be understood by one of ordinary skill in the art that in one embodiment the sequences of the invention are not meant to include the sequences of the prior art, such as those Genbank sequences set forth in Tables 2 or 4 which were available prior to the present invention. In one embodiment, the invention includes nucleotide and amino acid sequences having a percent identity to a nucleotide or amino acid sequence of the invention which is greater than that of a sequence of the prior art (*e.g.*, a Genbank sequence (or the protein encoded by such a sequence) set forth in Tables 2 or 4). For example, the invention includes a nucleotide sequence which is greater than and/or at least 44% identical to the nucleotide sequence designated RXA01503 (SEQ ID NO:5), a nucleotide sequence which is greater than and/or at least 41% identical to the nucleotide sequence designated RXA00951 (SEQ ID NO:15), and a nucleotide sequence which is greater than and/or at least 38% identical to the nucleotide sequence designated RXA01300 (SEQ ID NO:21). One of ordinary skill in the art would be able to calculate the lower threshold of percent identity for any given sequence of the invention by examining the GAP-calculated percent identity scores set forth in Table 4 for each of the three top hits for the given sequence, and by subtracting the highest GAP-calculated percent identity from 100 percent. One of ordinary skill in the art will also appreciate that nucleic acid and amino acid sequences having percent identities greater than the lower threshold so calculated (*e.g.*, at least 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, or 60%, preferably at least about 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, or 70%, more preferably at least about 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, or 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, or

90%, or 91%, 92%, 93%, 94%, and even more preferably at least about 95%, 96%, 97%, 98%, 99% or more identical) are also encompassed by the invention.

In addition to the *C. glutamicum* PTS nucleotide sequences shown in Appendix A, it will be appreciated by those of ordinary skill in the art that DNA sequence polymorphisms that lead to changes in the amino acid sequences of PTS proteins may exist within a population (e.g., the *C. glutamicum* population). Such genetic polymorphism in the PTS gene may exist among individuals within a population due to natural variation. As used herein, the terms "gene" and "recombinant gene" refer to nucleic acid molecules comprising an open reading frame encoding a PTS protein, preferably a *C. glutamicum* PTS protein. Such natural variations can typically result in 1-5% variance in the nucleotide sequence of the PTS gene. Any and all such nucleotide variations and resulting amino acid polymorphisms in PTS that are the result of natural variation and that do not alter the functional activity of PTS proteins are intended to be within the scope of the invention.

Nucleic acid molecules corresponding to natural variants and non-*C. glutamicum* homologues of the *C. glutamicum* PTS DNA of the invention can be isolated based on their homology to the *C. glutamicum* PTS nucleic acid disclosed herein using the *C. glutamicum* DNA, or a portion thereof, as a hybridization probe according to standard hybridization techniques under stringent hybridization conditions. Accordingly, in another embodiment, an isolated nucleic acid molecule of the invention is at least 15 nucleotides in length and hybridizes under stringent conditions to the nucleic acid molecule comprising a nucleotide sequence of Appendix A. In other embodiments, the nucleic acid is at least 30, 50, 100, 250 or more nucleotides in length. As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences at least 60% homologous to each other typically remain hybridized to each other. Preferably, the conditions are such that sequences at least about 65%, more preferably at least about 70%, and even more preferably at least about 75% or more homologous to each other typically remain hybridized to each other. Such stringent conditions are known to those of ordinary skill in the art and can be found in Ausubel *et al.*, *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. A preferred, non-limiting example of stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50-65°C. Preferably, an isolated nucleic acid molecule of the invention that hybridizes under stringent conditions to a sequence of Appendix A corresponds to a naturally-occurring nucleic acid molecule. As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence

09604231.052700

that occurs in nature (e.g., encodes a natural protein). In one embodiment, the nucleic acid encodes a natural *C. glutamicum* PTS protein.

In addition to naturally-occurring variants of the PTS sequence that may exist in the population, one of ordinary skill in the art will further appreciate that changes can be introduced by mutation into a nucleotide sequence of Appendix A, thereby leading to changes in the amino acid sequence of the encoded PTS protein, without altering the functional ability of the PTS protein. For example, nucleotide substitutions leading to amino acid substitutions at "non-essential" amino acid residues can be made in a sequence of Appendix A. A "non-essential" amino acid residue is a residue that can be altered from the wild-type sequence of one of the PTS proteins (Appendix B) without altering the activity of said PTS protein, whereas an "essential" amino acid residue is required for PTS protein activity. Other amino acid residues, however, (e.g., those that are not conserved or only semi-conserved in the domain having PTS activity) may not be essential for activity and thus are likely to be amenable to alteration without altering PTS activity.

Accordingly, another aspect of the invention pertains to nucleic acid molecules encoding PTS proteins that contain changes in amino acid residues that are not essential for PTS activity. Such PTS proteins differ in amino acid sequence from a sequence contained in Appendix B yet retain at least one of the PTS activities described herein. In one embodiment, the isolated nucleic acid molecule comprises a nucleotide sequence encoding a protein, wherein the protein comprises an amino acid sequence at least about 50% homologous to an amino acid sequence of Appendix B and is capable of transporting high-energy carbon-containing molecules such as glucose into *C. glutamicum*, or of participating in intracellular signal transduction in this microorganism, or has one or more activities set forth in Table 1. Preferably, the protein encoded by the nucleic acid molecule is at least about 50-60% homologous to one of the sequences in Appendix B, more preferably at least about 60-70% homologous to one of the sequences in Appendix B, even more preferably at least about 70-80%, 80-90%, 90-95% homologous to one of the sequences in Appendix B, and most preferably at least about 96%, 97%, 98%, or 99% homologous to one of the sequences in Appendix B.

To determine the percent homology of two amino acid sequences (e.g., one of the sequences of Appendix B and a mutant form thereof) or of two nucleic acids, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in the sequence of one protein or nucleic acid for optimal alignment with the other protein or nucleic acid). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in one sequence (e.g., one of the sequences of Appendix B) is occupied by the same amino acid residue

or nucleotide as the corresponding position in the other sequence (e.g., a mutant form of the sequence selected from Appendix B), then the molecules are homologous at that position (i.e., as used herein amino acid or nucleic acid "homology" is equivalent to amino acid or nucleic acid "identity"). The percent homology between the two sequences is a function of the number of identical positions shared by the sequences (i.e., % homology = # of identical positions/total # of positions x 100).

An isolated nucleic acid molecule encoding a PTS protein homologous to a protein sequence of Appendix B can be created by introducing one or more nucleotide substitutions, additions or deletions into a nucleotide sequence of Appendix A such that one or more amino acid substitutions, additions or deletions are introduced into the encoded protein. Mutations can be introduced into one of the sequences of Appendix A by standard techniques, such as site-directed mutagenesis and PCR-mediated mutagenesis. Preferably, conservative amino acid substitutions are made at one or more predicted non-essential amino acid residues. A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). Thus, a predicted nonessential amino acid residue in a PTS protein is preferably replaced with another amino acid residue from the same side chain family. Alternatively, in another embodiment, mutations can be introduced randomly along all or part of a PTS coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for a PTS activity described herein to identify mutants that retain PTS activity. Following mutagenesis of one of the sequences of Appendix A, the encoded protein can be expressed recombinantly and the activity of the protein can be determined using, for example, assays described herein (see Example 8 of the Exemplification).

In addition to the nucleic acid molecules encoding PTS proteins described above, another aspect of the invention pertains to isolated nucleic acid molecules which are antisense thereto. An "antisense" nucleic acid comprises a nucleotide sequence which is complementary to a "sense" nucleic acid encoding a protein, e.g., complementary to the coding strand of a double-stranded DNA molecule or complementary to an mRNA sequence. Accordingly, an antisense nucleic acid can hydrogen bond to a sense nucleic

10

35

amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (i.e., RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

The antisense nucleic acid molecules of the invention are typically administered to a cell or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding a PTS protein to thereby inhibit expression of the protein, e.g., by inhibiting transcription and/or translation. The hybridization can be by conventional nucleotide complementarity to form a stable duplex, or, for example, in the case of an antisense nucleic acid molecule which binds to DNA duplexes, through specific interactions in the major groove of the double helix. The antisense molecule can be modified such that it specifically binds to a receptor or an antigen expressed on a selected cell surface, e.g., by linking the antisense nucleic acid molecule to a peptide or an antibody which binds to a cell surface receptor or antigen. The antisense nucleic acid molecule can also be delivered to cells using the vectors described herein. To achieve sufficient intracellular concentrations of the antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong prokaryotic, viral, or eukaryotic promoter are preferred.

In yet another embodiment, the antisense nucleic acid molecule of the invention is an α -anomeric nucleic acid molecule. An α -anomeric nucleic acid molecule forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gaultier *et al.* (1987) *Nucleic Acids Res.* 15:6625-6641). The antisense nucleic acid molecule can also comprise a 2'-o-methylribonucleotide (Inoue *et al.* (1987) *Nucleic Acids Res.* 15:6131-6148) or a chimeric RNA-DNA analogue (Inoue *et al.* (1987) *FEBS Lett.* 215:327-330).

In still another embodiment, an antisense nucleic acid of the invention is a ribozyme. Ribozymes are catalytic RNA molecules with ribonuclease activity which are capable of cleaving a single-stranded nucleic acid, such as an mRNA, to which they have a complementary region. Thus, ribozymes (e.g., hammerhead ribozymes (described in Haselhoff and Gerlach (1988) *Nature* 334:585-591)) can be used to catalytically cleave PTS mRNA transcripts to thereby inhibit translation of PTS mRNA. A ribozyme having specificity for a PTS-encoding nucleic acid can be designed based upon the nucleotide sequence of a PTS DNA disclosed herein (i.e., SEQ ID NO:5 (RXA01503 in Appendix A)). For example, a derivative of a *Tetrahymena* L-19 IVS RNA can be constructed in which the nucleotide sequence of the active site is complementary to the nucleotide sequence to be cleaved in a PTS-encoding mRNA.

See, e.g., Cech *et al.* U.S. Patent No. 4,987,071 and Cech *et al.* U.S. Patent No. 5,116,742. Alternatively, PTS mRNA can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules. See, e.g., Bartel, D. and Szostak, J.W. (1993) *Science* 261:1411-1418.

- 5 Alternatively, PTS gene expression can be inhibited by targeting nucleotide sequences complementary to the regulatory region of a PTS nucleotide sequence (e.g., a PTS promoter and/or enhancers) to form triple helical structures that prevent transcription of a PTS gene in target cells. See generally, Helene, C. (1991) *Anticancer Drug Des.* 6(6):569-84; Helene, C. *et al.* (1992) *Ann. N.Y. Acad. Sci.* 660:27-36; and
- 10 Maher, L.J. (1992) *Bioassays* 14(12):807-15.

B. Recombinant Expression Vectors and Host Cells

- Another aspect of the invention pertains to vectors, preferably expression vectors, containing a nucleic acid encoding a PTS protein (or a portion thereof). As
- 15 used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments can be ligated. Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome. Certain vectors are capable of
- 20 autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to
- 25 which they are operatively linked. Such vectors are referred to herein as "expression vectors". In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, "plasmid" and "vector" can be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors,
- 30 such as viral vectors (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

- The recombinant expression vectors of the invention comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell, which means that the recombinant expression vectors include one or more regulatory
- 35 sequences, selected on the basis of the host cells to be used for expression, which is operatively linked to the nucleic acid sequence to be expressed. Within a recombinant expression vector, "operably linked" is intended to mean that the nucleotide sequence of

- interest is linked to the regulatory sequence(s) in a manner which allows for expression of the nucleotide sequence (e.g., in an *in vitro* transcription/translation system or in a host cell when the vector is introduced into the host cell). The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (e.g., polyadenylation signals). Such regulatory sequences are described, for example, in Goeddel; *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). Regulatory sequences include those which direct constitutive expression of a nucleotide sequence in many types of host cell and those which direct expression of the nucleotide sequence only in certain host cells.
- 10 Preferred regulatory sequences are, for example, promoters such as cos-, tac-, trp-, tet-, trp-tet-, lpp-, lac-, lpp-lac-, lacI^q-, T7-, T5-, T3-, gal-, trc-, ara-, SP6-, amy, SPO2, λ -P_R- or λ P_L, which are used preferably in bacteria. Additional regulatory sequences are, for example, promoters from yeasts and fungi, such as ADC1, MFa, AC, P-60, CYC1, GAPDH, TEF, rp28, ADH, promoters from plants such as CaMV/35S, SSU, OCS, lib4,
- 15 usp, STLS1, B33, nos or ubiquitin- or phaseolin-promoters. It is also possible to use artificial promoters. It will be appreciated by one of ordinary skill in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. The expression vectors of the invention can be introduced into host cells to thereby produce proteins or
- 20 peptides, including fusion proteins or peptides, encoded by nucleic acids as described herein (e.g., PTS proteins, mutant forms of PTS proteins, fusion proteins, etc.).
- The recombinant expression vectors of the invention can be designed for expression of PTS proteins in prokaryotic or eukaryotic cells. For example, PTS genes can be expressed in bacterial cells such as *C. glutamicum*, insect cells (using baculovirus
- 25 expression vectors), yeast and other fungal cells (see Romanos, M.A. *et al.* (1992) "Foreign gene expression in yeast: a review", *Yeast* 8: 423-488; van den Hondel, C.A.M.J.J. *et al.* (1991) "Heterologous gene expression in filamentous fungi" in: More Gene Manipulations in Fungi, J.W. Bennet & L.L. Lasure, eds., p. 396-428: Academic Press: San Diego; and van den Hondel, C.A.M.J.J. & Punt, P.J. (1991) "Gene transfer
- 30 systems and vector development for filamentous fungi, in: Applied Molecular Genetics of Fungi, Peberdy, J.F. *et al.*, eds., p. 1-28, Cambridge University Press: Cambridge), algae and multicellular plant cells (see Schmidt, R. and Willmitzer, L. (1988) High efficiency *Agrobacterium tumefaciens* -mediated transformation of *Arabidopsis thaliana* leaf and cotyledon explants" *Plant Cell Rep.*: 583-586), or mammalian cells.
- 35 Suitable host cells are discussed further in Goeddel, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). Alternatively, the

recombinant expression vector can be transcribed and translated *in vitro*, for example using T7 promoter regulatory sequences and T7 polymerase.

- Expression of proteins in prokaryotes is most often carried out with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion proteins. Fusion vectors add a number of amino acids to a protein encoded therein, usually to the amino terminus of the recombinant protein. Such fusion vectors typically serve three purposes: 1) to increase expression of recombinant protein; 2) to increase the solubility of the recombinant protein; and 3) to aid in the purification of the recombinant protein by acting as a ligand in affinity purification. Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the fusion moiety and the recombinant protein to enable separation of the recombinant protein from the fusion moiety subsequent to purification of the fusion protein. Such enzymes, and their cognate recognition sequences, include Factor Xa, thrombin and enterokinase.

- Typical fusion expression vectors include pGEX (Pharmacia Biotech Inc; Smith, D.B. and Johnson, K.S. (1988) *Gene* 67:31-40), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein. In one embodiment, the coding sequence of the PTS protein is cloned into a pGEX expression vector to create a vector encoding a fusion protein comprising, from the N-terminus to the C-terminus, GST-thrombin cleavage site-X protein. The fusion protein can be purified by affinity chromatography using glutathione-agarose resin. Recombinant PTS protein unfused to GST can be recovered by cleavage of the fusion protein with thrombin.

- Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amann *et al.*, (1988) *Gene* 69:301-315) pLG338, pACYC184, pBR322, pUC18, pUC19, pKC30, pRep4, pHS1, pHS2, pPLc236, pMBL24, pLG200, pUR290, pIN-III113-B1. λ gt11, pBdCl, and pET 11d (Studier *et al.*, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 60-89 ; and Pouwels *et al.*, eds. (1985) *Cloning Vectors*. Elsevier: New York ISBN 0 444 904018).
- Target gene expression from the pTrc vector relies on host RNA polymerase transcription from a hybrid trp-lac fusion promoter. Target gene expression from the pET 11d vector relies on transcription from a T7 gn10-lac fusion promoter mediated by a coexpressed viral RNA polymerase (T7 gn1). This viral polymerase is supplied by host strains BL21(DE3) or HMS174(DE3) from a resident λ prophage harboring a T7 gn1 gene under the transcriptional control of the lacUV 5 promoter. For transformation of other varieties of bacteria, appropriate vectors may be selected. For example, the plasmids pIJ101, pIJ364, pIJ702 and pIJ361 are known to be useful in transforming

Streptomyces, while plasmids pUB110, pC194, or pBD214 are suited for transformation of *Bacillus* species. Several plasmids of use in the transfer of genetic information into *Corynebacterium* include pHM1519, pBL1, pSA77, or pAJ667 (Pouwels *et al.*, eds. (1985) *Cloning Vectors*. Elsevier: New York ISBN 0 444 904018).

- 5 One strategy to maximize recombinant protein expression is to express the protein in a host bacteria with an impaired capacity to proteolytically cleave the recombinant protein (Gottesman, S., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 119-128). Another strategy is to alter the nucleic acid sequence of the nucleic acid to be inserted into an expression vector so that the individual codons for each amino acid are those preferentially utilized in the bacterium chosen for expression, such as *C. glutamicum* (Wada *et al.* (1992) *Nucleic Acids Res.* 20:2111-2118). Such alteration of nucleic acid sequences of the invention can be carried out by standard DNA synthesis techniques.

- 10 In another embodiment, the PTS protein expression vector is a yeast expression vector. Examples of vectors for expression in yeast *S. cerevisiae* include pYepSec1 (Baldari, *et al.*, (1987) *Embo J.* 6:229-234), 2 μ , pAG-1, Yep6, Yep13, pEMBLYe23, pMFa (Kurjan and Herskowitz, (1982) *Cell* 30:933-943), pJRY88 (Schultz *et al.*, (1987) *Gene* 54:113-123), and pYES2 (Invitrogen Corporation, San Diego, CA). Vectors and methods for the construction of vectors appropriate for use in other fungi, such as the filamentous fungi, include those detailed in: van den Hondel, C.A.M.J.J. & Punt, P.J. (1991) "Gene transfer systems and vector development for filamentous fungi, in: Applied Molecular Genetics of Fungi, J.F. Peberdy, *et al.*, eds., p. 1-28, Cambridge University Press: Cambridge, and Pouwels *et al.*, eds. (1985) *Cloning Vectors*. Elsevier: New York (ISBN 0 444 904018).

- 25 Alternatively, the PTS proteins of the invention can be expressed in insect cells using baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (*e.g.*, Sf 9 cells) include the pAc series (Smith *et al.* (1983) *Mol. Cell Biol.* 3:2156-2165) and the pVL series (Lucklow and Summers (1989) *Virology* 170:31-39).

- 30 In another embodiment, the PTS proteins of the invention may be expressed in unicellular plant cells (such as algae) or in plant cells from higher plants (*e.g.*, the spermatophytes, such as crop plants). Examples of plant expression vectors include those detailed in: Becker, D., Kemper, E., Schell, J. and Masterson, R. (1992) "New plant binary vectors with selectable markers located proximal to the left border", *Plant Mol. Biol.* 20: 1195-1197; and Bevan, M.W. (1984) "Binary *Agrobacterium* vectors for plant transformation", *Nucl. Acid. Res.* 12: 8711-8721, and include pLGV23, pGHlac+,

09674231.062700

pBIN19, pAK2004, and pDH51 (Pouwels *et al.*, eds. (1985) Cloning Vectors. Elsevier: New York ISBN 0 444 904018).

In yet another embodiment, a nucleic acid of the invention is expressed in mammalian cells using a mammalian expression vector. Examples of mammalian expression vectors include pCDM8 (Seed, B. (1987) *Nature* 329:840) and pMT2PC (Kaufman *et al.* (1987) *EMBO J.* 6:187-195). When used in mammalian cells, the expression vector's control functions are often provided by viral regulatory elements. For example, commonly used promoters are derived from polyoma, Adenovirus 2, cytomegalovirus and Simian Virus 40. For other suitable expression systems for both prokaryotic and eukaryotic cells see chapters 16 and 17 of Sambrook, J., Fritsch, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual*. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989.

In another embodiment, the recombinant mammalian expression vector is capable of directing expression of the nucleic acid preferentially in a particular cell type (e.g., tissue-specific regulatory elements are used to express the nucleic acid). Tissue-specific regulatory elements are known in the art. Non-limiting examples of suitable tissue-specific promoters include the albumin promoter (liver-specific; Pinkert *et al.* (1987) *Genes Dev.* 1:268-277), lymphoid-specific promoters (Calame and Eaton (1988) *Adv. Immunol.* 43:235-275), in particular promoters of T cell receptors (Winoto and Baltimore (1989) *EMBO J.* 8:729-733) and immunoglobulins (Banerji *et al.* (1983) *Cell* 33:729-740; Queen and Baltimore (1983) *Cell* 33:741-748), neuron-specific promoters (e.g., the neurofilament promoter; Byrne and Ruddle (1989) *PNAS* 86:5473-5477), pancreas-specific promoters (Edlund *et al.* (1985) *Science* 230:912-916), and mammary gland-specific promoters (e.g., milk whey promoter; U.S. Patent No. 4,873,316 and European Application Publication No. 264,166). Developmentally-regulated promoters are also encompassed, for example the murine hox promoters (Kessel and Gruss (1990) *Science* 249:374-379) and the α -fetoprotein promoter (Campes and Tilghman (1989) *Genes Dev.* 3:537-546).

The invention further provides a recombinant expression vector comprising a DNA molecule of the invention cloned into the expression vector in an antisense orientation. That is, the DNA molecule is operatively linked to a regulatory sequence in a manner which allows for expression (by transcription of the DNA molecule) of an RNA molecule which is antisense to PTS mRNA. Regulatory sequences operatively linked to a nucleic acid cloned in the antisense orientation can be chosen which direct the continuous expression of the antisense RNA molecule in a variety of cell types, for instance viral promoters and/or enhancers, or regulatory sequences can be chosen which

03604221.062700

direct constitutive, tissue specific or cell type specific expression of antisense RNA. The antisense expression vector can be in the form of a recombinant plasmid, phagemid or attenuated virus in which antisense nucleic acids are produced under the control of a high efficiency regulatory region, the activity of which can be determined by the cell type into which the vector is introduced. For a discussion of the regulation of gene expression using antisense genes see Weintraub, H. *et al.*, Antisense RNA as a molecular tool for genetic analysis, *Reviews - Trends in Genetics*, Vol. 1(1) 1986.

Another aspect of the invention pertains to host cells into which a recombinant expression vector of the invention has been introduced. The terms "host cell" and "recombinant host cell" are used interchangeably herein. It is understood that such terms refer not only to the particular subject cell but to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

A host cell can be any prokaryotic or eukaryotic cell. For example, a PTS protein can be expressed in bacterial cells such as *C. glutamicum*, insect cells, yeast or mammalian cells (such as Chinese hamster ovary cells (CHO) or COS cells). Other suitable host cells are known to one of ordinary skill in the art. Microorganisms related to *Corynebacterium glutamicum* which may be conveniently used as host cells for the nucleic acid and protein molecules of the invention are set forth in Table 3.

Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. As used herein, the terms "transformation" and "transfection" are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid (*e.g.*, linear DNA or RNA (*e.g.*, a linearized vector or a gene construct alone without a vector) or nucleic acid in the form of a vector (*e.g.*, a plasmid, phage, phasmid, phagemid, transposon or other DNA) into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, or electroporation. Suitable methods for transforming or transfecting host cells can be found in Sambrook, *et al.* (*Molecular Cloning: A Laboratory Manual*, 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989), and other laboratory manuals.

For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (*e.g.*, resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Preferred

05604231, 0562700

- selectable markers include those which confer resistance to drugs, such as G418, hygromycin and methotrexate. Nucleic acid encoding a selectable marker can be introduced into a host cell on the same vector as that encoding a PTS protein or can be introduced on a separate vector. Cells stably transfected with the introduced nucleic acid can be identified by drug selection (e.g., cells that have incorporated the selectable marker gene will survive, while the other cells die).

- To create a homologous recombinant microorganism, a vector is prepared which contains at least a portion of a PTS gene into which a deletion, addition or substitution has been introduced to thereby alter, e.g., functionally disrupt, the PTS gene.
- 10 Preferably, this PTS gene is a *Corynebacterium glutamicum* PTS gene, but it can be a homologue from a related bacterium or even from a mammalian, yeast, or insect source. In a preferred embodiment, the vector is designed such that, upon homologous recombination, the endogenous PTS gene is functionally disrupted (i.e., no longer encodes a functional protein; also referred to as a "knock out" vector). Alternatively,
- 15 the vector can be designed such that, upon homologous recombination, the endogenous PTS gene is mutated or otherwise altered but still encodes functional protein (e.g., the upstream regulatory region can be altered to thereby alter the expression of the endogenous PTS protein). In the homologous recombination vector, the altered portion of the PTS gene is flanked at its 5' and 3' ends by additional nucleic acid of the PTS
- 20 gene to allow for homologous recombination to occur between the exogenous PTS gene carried by the vector and an endogenous PTS gene in a microorganism. The additional flanking PTS nucleic acid is of sufficient length for successful homologous recombination with the endogenous gene. Typically, several kilobases of flanking DNA (both at the 5' and 3' ends) are included in the vector (see e.g., Thomas, K.R., and
- 25 Capecchi, M.R. (1987) Cell 51: 503 for a description of homologous recombination vectors). The vector is introduced into a microorganism (e.g., by electroporation) and cells in which the introduced PTS gene has homologously recombined with the endogenous PTS gene are selected, using art-known techniques.

- In another embodiment, recombinant microorganisms can be produced which
- 30 contain selected systems which allow for regulated expression of the introduced gene. For example, inclusion of a PTS gene on a vector placing it under control of the lac operon permits expression of the PTS gene only in the presence of IPTG. Such regulatory systems are well known in the art.

- In another embodiment, an endogenous PTS gene in a host cell is disrupted (e.g.,
- 35 by homologous recombination or other genetic means known in the art) such that expression of its protein product does not occur. In another embodiment, an endogenous or introduced PTS gene in a host cell has been altered by one or more point mutations,

000004231.062700

deletions, or inversions, but still encodes a functional PTS protein. In still another embodiment, one or more of the regulatory regions (*e.g.*, a promoter, repressor, or inducer) of a PTS gene in a microorganism has been altered (*e.g.*, by deletion, truncation, inversion, or point mutation) such that the expression of the PTS gene is modulated. One of ordinary skill in the art will appreciate that host cells containing more than one of the described PTS gene and protein modifications may be readily produced using the methods of the invention, and are meant to be included in the present invention.

A host cell of the invention, such as a prokaryotic or eukaryotic host cell in culture, can be used to produce (*i.e.*, express) a PTS protein. Accordingly, the invention further provides methods for producing PTS proteins using the host cells of the invention. In one embodiment, the method comprises culturing the host cell of invention (into which a recombinant expression vector encoding a PTS protein has been introduced, or into which genome has been introduced a gene encoding a wild-type or altered PTS protein) in a suitable medium until PTS protein is produced. In another embodiment, the method further comprises isolating PTS proteins from the medium or the host cell.

C. Isolated PTS Proteins

Another aspect of the invention pertains to isolated PTS proteins, and biologically active portions thereof. An "isolated" or "purified" protein or biologically active portion thereof is substantially free of cellular material when produced by recombinant DNA techniques, or chemical precursors or other chemicals when chemically synthesized. The language "substantially free of cellular material" includes preparations of PTS protein in which the protein is separated from cellular components of the cells in which it is naturally or recombinantly produced. In one embodiment, the language "substantially free of cellular material" includes preparations of PTS protein having less than about 30% (by dry weight) of non-PTS protein (also referred to herein as a "contaminating protein"), more preferably less than about 20% of non-PTS protein, still more preferably less than about 10% of non-PTS protein, and most preferably less than about 5% non-PTS protein. When the PTS protein or biologically active portion thereof is recombinantly produced, it is also preferably substantially free of culture medium, *i.e.*, culture medium represents less than about 20%, more preferably less than about 10%, and most preferably less than about 5% of the volume of the protein preparation. The language "substantially free of chemical precursors or other chemicals" includes preparations of PTS protein in which the protein is separated from chemical precursors or other chemicals which are involved in the synthesis of the

protein. In one embodiment, the language "substantially free of chemical precursors or other chemicals" includes preparations of PTS protein having less than about 30% (by dry weight) of chemical precursors or non-PTS chemicals, more preferably less than about 20% chemical precursors or non-PTS chemicals, still more preferably less than about 10% chemical precursors or non-PTS chemicals, and most preferably less than about 5% chemical precursors or non-PTS chemicals. In preferred embodiments, isolated proteins or biologically active portions thereof lack contaminating proteins from the same organism from which the PTS protein is derived. Typically, such proteins are produced by recombinant expression of, for example, a *C. glutamicum* PTS protein in a microorganism such as *C. glutamicum*.

An isolated PTS protein or a portion thereof of the invention can participate in the transport of high-energy carbon-containing molecules such as glucose into *C. glutamicum*, and may also participate in intracellular signal transduction in this microorganism, or has one or more of the activities set forth in Table 1. In preferred embodiments, the protein or portion thereof comprises an amino acid sequence which is sufficiently homologous to an amino acid sequence of Appendix B such that the protein or portion thereof maintains the ability to transport high-energy carbon-containing molecules such as glucose into *C. glutamicum*, or to participate in intracellular signal transduction in this microorganism. The portion of the protein is preferably a biologically active portion as described herein. In another preferred embodiment, a PTS protein of the invention has an amino acid sequence shown in Appendix B. In yet another preferred embodiment, the PTS protein has an amino acid sequence which is encoded by a nucleotide sequence which hybridizes, *e.g.*, hybridizes under stringent conditions, to a nucleotide sequence of Appendix A. In still another preferred embodiment, the PTS protein has an amino acid sequence which is encoded by a nucleotide sequence that is at least about 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, or 60%, preferably at least about 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, or 70%, more preferably at least about 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, or 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, or 90%, or 91%, 92%, 93%, 94%, and even more preferably at least about 95%, 96%, 97%, 98%, 99% or more homologous to one of the nucleic acid sequences of Appendix A, or a portion thereof. Ranges and identity values intermediate to the above-recited values, (*e.g.*, 70-90% identical or 80-95% identical) are also intended to be encompassed by the present invention. For example, ranges of identity values using a combination of any of the above values recited as upper and/or lower limits are intended to be included. The preferred PTS proteins of the present invention also preferably possess at least one of the PTS activities described herein. For example, a preferred PTS protein of the present

05604731.062700

invention includes an amino acid sequence encoded by a nucleotide sequence which hybridizes, e.g., hybridizes under stringent conditions, to a nucleotide sequence of Appendix A, and which can participate in the transport of high-energy carbon-containing molecules such as glucose into *C. glutamicum*, and may also participate in intracellular signal transduction in this microorganism, or which has one or more of the activities set forth in Table 1.

In other embodiments, the PTS protein is substantially homologous to an amino acid sequence of Appendix B and retains the functional activity of the protein of one of the sequences of Appendix B yet differs in amino acid sequence due to natural variation or mutagenesis, as described in detail in subsection I above. Accordingly, in another embodiment, the PTS protein is a protein which comprises an amino acid sequence which is at least about 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, or 60%, preferably at least about 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, or 70%, more preferably at least about 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, or 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, or 90%, or 91%, 92%, 93%, 94%, and even more preferably at least about 95%, 96%, 97%, 98%, 99% or more homologous to an entire amino acid sequence of Appendix B and which has at least one of the PTS activities described herein. Ranges and identity values intermediate to the above-recited values, (e.g., 70-90% identical or 80-95% identical) are also intended to be encompassed by the present invention. For example, ranges of identity values using a combination of any of the above values recited as upper and/or lower limits are intended to be included. In another embodiment, the invention pertains to a full length *C. glutamicum* protein which is substantially homologous to an entire amino acid sequence of Appendix B.

Biologically active portions of a PTS protein include peptides comprising amino acid sequences derived from the amino acid sequence of a PTS protein, e.g., the amino acid sequence shown in Appendix B or the amino acid sequence of a protein homologous to a PTS protein, which include fewer amino acids than a full length PTS protein or the full length protein which is homologous to a PTS protein, and exhibit at least one activity of a PTS protein. Typically, biologically active portions (peptides, e.g., peptides which are, for example, 5, 10, 15, 20, 30, 35, 36, 37, 38, 39, 40, 50, 100 or more amino acids in length) comprise a domain or motif with at least one activity of a PTS protein. Moreover, other biologically active portions, in which other regions of the protein are deleted, can be prepared by recombinant techniques and evaluated for one or more of the activities described herein. Preferably, the biologically active portions of a PTS protein include one or more selected domains/motifs or portions thereof having biological activity.

PTS proteins are preferably produced by recombinant DNA techniques. For example, a nucleic acid molecule encoding the protein is cloned into an expression vector (as described above), the expression vector is introduced into a host cell (as described above) and the PTS protein is expressed in the host cell. The PTS protein can then be isolated from the cells by an appropriate purification scheme using standard protein purification techniques. Alternative to recombinant expression, a PTS protein, polypeptide, or peptide can be synthesized chemically using standard peptide synthesis techniques. Moreover, native PTS protein can be isolated from cells (e.g., endothelial cells), for example using an anti-PTS antibody, which can be produced by standard techniques utilizing a PTS protein or fragment thereof of this invention.

The invention also provides PTS chimeric or fusion proteins. As used herein, a PTS "chimeric protein" or "fusion protein" comprises a PTS polypeptide operatively linked to a non-PTS polypeptide. An "PTS polypeptide" refers to a polypeptide having an amino acid sequence corresponding to PTS, whereas a "non-PTS polypeptide" refers to a polypeptide having an amino acid sequence corresponding to a protein which is not substantially homologous to the PTS protein, e.g., a protein which is different from the PTS protein and which is derived from the same or a different organism. Within the fusion protein, the term "operatively linked" is intended to indicate that the PTS polypeptide and the non-PTS polypeptide are fused in-frame to each other. The non-PTS polypeptide can be fused to the N-terminus or C-terminus of the PTS polypeptide. For example, in one embodiment the fusion protein is a GST-PTS fusion protein in which the PTS sequences are fused to the C-terminus of the GST sequences. Such fusion proteins can facilitate the purification of recombinant PTS proteins. In another embodiment, the fusion protein is a PTS protein containing a heterologous signal sequence at its N-terminus. In certain host cells (e.g., mammalian host cells), expression and/or secretion of a PTS protein can be increased through use of a heterologous signal sequence.

Preferably, a PTS chimeric or fusion protein of the invention is produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different polypeptide sequences are ligated together in-frame in accordance with conventional techniques, for example by employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene

00604231-002700

fragments which can subsequently be annealed and reamplified to generate a chimeric gene sequence (see, for example, *Current Protocols in Molecular Biology*, eds. Ausubel *et al.* John Wiley & Sons: 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (e.g., a GST polypeptide). A PTS-

- 5 encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the PTS protein.

Homologues of the PTS protein can be generated by mutagenesis, e.g., discrete point mutation or truncation of the PTS protein. As used herein, the term "homologue" refers to a variant form of the PTS protein which acts as an agonist or antagonist of the activity of the PTS protein. An agonist of the PTS protein can retain substantially the same, or a subset, of the biological activities of the PTS protein. An antagonist of the PTS protein can inhibit one or more of the activities of the naturally occurring form of the PTS protein, by, for example, competitively binding to a downstream or upstream member of the PTS system which includes the PTS protein. Thus, the *C. glutamicum* 10 PTS protein and homologues thereof of the present invention may modulate the activity of one or more sugar transport pathways or intracellular signal transduction pathways in which PTS proteins play a role in this microorganism.

In an alternative embodiment, homologues of the PTS protein can be identified by screening combinatorial libraries of mutants, e.g., truncation mutants, of the PTS 20 protein for PTS protein agonist or antagonist activity. In one embodiment, a variegated library of PTS variants is generated by combinatorial mutagenesis at the nucleic acid level and is encoded by a variegated gene library. A variegated library of PTS variants can be produced by, for example, enzymatically ligating a mixture of synthetic oligonucleotides into gene sequences such that a degenerate set of potential PTS 25 sequences is expressible as individual polypeptides, or alternatively, as a set of larger fusion proteins (e.g., for phage display) containing the set of PTS sequences therein. There are a variety of methods which can be used to produce libraries of potential PTS homologues from a degenerate oligonucleotide sequence. Chemical synthesis of a degenerate gene sequence can be performed in an automatic DNA synthesizer, and the 30 synthetic gene then ligated into an appropriate expression vector. Use of a degenerate set of genes allows for the provision, in one mixture, of all of the sequences encoding the desired set of potential PTS sequences. Methods for synthesizing degenerate oligonucleotides are known in the art (see, e.g., Narang, S.A. (1983) *Tetrahedron* 39:3; Itakura *et al.* (1984) *Annu. Rev. Biochem.* 53:323; Itakura *et al.* (1984) *Science* 35 198:1056; Ike *et al.* (1983) *Nucleic Acid Res.* 11:477).

In addition, libraries of fragments of the PTS protein coding can be used to generate a variegated population of PTS fragments for screening and subsequent

09604231.062700

selection of homologues of a PTS protein. In one embodiment, a library of coding sequence fragments can be generated by treating a double stranded PCR fragment of a PTS coding sequence with a nuclease under conditions wherein nicking occurs only about once per molecule, denaturing the double stranded DNA, renaturing the DNA to form double stranded DNA which can include sense/antisense pairs from different nicked products, removing single stranded portions from reformed duplexes by treatment with S1 nuclease, and ligating the resulting fragment library into an expression vector. By this method, an expression library can be derived which encodes N-terminal, C-terminal and internal fragments of various sizes of the PTS protein.

Several techniques are known in the art for screening gene products of combinatorial libraries made by point mutations or truncation, and for screening cDNA libraries for gene products having a selected property. Such techniques are adaptable for rapid screening of the gene libraries generated by the combinatorial mutagenesis of PTS homologues. The most widely used techniques, which are amenable to high through-put analysis, for screening large gene libraries typically include cloning the gene library into replicable expression vectors, transforming appropriate cells with the resulting library of vectors, and expressing the combinatorial genes under conditions in which detection of a desired activity facilitates isolation of the vector encoding the gene whose product was detected. Recursive ensemble mutagenesis (REM), a new technique which enhances the frequency of functional mutants in the libraries, can be used in combination with the screening assays to identify PTS homologues (Arkin and Yourvan (1992) *PNAS* 89:7811-7815; Delgrave *et al.* (1993) *Protein Engineering* 6(3):327-331).

In another embodiment, cell based assays can be exploited to analyze a variegated PTS library, using methods well known in the art.

D. Uses and Methods of the Invention

The nucleic acid molecules, proteins, protein homologues, fusion proteins, primers, vectors, and host cells described herein can be used in one or more of the following methods: identification of *C. glutamicum* and related organisms; mapping of genomes of organisms related to *C. glutamicum*; identification and localization of *C. glutamicum* sequences of interest; evolutionary studies; determination of PTS protein regions required for function; modulation of a PTS protein activity; modulation of the activity of a PTS pathway; and modulation of cellular production of a desired compound, such as a fine chemical.

The PTS nucleic acid molecules of the invention have a variety of uses. First, they may be used to identify an organism as being *Corynebacterium glutamicum* or a close relative thereof. Also, they may be used to identify the presence of *C. glutamicum*

or a relative thereof in a mixed population of microorganisms. The invention provides the nucleic acid sequences of a number of *C. glutamicum* genes; by probing the extracted genomic DNA of a culture of a unique or mixed population of microorganisms under stringent conditions with a probe spanning a region of a *C. glutamicum* gene which is unique to this organism, one can ascertain whether this organism is present.

Although *Corynebacterium glutamicum* itself is nonpathogenic, it is related to pathogenic species, such as *Corynebacterium diphtheriae*. *Corynebacterium diphtheriae* is the causative agent of diphtheria, a rapidly developing, acute, febrile infection which involves both local and systemic pathology. In this disease, a local lesion develops in the upper respiratory tract and involves necrotic injury to epithelial cells; the bacilli secrete toxin which is disseminated through this lesion to distal susceptible tissues of the body. Degenerative changes brought about by the inhibition of protein synthesis in these tissues, which include heart, muscle, peripheral nerves, adrenals, kidneys, liver and spleen, result in the systemic pathology of the disease. Diphtheria continues to have high incidence in many parts of the world, including Africa, Asia, Eastern Europe and the independent states of the former Soviet Union. An ongoing epidemic of diphtheria in the latter two regions has resulted in at least 5,000 deaths since 1990.

In one embodiment, the invention provides a method of identifying the presence or activity of *Corynebacterium diphtheriae* in a subject. This method includes detection of one or more of the nucleic acid or amino acid sequences of the invention (e.g., the sequences set forth in Appendix A or Appendix B) in a subject, thereby detecting the presence or activity of *Corynebacterium diphtheriae* in the subject. *C. glutamicum* and *C. diphtheriae* are related bacteria, and many of the nucleic acid and protein molecules in *C. glutamicum* are homologous to *C. diphtheriae* nucleic acid and protein molecules, and can therefore be used to detect *C. diphtheriae* in a subject.

The nucleic acid and protein molecules of the invention may also serve as markers for specific regions of the genome. This has utility not only in the mapping of the genome, but also for functional studies of *C. glutamicum* proteins. For example, to identify the region of the genome to which a particular *C. glutamicum* DNA-binding protein binds, the *C. glutamicum* genome could be digested, and the fragments incubated with the DNA-binding protein. Those which bind the protein may be additionally probed with the nucleic acid molecules of the invention, preferably with readily detectable labels; binding of such a nucleic acid molecule to the genome fragment enables the localization of the fragment to the genome map of *C. glutamicum*, and, when performed multiple times with different enzymes, facilitates a rapid determination of the nucleic acid sequence to which the protein binds. Further, the nucleic acid molecules of the

invention may be sufficiently homologous to the sequences of related species such that these nucleic acid molecules may serve as markers for the construction of a genomic map in related bacteria, such as *Brevibacterium lactofermentum*.

5 The PTS nucleic acid molecules of the invention are also useful for evolutionary and protein structural studies. The sugar uptake system in which the molecules of the invention participate are utilized by a wide variety of bacteria; by comparing the sequences of the nucleic acid molecules of the present invention to those encoding similar enzymes from other organisms, the evolutionary relatedness of the organisms can be assessed. Similarly, such a comparison permits an assessment of which regions
10 of the sequence are conserved and which are not, which may aid in determining those regions of the protein which are essential for the functioning of the enzyme. This type of determination is of value for protein engineering studies and may give an indication of what the protein can tolerate in terms of mutagenesis without losing function.

15 Manipulation of the PTS nucleic acid molecules of the invention may result in the production of PTS proteins having functional differences from the wild-type PTS proteins. These proteins may be improved in efficiency or activity, may be present in greater numbers in the cell than is usual, or may be decreased in efficiency or activity.

The invention provides methods for screening molecules which modulate the activity of a PTS protein, either by interacting with the protein itself or a substrate or
20 binding partner of the PTS protein, or by modulating the transcription or translation of a PTS nucleic acid molecule of the invention. In such methods, a microorganism expressing one or more PTS proteins of the invention is contacted with one or more test compounds, and the effect of each test compound on the activity or level of expression of the PTS protein is assessed.

25 The PTS molecules of the invention may be modified such that the yield, production, and/or efficiency of production of one or more fine chemicals is improved. For example, by modifying a PTS protein involved in the uptake of glucose such that it is optimized in activity, the quantity of glucose uptake or the rate at which glucose is translocated into the cell may be increased. The breakdown of glucose and other sugars
30 within the cell provides energy that may be used to drive energetically unfavorable biochemical reactions, such as those involved in the biosynthesis of fine chemicals. This breakdown also provides intermediate and precursor molecules necessary for the biosynthesis of certain fine chemicals, such as amino acids, vitamins and cofactors. By increasing the amount of intracellular high-energy carbon molecules through
35 modification of the PTS molecules of the invention, one may therefore increase both the energy available to perform metabolic pathways necessary for the production of one or more fine chemicals, and also the intracellular pools of metabolites necessary for such

00004231 002700

production. Conversely, by decreasing the importation of a sugar whose breakdown products include a compound which is used solely in metabolic pathways which compete with pathways utilized for the production of a desired fine chemical for enzymes, cofactors, or intermediates, one may downregulate this pathway and thus perhaps increase production through the desired biosynthetic pathway.

Further, the PTS molecules of the invention may be involved in one or more intracellular signal transduction pathways which may affect the yields and/or rate of production of one or more fine chemical from *C. glutamicum*. For example, proteins necessary for the import of one or more sugars from the extracellular medium (e.g., HPr, Enzyme I, or a member of an Enzyme II complex) are frequently posttranslationally modified upon the presence of a sufficient quantity of the sugar in the cell, such that they are no longer able to import that sugar. An example of this occurs in *E. coli*, where high intracellular levels of fructose 1,6-bisphosphate result in the phosphorylation of HPr at serine-46, upon which this molecule is no longer able to participate in the transport of any sugar. While this intracellular level of sugar at which the transport system is shut off may be sufficient to sustain the normal functioning of the cell, it may be limiting for the overproduction of the desired fine chemical. Thus, it may be desirable to modify the PTS proteins of the invention such that they are no longer responsive to such negative regulation, thereby permitting greater intracellular concentrations of one or more sugars to be achieved, and, by extension, more efficient production or greater yields of one or more fine chemicals from organisms containing such mutant PTS proteins.

This aforementioned list of mutagenesis strategies for PTS proteins to result in increased yields of a desired compound is not meant to be limiting; variations on these mutagenesis strategies will be readily apparent to one of ordinary skill in the art. By these mechanisms, the nucleic acid and protein molecules of the invention may be utilized to generate *C. glutamicum* or related strains of bacteria expressing mutated PTS nucleic acid and protein molecules such that the yield, production, and/or efficiency of production of a desired compound is improved. This desired compound may be any natural product of *C. glutamicum*, which includes the final products of biosynthesis pathways and intermediates of naturally-occurring metabolic pathways, as well as molecules which do not naturally occur in the metabolism of *C. glutamicum*, but which are produced by a *C. glutamicum* strain of the invention.

This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patent applications, patents, published patent applications, Tables, Appendices, and the sequence listing cited throughout this application are hereby incorporated by reference.

Exemplification**Example 1: Preparation of total genomic DNA of *Corynebacterium glutamicum* ATCC 13032**

- 5 A culture of *Corynebacterium glutamicum* (ATCC 13032) was grown overnight at 30°C with vigorous shaking in BHI medium (Difco). The cells were harvested by centrifugation, the supernatant was discarded and the cells were resuspended in 5 ml buffer-I (5% of the original volume of the culture — all indicated volumes have been calculated for 100 ml of culture volume). Composition of buffer-I: 140.34 g/l sucrose,
- 10 2.46 g/l $\text{MgSO}_4 \times 7\text{H}_2\text{O}$, 10 ml/l KH_2PO_4 solution (100 g/l, adjusted to pH 6.7 with KOH), 50 ml/l M12 concentrate (10 g/l $(\text{NH}_4)_2\text{SO}_4$, 1 g/l NaCl, 2 g/l $\text{MgSO}_4 \times 7\text{H}_2\text{O}$, 0.2 g/l CaCl_2 , 0.5 g/l yeast extract (Difco), 10 ml/l trace-elements-mix (200 mg/l $\text{FeSO}_4 \times \text{H}_2\text{O}$, 10 mg/l $\text{ZnSO}_4 \times 7\text{H}_2\text{O}$, 3 mg/l $\text{MnCl}_2 \times 4\text{H}_2\text{O}$, 30 mg/l H_3BO_3 , 20 mg/l $\text{CoCl}_2 \times 6\text{H}_2\text{O}$, 1 mg/l $\text{NiCl}_2 \times 6\text{H}_2\text{O}$, 3 mg/l $\text{Na}_2\text{MoO}_4 \times 2\text{H}_2\text{O}$, 500 mg/l complexing agent
- 15 (EDTA or citric acid), 100 ml/l vitamins-mix (0.2 mg/l biotin, 0.2 mg/l folic acid, 20 mg/l p-amino benzoic acid, 20 mg/l riboflavin, 40 mg/l ca-pantothenate, 140 mg/l nicotinic acid, 40 mg/l pyridoxole hydrochloride, 200 mg/l myo-inositol). Lysozyme was added to the suspension to a final concentration of 2.5 mg/ml. After an approximately 4 h incubation at 37°C, the cell wall was degraded and the resulting
- 20 protoplasts are harvested by centrifugation. The pellet was washed once with 5 ml buffer-I and once with 5 ml TE-buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8). The pellet was resuspended in 4 ml TE-buffer and 0.5 ml SDS solution (10%) and 0.5 ml NaCl solution (5 M) are added. After adding of proteinase K to a final concentration of 200 µg/ml, the suspension is incubated for ca.18 h at 37°C. The DNA was purified by
- 25 extraction with phenol, phenol-chloroform-isoamylalcohol and chloroform-isoamylalcohol using standard procedures. Then, the DNA was precipitated by adding 1/50 volume of 3 M sodium acetate and 2 volumes of ethanol, followed by a 30 min incubation at -20°C and a 30 min centrifugation at 12,000 rpm in a high speed centrifuge using a SS34 rotor (Sorvall). The DNA was dissolved in 1 ml TE-buffer containing 20
- 30 µg/ml RNaseA and dialysed at 4°C against 1000 ml TE-buffer for at least 3 hours. During this time, the buffer was exchanged 3 times. To aliquots of 0.4 ml of the dialysed DNA solution, 0.4 ml of 2 M LiCl and 0.8 ml of ethanol are added. After a 30 min incubation at -20°C, the DNA was collected by centrifugation (13,000 rpm, Biofuge Fresco, Heraeus, Hanau, Germany). The DNA pellet was dissolved in TE-buffer. DNA
- 35 prepared by this procedure could be used for all purposes, including southern blotting or construction of genomic libraries.

00604231.062700

Example 2: Construction of genomic libraries in *Escherichia coli* of *Corynebacterium glutamicum* ATCC13032.

Using DNA prepared as described in Example 1, cosmid and plasmid libraries were constructed according to known and well established methods (see e.g., Sambrook, J. et al. (1989) "Molecular Cloning : A Laboratory Manual", Cold Spring Harbor Laboratory Press, or Ausubel, F.M. et al. (1994) "Current Protocols in Molecular Biology", John Wiley & Sons.)

Any plasmid or cosmid could be used. Of particular use were the plasmids pBR322 (Sutcliffe, J.G. (1979) *Proc. Natl. Acad. Sci. USA*, 75:3737-3741); pACYC177 (Change & Cohen (1978) *J. Bacteriol* 134:1141-1156), plasmids of the pBS series (pBSSK+, pBSSK- and others; Stratagene, LaJolla, USA), or cosmids as SuperCos1 (Stratagene, LaJolla, USA) or Lorist6 (Gibson, T.J., Rosenthal A. and Waterson, R.H. (1987) *Gene* 53:283-286. Gene libraries specifically for use in *C. glutamicum* may be constructed using plasmid pSL109 (Lee, H.-S. and A. J. Sinskey (1994) *J. Microbiol. Biotechnol.* 4: 256-263).

Example 3: DNA Sequencing and Computational Functional Analysis

Genomic libraries as described in Example 2 were used for DNA sequencing according to standard methods, in particular by the chain termination method using ABI377 sequencing machines (see e.g., Fleischman, R.D. et al. (1995) "Whole-genome Random Sequencing and Assembly of *Haemophilus Influenzae* Rd., *Science*, 269:496-512). Sequencing primers with the following nucleotide sequences were used: 5'-GGAAACAGTATGACCATG-3' or 5'-GTAAACGACGCCAGT-3'.

Example 4: *In vivo* Mutagenesis

In vivo mutagenesis of *Corynebacterium glutamicum* can be performed by passage of plasmid (or other vector) DNA through *E. coli* or other microorganisms (e.g. *Bacillus* spp. or yeasts such as *Saccharomyces cerevisiae*) which are impaired in their capabilities to maintain the integrity of their genetic information. Typical mutator strains have mutations in the genes for the DNA repair system (e.g., mutHLS, mutD, mutT, etc.; for reference, see Rupp, W.D. (1996) DNA repair mechanisms, in: *Escherichia coli* and *Salmonella*, p. 2277-2294, ASM: Washington.) Such strains are well known to one of ordinary skill in the art. The use of such strains is illustrated, for example, in Greener, A. and Callahan, M. (1994) *Strategies* 7: 32-34.

Example 5: DNA Transfer Between *Escherichia coli* and *Corynebacterium glutamicum*

Several *Corynebacterium* and *Brevibacterium* species contain endogenous plasmids (as e.g., pHM1519 or pBL1) which replicate autonomously (for review see, e.g.,

- Martin, J.F. *et al.* (1987) *Biotechnology*, 5:137-146). Shuttle vectors for *Escherichia coli* and *Corynebacterium glutamicum* can be readily constructed by using standard vectors for *E. coli* (Sambrook, J. *et al.* (1989), "Molecular Cloning: A Laboratory Manual", Cold Spring Harbor Laboratory Press or Ausubel, F.M. *et al.* (1994) "Current Protocols in Molecular Biology", John Wiley & Sons) to which a origin or replication for and a suitable marker from *Corynebacterium glutamicum* is added. Such origins of replication are preferably taken from endogenous plasmids isolated from *Corynebacterium* and *Brevibacterium* species. Of particular use as transformation markers for these species are genes for kanamycin resistance (such as those derived from the Tn5 or Tn903 transposons) or chloramphenicol (Winnacker, E.L. (1987) "From Genes to Clones — Introduction to Gene Technology, VCH, Weinheim). There are numerous examples in the literature of the construction of a wide variety of shuttle vectors which replicate in both *E. coli* and *C. glutamicum*, and which can be used for several purposes, including gene overexpression (for reference, see *e.g.*, Yoshihama, M. *et al.* (1985) *J. Bacteriol.* 162:591-597, Martin J.F. *et al.* (1987) *Biotechnology*, 5:137-146 and Eikmanns, B.J. *et al.* (1991) *Gene*, 102:93-98).

Using standard methods, it is possible to clone a gene of interest into one of the shuttle vectors described above and to introduce such a hybrid vectors into strains of *Corynebacterium glutamicum*. Transformation of *C. glutamicum* can be achieved by protoplast transformation (Kastsumata, R. *et al.* (1984) *J. Bacteriol.* 159:306-311), electroporation (Liebl, E. *et al.* (1989) *FEMS Microbiol. Letters*, 53:399-303) and in cases where special vectors are used, also by conjugation (as described *e.g.* in Schäfer, A *et al.* (1990) *J. Bacteriol.* 172:1663-1666). It is also possible to transfer the shuttle vectors for *C. glutamicum* to *E. coli* by preparing plasmid DNA from *C. glutamicum* (using standard methods well-known in the art) and transforming it into *E. coli*. This transformation step can be performed using standard methods, but it is advantageous to use an Mcr-deficient *E. coli* strain, such as NM522 (Gough & Murray (1983) *J. Mol. Biol.* 166:1-19).

Genes may be overexpressed in *C. glutamicum* strains using plasmids which comprise pCG1 (U.S. Patent No. 4,617,267) or fragments thereof, and optionally the gene for kanamycin resistance from TN903 (Grindley, N.D. and Joyce, C.M. (1980) *Proc. Natl. Acad. Sci. USA* 77(12): 7176-7180). In addition, genes may be overexpressed in *C. glutamicum* strains using plasmid pSL109 (Lee, H.-S. and A. J. Sinskey (1994) *J. Microbiol. Biotechnol.* 4: 256-263).

Aside from the use of replicative plasmids, gene overexpression can also be achieved by integration into the genome. Genomic integration in *C. glutamicum* or other *Corynebacterium* or *Brevibacterium* species may be accomplished by well-known methods, such as homologous recombination with genomic region(s), restriction

09604231-062700

endonuclease mediated integration (REMI) (see, *e.g.*, DE Patent 19823834), or through the use of transposons. It is also possible to modulate the activity of a gene of interest by modifying the regulatory regions (*e.g.*, a promoter, a repressor, and/or an enhancer) by sequence modification, insertion, or deletion using site-directed methods (such as homologous recombination) or methods based on random events (such as transposon mutagenesis or REMI). Nucleic acid sequences which function as transcriptional terminators may also be inserted 3' to the coding region of one or more genes of the invention; such terminators are well-known in the art and are described, for example, in Winnacker, E.L. (1987) *From Genes to Clones – Introduction to Gene Technology*. VCH: Weinheim.

Example 6: Assessment of the Expression of the Mutant Protein

Observations of the activity of a mutated protein in a transformed host cell rely on the fact that the mutant protein is expressed in a similar fashion and in a similar quantity to that of the wild-type protein. A useful method to ascertain the level of transcription of the mutant gene (an indicator of the amount of mRNA available for translation to the gene product) is to perform a Northern blot (for reference see, for example, Ausubel *et al.* (1988) *Current Protocols in Molecular Biology*, Wiley: New York), in which a primer designed to bind to the gene of interest is labeled with a detectable tag (usually radioactive or chemiluminescent), such that when the total RNA of a culture of the organism is extracted, run on gel, transferred to a stable matrix and incubated with this probe, the binding and quantity of binding of the probe indicates the presence and also the quantity of mRNA for this gene. This information is evidence of the degree of transcription of the mutant gene. Total cellular RNA can be prepared from *Corynebacterium glutamicum* by several methods, all well-known in the art, such as that described in Bormann, E.R. *et al.* (1992) *Mol. Microbiol.* 6: 317-326.

To assess the presence or relative quantity of protein translated from this mRNA, standard techniques, such as a Western blot, may be employed (see, for example, Ausubel *et al.* (1988) *Current Protocols in Molecular Biology*, Wiley: New York). In this process, total cellular proteins are extracted, separated by gel electrophoresis, transferred to a matrix such as nitrocellulose, and incubated with a probe, such as an antibody, which specifically binds to the desired protein. This probe is generally tagged with a chemiluminescent or colorimetric label which may be readily detected. The presence and quantity of label observed indicates the presence and quantity of the desired mutant protein present in the cell.

Example 7: Growth of Genetically Modified *Corynebacterium glutamicum* — Media and Culture Conditions

- Genetically modified *Corynebacteria* are cultured in synthetic or natural growth media. A number of different growth media for *Corynebacteria* are both well-known and readily available (Lieb *et al.* (1989) *Appl. Microbiol. Biotechnol.*, 32:205-210; von der Osten *et al.* (1998) *Biotechnology Letters*, 11:11-16; Patent DE 4,120,867; Liebl (1992) "The Genus *Corynebacterium*, in: The Prokaryotes, Volume II, Balows, A. *et al.*, eds. Springer-Verlag). These media consist of one or more carbon sources, nitrogen sources, inorganic salts, vitamins and trace elements. Preferred carbon sources are sugars, such as mono-, di-, or polysaccharides. For example, glucose, fructose, mannose, galactose, ribose, sorbose, ribulose, lactose, maltose, sucrose, raffinose, starch or cellulose serve as very good carbon sources. It is also possible to supply sugar to the media via complex compounds such as molasses or other by-products from sugar refinement. It can also be advantageous to supply mixtures of different carbon sources. Other possible carbon sources are alcohols and organic acids, such as methanol, ethanol, acetic acid or lactic acid. Nitrogen sources are usually organic or inorganic nitrogen compounds, or materials which contain these compounds. Exemplary nitrogen sources include ammonia gas or ammonia salts, such as NH_4Cl or $(\text{NH}_4)_2\text{SO}_4$, NH_4OH , nitrates, urea, amino acids or complex nitrogen sources like corn steep liquor, soy bean flour, soy bean protein, yeast extract, meat extract and others.

- Inorganic salt compounds which may be included in the media include the chloride-, phosphorous- or sulfate- salts of calcium, magnesium, sodium, cobalt, molybdenum, potassium, manganese, zinc, copper and iron. Chelating compounds can be added to the medium to keep the metal ions in solution. Particularly useful chelating compounds include dihydroxyphenols, like catechol or protocatechuic acid, or organic acids, such as citric acid. It is typical for the media to also contain other growth factors, such as vitamins or growth promoters, examples of which include biotin, riboflavin, thiamin, folic acid, nicotinic acid, pantothenate and pyridoxin. Growth factors and salts frequently originate from complex media components such as yeast extract, molasses, corn steep liquor and others. The exact composition of the media compounds depends strongly on the immediate experiment and is individually decided for each specific case. Information about media optimization is available in the textbook "Applied Microbiol. Physiology, A Practical Approach" (eds. P.M. Rhodes, P.F. Stanbury, IRL Press (1997) pp. 53-73, ISBN 0 19 963577 3). It is also possible to select growth media from commercial suppliers, like standard 1 (Merck) or BHI (grain heart infusion, DIFCO) or others.

All medium components are sterilized, either by heat (20 minutes at 1.5 bar and 121°C) or by sterile filtration. The components can either be sterilized together or, if necessary, separately. All media components can be present at the beginning of growth, or they can optionally be added continuously or batchwise.

- 5 Culture conditions are defined separately for each experiment. The temperature should be in a range between 15°C and 45°C. The temperature can be kept constant or can be altered during the experiment. The pH of the medium should be in the range of 5 to 8.5, preferably around 7.0, and can be maintained by the addition of buffers to the media. An exemplary buffer for this purpose is a potassium phosphate buffer. Synthetic buffers
10 such as MOPS, HEPES, ACES and others can alternatively or simultaneously be used. It is also possible to maintain a constant culture pH through the addition of NaOH or NH₄OH during growth. If complex medium components such as yeast extract are utilized, the necessity for additional buffers may be reduced, due to the fact that many complex compounds have high buffer capacities. If a fermentor is utilized for culturing the micro-organisms, the pH can also be controlled using gaseous ammonia.

- The incubation time is usually in a range from several hours to several days. This time is selected in order to permit the maximal amount of product to accumulate in the broth. The disclosed growth experiments can be carried out in a variety of vessels, such as microtiter plates, glass tubes, glass flasks or glass or metal fermentors of different sizes.
20 For screening a large number of clones, the microorganisms should be cultured in microtiter plates, glass tubes or shake flasks, either with or without baffles. Preferably 100 ml shake flasks are used, filled with 10% (by volume) of the required growth medium. The flasks should be shaken on a rotary shaker (amplitude 25 mm) using a speed-range of 100 – 300 rpm. Evaporation losses can be diminished by the maintenance
25 of a humid atmosphere; alternatively, a mathematical correction for evaporation losses should be performed.

- If genetically modified clones are tested, an unmodified control clone or a control clone containing the basic plasmid without any insert should also be tested. The medium is inoculated to an OD₆₀₀ of 0.5 – 1.5 using cells grown on agar plates, such as CM plates
30 (10 g/l glucose, 2.5 g/l NaCl, 2 g/l urea, 10 g/l polypeptone, 5 g/l yeast extract, 5 g/l meat extract, 22 g/l NaCl, 2 g/l urea, 10 g/l polypeptone, 5 g/l yeast extract, 5 g/l meat extract, 22 g/l agar, pH 6.8 with 2M NaOH) that had been incubated at 30°C. Inoculation of the media is accomplished by either introduction of a saline suspension of *C. glutamicum* cells from CM plates or addition of a liquid preculture of this bacterium.

Example 8 – *In vitro* Analysis of the Function of Mutant Proteins

- The determination of activities and kinetic parameters of enzymes is well established in the art. Experiments to determine the activity of any given altered enzyme must be tailored to the specific activity of the wild-type enzyme, which is well within the ability of one of ordinary skill in the art. Overviews about enzymes in general, as well as specific details concerning structure, kinetics, principles, methods, applications and examples for the determination of many enzyme activities may be found, for example, in the following references: Dixon, M., and Webb, E.C., (1979) *Enzymes*. Longmans: London; Fersht, (1985) *Enzyme Structure and Mechanism*. Freeman: New York; Walsh, (1979) *Enzymatic Reaction Mechanisms*. Freeman: San Francisco; Price, N.C., Stevens, L. (1982) *Fundamentals of Enzymology*. Oxford Univ. Press: Oxford; Boyer, P.D., ed. (1983) *The Enzymes*, 3rd ed. Academic Press: New York; Bisswanger, H., (1994) *Enzymkinetik*, 2nd ed. VCH: Weinheim (ISBN 3527300325); Bergmeyer, H.U., Bergmeyer, J., Graßl, M., eds. (1983-1986) *Methods of Enzymatic Analysis*, 3rd ed., vol. I-XII, Verlag Chemie: Weinheim; and Ullmann's Encyclopedia of Industrial Chemistry (1987) vol. A9, "Enzymes". VCH: Weinheim, p. 352-363.

- The activity of proteins which bind to DNA can be measured by several well-established methods, such as DNA band-shift assays (also called gel retardation assays). The effect of such proteins on the expression of other molecules can be measured using reporter gene assays (such as that described in Kolmar, H. *et al.* (1995) *EMBO J.* 14: 3895-3904 and references cited therein). Reporter gene test systems are well known and established for applications in both pro- and eukaryotic cells, using enzymes such as beta-galactosidase, green fluorescent protein, and several others.
- The determination of activity of membrane-transport proteins can be performed according to techniques such as those described in Gennis, R.B. (1989) "Pores, Channels and Transporters", in *Biomembranes, Molecular Structure and Function*, Springer: Heidelberg, p. 85-137; 199-234; and 270-322.

Example 9: Analysis of Impact of Mutant Protein on the Production of the Desired Product

- The effect of the genetic modification in *C. glutamicum* on production of a desired compound (such as an amino acid) can be assessed by growing the modified microorganism under suitable conditions (such as those described above) and analyzing the medium and/or the cellular component for increased production of the desired product (i.e., an amino acid). Such analysis techniques are well known to one of ordinary skill in the art, and include spectroscopy, thin layer chromatography, staining

- methods of various kinds, enzymatic and microbiological methods, and analytical chromatography such as high performance liquid chromatography (see, for example, Ullman, Encyclopedia of Industrial Chemistry, vol. A2, p. 89-90 and p. 443-613, VCH: Weinheim (1985); Fallon, A. *et al.*, (1987) "Applications of HPLC in Biochemistry" in:
- 5 Laboratory Techniques in Biochemistry and Molecular Biology, vol. 17; Rehm *et al.* (1993) Biotechnology, vol. 3, Chapter III: "Product recovery and purification", page 469-714, VCH: Weinheim; Belter, P.A. *et al.* (1988) Bioseparations: downstream processing for biotechnology, John Wiley and Sons; Kennedy, J.F. and Cabral, J.M.S. (1992) Recovery processes for biological materials, John Wiley and Sons; Shaeiwitz,
- 10 J.A. and Henry, J.D. (1988) Biochemical separations, in: Ulmann's Encyclopedia of Industrial Chemistry, vol. B3, Chapter 11, page 1-27, VCH: Weinheim; and Dechow, F.J. (1989) Separation and purification techniques in biotechnology, Noyes Publications.)

- In addition to the measurement of the final product of fermentation, it is also
- 15 possible to analyze other components of the metabolic pathways utilized for the production of the desired compound, such as intermediates and side-products, to determine the overall productivity of the organism, yield, and/or efficiency of production of the compound. Analysis methods include measurements of nutrient levels in the medium (*e.g.*, sugars, hydrocarbons, nitrogen sources, phosphate, and other ions),
- 20 measurements of biomass composition and growth, analysis of the production of common metabolites of biosynthetic pathways, and measurement of gasses produced during fermentation. Standard methods for these measurements are outlined in Applied Microbial Physiology, A Practical Approach, P.M. Rhodes and P.F. Stanbury, eds., IRL Press, p. 103-129; 131-163; and 165-192 (ISBN: 0199635773) and references cited
- 25 therein.

Example 10: Purification of the Desired Product from *C. glutamicum* Culture

- Recovery of the desired product from the *C. glutamicum* cells or supernatant of the above-described culture can be performed by various methods well known in the art.
- 30 If the desired product is not secreted from the cells, the cells can be harvested from the culture by low-speed centrifugation, the cells can be lysed by standard techniques, such as mechanical force or sonication. The cellular debris is removed by centrifugation, and the supernatant fraction containing the soluble proteins is retained for further purification of the desired compound. If the product is secreted from the *C. glutamicum*
- 35 cells, then the cells are removed from the culture by low-speed centrifugation, and the supernate fraction is retained for further purification.

The supernatant fraction from either purification method is subjected to chromatography with a suitable resin, in which the desired molecule is either retained on a chromatography resin while many of the impurities in the sample are not, or where the impurities are retained by the resin while the sample is not. Such chromatography steps may be repeated as necessary, using the same or different chromatography resins. One of ordinary skill in the art would be well-versed in the selection of appropriate chromatography resins and in their most efficacious application for a particular molecule to be purified. The purified product may be concentrated by filtration or ultrafiltration, and stored at a temperature at which the stability of the product is maximized.

There are a wide array of purification methods known to the art and the preceding method of purification is not meant to be limiting. Such purification techniques are described, for example, in Bailey, J.E. & Ollis, D.F. *Biochemical Engineering Fundamentals*, McGraw-Hill: New York (1986).

The identity and purity of the isolated compounds may be assessed by techniques standard in the art. These include high-performance liquid chromatography (HPLC), spectroscopic methods, staining methods, thin layer chromatography, NIRS, enzymatic assay, or microbiologically. Such analysis methods are reviewed in: Patek *et al.* (1994) *Appl. Environ. Microbiol.* 60: 133-140; Malakhova *et al.* (1996) *Biotechnologia* 11: 27-32; and Schmidt *et al.* (1998) *Bioprocess Engineer.* 19: 67-70. Ulmann's Encyclopedia of Industrial Chemistry, (1996) vol. A27, VCH: Weinheim, p. 89-90, p. 521-540, p. 540-547, p. 559-566, 575-581 and p. 581-587; Michal, G. (1999) *Biochemical Pathways: An Atlas of Biochemistry and Molecular Biology*, John Wiley and Sons; Fallon, A. *et al.* (1987) *Applications of HPLC in Biochemistry in: Laboratory Techniques in Biochemistry and Molecular Biology*, vol. 17.

Example 11: Analysis of the Gene Sequences of the Invention

The comparison of sequences and determination of percent homology between two sequences are art-known techniques, and can be accomplished using a mathematical algorithm, such as the algorithm of Karlin and Altschul (1990) *Proc. Natl. Acad. Sci. USA* 87:2264-68, modified as in Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90:5873-77. Such an algorithm is incorporated into the NBLAST and XBLAST programs (version 2.0) of Altschul, *et al.* (1990) *J. Mol. Biol.* 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to PTS nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to PTS protein molecules of the invention. To obtain gapped alignments

for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, (1997) *Nucleic Acids Res.* 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, one of ordinary skill in the art will know how to optimize the parameters of the program (*e.g.*, XBLAST and NBLAST) for the specific sequence being analyzed.

Another example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Meyers and Miller ((1988) *Comput. Appl. Biosci.* 4: 11-17). Such an algorithm is incorporated into the ALIGN program (version 2.0) which is part of the GCG sequence alignment software package. When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used. Additional algorithms for sequence analysis are known in the art, and include ADVANCE and ADAM. described in Torelli and Robotti (1994) *Comput. Appl. Biosci.* 10:3-5; and FASTA, described in Pearson and Lipman (1988) *P.N.A.S.* 85:2444-8.

The percent homology between two amino acid sequences can also be accomplished using the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blosum 62 matrix or a PAM250 matrix, and a gap weight of 12, 10, 8, 6, or 4 and a length weight of 2, 3, or 4. The percent homology between two nucleic acid sequences can be accomplished using the GAP program in the GCG software package, using standard parameters, such as a gap weight of 50 and a length weight of 3.

A comparative analysis of the gene sequences of the invention with those present in Genbank has been performed using techniques known in the art (see, *e.g.*, Bexevanis and Ouellette, eds. (1998) *Bioinformatics: A Practical Guide to the Analysis of Genes and Proteins.* John Wiley and Sons: New York). The gene sequences of the invention were compared to genes present in Genbank in a three-step process. In a first step, a BLASTN analysis (*e.g.*, a local alignment analysis) was performed for each of the sequences of the invention against the nucleotide sequences present in Genbank, and the top 500 hits were retained for further analysis. A subsequent FASTA search (*e.g.*, a combined local and global alignment analysis, in which limited regions of the sequences are aligned) was performed on these 500 hits. Each gene sequence of the invention was subsequently globally aligned to each of the top three FASTA hits, using the GAP program in the GCG software package (using standard parameters). In order to obtain correct results, the length of the sequences extracted from Genbank were adjusted to the length of the query sequences by methods well-known in the art. The results of this analysis are set forth in Table 4. The resulting data is identical to that which would have been obtained had a GAP (global) analysis alone been performed on each of the genes of

the invention in comparison with each of the references in Genbank, but required significantly reduced computational time as compared to such a database-wide GAP (global) analysis. Sequences of the invention for which no alignments above the cutoff values were obtained are indicated on Table 4 by the absence of alignment information.

- 5 It will further be understood by one of ordinary skill in the art that the GAP alignment homology percentages set forth in Table 4 under the heading "% homology (GAP)" are listed in the European numerical format, wherein a ',' represents a decimal point. For example, a value of "40,345" in this column represents "40.345%".

10 **Example 12: Construction and Operation of DNA Microarrays**

The sequences of the invention may additionally be used in the construction and application of DNA microarrays (the design, methodology, and uses of DNA arrays are well known in the art, and are described, for example, in Schena, M. *et al.* (1995) *Science* 270: 467-470; Wodicka, L. *et al.* (1997) *Nature Biotechnology* 15: 1359-1367;

- 15 DeSaizieu, A. *et al.* (1998) *Nature Biotechnology* 16: 45-48; and DeRisi, J.L. *et al.* (1997) *Science* 278: 680-686).

DNA microarrays are solid or flexible supports consisting of nitrocellulose, nylon, glass, silicone, or other materials. Nucleic acid molecules may be attached to the surface in an ordered manner. After appropriate labeling, other nucleic acids or nucleic acid mixtures can be hybridized to the immobilized nucleic acid molecules, and the label may be used to monitor and measure the individual signal intensities of the hybridized molecules at defined regions. This methodology allows the simultaneous quantification of the relative or absolute amount of all or selected nucleic acids in the applied nucleic acid sample or mixture. DNA microarrays, therefore, permit an analysis of the

20 expression of multiple (as many as 6800 or more) nucleic acids in parallel (see, e.g., Schena, M. (1996) *BioEssays* 18(5): 427-431).

The sequences of the invention may be used to design oligonucleotide primers which are able to amplify defined regions of one or more *C. glutamicum* genes by a nucleic acid amplification reaction such as the polymerase chain reaction. The choice and design of the 5' or 3' oligonucleotide primers or of appropriate linkers allows the covalent attachment of the resulting PCR products to the surface of a support medium described above (and also described, for example, Schena, M. *et al.* (1995) *Science* 270: 467-470).

Nucleic acid microarrays may also be constructed by *in situ* oligonucleotide synthesis as described by Wodicka, L. *et al.* (1997) *Nature Biotechnology* 15: 1359-1367. By photolithographic methods, precisely defined regions of the matrix are exposed to light. Protective groups which are photolabile are thereby activated and

09604551.062700

undergo nucleotide addition, whereas regions that are masked from light do not undergo any modification. Subsequent cycles of protection and light activation permit the synthesis of different oligonucleotides at defined positions. Small, defined regions of the genes of the invention may be synthesized on microarrays by solid phase

5 oligonucleotide synthesis.

The nucleic acid molecules of the invention present in a sample or mixture of nucleotides may be hybridized to the microarrays. These nucleic acid molecules can be labeled according to standard methods. In brief, nucleic acid molecules (*e.g.*, mRNA molecules or DNA molecules) are labeled by the incorporation of isotopically or

10 fluorescently labeled nucleotides, *e.g.*, during reverse transcription or DNA synthesis. Hybridization of labeled nucleic acids to microarrays is described (*e.g.*, in Schena, M. *et al.* (1995) *supra*; Wodicka, L. *et al.* (1997), *supra*; and DeSaizieu A. *et al.* (1998), *supra*). The detection and quantification of the hybridized molecule are tailored to the specific incorporated label. Radioactive labels can be detected, for example, as

15 described in Schena, M. *et al.* (1995) *supra* and fluorescent labels may be detected, for example, by the method of Shalon *et al.* (1996) *Genome Research* 6: 639-645).

The application of the sequences of the invention to DNA microarray technology, as described above, permits comparative analyses of different strains of *C. glutamicum* or other Corynebacteria. For example, studies of inter-strain variations

20 based on individual transcript profiles and the identification of genes that are important for specific and/or desired strain properties such as pathogenicity, productivity and stress tolerance are facilitated by nucleic acid array methodologies. Also, comparisons of the profile of expression of genes of the invention during the course of a fermentation reaction are possible using nucleic acid array technology.

25

Example 13: Analysis of the Dynamics of Cellular Protein Populations (Proteomics)

The genes, compositions, and methods of the invention may be applied to study the interactions and dynamics of populations of proteins, termed 'proteomics'. Protein

30 populations of interest include, but are not limited to, the total protein population of *C. glutamicum* (*e.g.*, in comparison with the protein populations of other organisms), those proteins which are active under specific environmental or metabolic conditions (*e.g.*, during fermentation, at high or low temperature, or at high or low pH), or those proteins which are active during specific phases of growth and development.

35 Protein populations can be analyzed by various well-known techniques, such as gel electrophoresis. Cellular proteins may be obtained, for example, by lysis or extraction, and may be separated from one another using a variety of electrophoretic

techniques. Sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) separates proteins largely on the basis of their molecular weight. Isoelectric focusing polyacrylamide gel electrophoresis (IEF-PAGE) separates proteins by their isoelectric point (which reflects not only the amino acid sequence but also posttranslational modifications of the protein). Another, more preferred method of protein analysis is the consecutive combination of both IEF-PAGE and SDS-PAGE, known as 2-D-gel electrophoresis (described, for example, in Hermann *et al.* (1998) *Electrophoresis* 19: 3217-3221; Fountoulakis *et al.* (1998) *Electrophoresis* 19: 1193-1202; Langen *et al.* (1997) *Electrophoresis* 18: 1184-1192; Antelmann *et al.* (1997) *Electrophoresis* 18: 1451-1463). Other separation techniques may also be utilized for protein separation, such as capillary gel electrophoresis; such techniques are well known in the art.

Proteins separated by these methodologies can be visualized by standard techniques, such as by staining or labeling. Suitable stains are known in the art, and include Coomassie Brilliant Blue, silver stain, or fluorescent dyes such as Sypro Ruby (Molecular Probes). The inclusion of radioactively labeled amino acids or other protein precursors (*e.g.*, ^{35}S -methionine, ^{35}S -cysteine, ^{14}C -labelled amino acids, ^{15}N -amino acids, $^{15}\text{NO}_3$ or $^{15}\text{NH}_4^+$ or ^{13}C -labelled amino acids) in the medium of *C. glutamicum* permits the labeling of proteins from these cells prior to their separation. Similarly, fluorescent labels may be employed. These labeled proteins can be extracted, isolated and separated according to the previously described techniques.

Proteins visualized by these techniques can be further analyzed by measuring the amount of dye or label used. The amount of a given protein can be determined quantitatively using, for example, optical methods and can be compared to the amount of other proteins in the same gel or in other gels. Comparisons of proteins on gels can be made, for example, by optical comparison, by spectroscopy, by image scanning and analysis of gels, or through the use of photographic films and screens. Such techniques are well-known in the art.

To determine the identity of any given protein, direct sequencing or other standard techniques may be employed. For example, N- and/or C-terminal amino acid sequencing (such as Edman degradation) may be used, as may mass spectrometry (in particular MALDI or ESI techniques (see, *e.g.*, Langen *et al.* (1997) *Electrophoresis* 18: 1184-1192)). The protein sequences provided herein can be used for the identification of *C. glutamicum* proteins by these techniques.

The information obtained by these methods can be used to compare patterns of protein presence, activity, or modification between different samples from various biological conditions (*e.g.*, different organisms, time points of fermentation, media conditions, or different biotopes, among others). Data obtained from such experiments

alone, or in combination with other techniques, can be used for various applications, such as to compare the behavior of various organisms in a given (*e.g.*, metabolic) situation, to increase the productivity of strains which produce fine chemicals or to increase the efficiency of the production of fine chemicals.

5

Equivalents

Those of ordinary skill in the art will recognize, or will be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

10

03604231.002700

What is claimed:

1. An isolated nucleic acid molecule from *Corynebacterium glutamicum* encoding a phosphoenolpyruvate: sugar phosphotransferase system protein, or a portion thereof, provided that the nucleic acid molecule does not consist of any of the F-designated genes set forth in Table 1.
2. The isolated nucleic acid molecule of claim 1, wherein said phosphoenolpyruvate: sugar phosphotransferase system protein is selected from the group consisting of proteins involved in the transport of glucose, sucrose, mannose, fructose, mannitol, raffinose, ribulose, ribose, lactose, maltose, sorbose, sorbitol, xylose, and galactose.
3. An isolated *Corynebacterium glutamicum* nucleic acid molecule selected from the group consisting of those sequences set forth in Appendix A, or a portion thereof, provided that the nucleic acid molecule does not consist of any of the F-designated genes set forth in Table 1.
4. An isolated nucleic acid molecule which encodes a polypeptide sequence selected from the group consisting of those sequences set forth in Appendix B, provided that the nucleic acid molecule does not consist of any of the F-designated genes set forth in Table 1.
5. An isolated nucleic acid molecule which encodes a naturally occurring allelic variant of a polypeptide selected from the group of amino acid sequences consisting of those sequences set forth in Appendix B, provided that the nucleic acid molecule does not consist of any of the F-designated genes set forth in Table 1.
6. An isolated nucleic acid molecule comprising a nucleotide sequence which is at least 50% homologous to a nucleotide sequence selected from the group consisting of those sequences set forth in Appendix A, or a portion thereof, provided that the nucleic acid molecule does not consist of any of the F-designated genes set forth in Table 1.
7. An isolated nucleic acid molecule comprising a fragment of at least 15 nucleotides of a nucleic acid comprising a nucleotide sequence selected from the group

consisting of those sequences set forth in Appendix A, provided that the nucleic acid molecule does not consist of any of the F-designated genes set forth in Table 1.

8. An isolated nucleic acid molecule which hybridizes to the nucleic acid molecule of any one of claims 1-7 under stringent conditions.

9. An isolated nucleic acid molecule comprising the nucleic acid molecule of claim 1 or a portion thereof and a nucleotide sequence encoding a heterologous polypeptide.

10. A vector comprising the nucleic acid molecule of claim 1.

11. The vector of claim 10, which is an expression vector.

12. A host cell transfected with the expression vector of claim 11.

13. The host cell of claim 12, wherein said cell is a microorganism.

14. The host cell of claim 13, wherein said cell belongs to the genus *Corynebacterium* or *Brevibacterium*.

15. The host cell of claim 12, wherein the expression of said nucleic acid molecule results in the modulation in production of a fine chemical from said cell.

16. The host cell of claim 15, wherein said fine chemical is selected from the group consisting of: organic acids, proteinogenic amino acids, nonproteinogenic amino acids, purine and pyrimidine bases, nucleosides, nucleotides, lipids, saturated and unsaturated fatty acids, diols, carbohydrates, aromatic compounds, vitamins, cofactors, polyketides, and enzymes.

17. A method of producing a polypeptide comprising culturing the host cell of claim 12 in an appropriate culture medium to, thereby, produce the polypeptide.

18. An isolated phosphoenolpyruvate: sugar phosphotransferase system polypeptide from *Corynebacterium glutamicum*, or a portion thereof.

19. The protein of claim 18, wherein said phosphoenolpyruvate: sugar phosphotransferase system protein is selected from the group consisting of proteins

03604231, 062700

involved in the transport of glucose, sucrose, mannose, fructose, mannitol, raffinose, ribulose, ribose, lactose, maltose, sorbose, and galactose.

20. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of those sequences set forth in Appendix B, provided that the amino acid sequence is not encoded by any of the F-designated genes set forth in Table 1.

21. An isolated polypeptide comprising a naturally occurring allelic variant of a polypeptide comprising an amino acid sequence selected from the group consisting of those sequences set forth in Appendix B, or a portion thereof, provided that the amino acid sequence is not encoded by any of the F-designated genes set forth in Table 1.

22. The isolated polypeptide of claim 18, further comprising heterologous amino acid sequences.

23. An isolated polypeptide which is encoded by a nucleic acid molecule comprising a nucleotide sequence which is at least 50% homologous to a nucleic acid selected from the group consisting of those sequences set forth in Appendix A, provided that the nucleic acid molecule does not consist of any of the F-designated nucleic acid molecules set forth in Table 1.

24. An isolated polypeptide comprising an amino acid sequence which is at least 50% homologous to an amino acid sequence selected from the group consisting of those sequences set forth in Appendix B, provided that the amino acid sequence is not encoded by any of the F-designated genes set forth in Table 1.

25. A method for producing a fine chemical, comprising culturing a cell containing a vector of claim 12 such that the fine chemical is produced.

26. The method of claim 25, wherein said method further comprises the step of recovering the fine chemical from said culture.

27. The method of claim 25, wherein said method further comprises the step of transfecting said cell with the vector of claim 11 to result in a cell containing said vector.

28. The method of claim 25, wherein said cell belongs to the genus *Corynebacterium* or *Brevibacterium*.

29. The method of claim 25, wherein said cell is selected from the group consisting of: *Corynebacterium glutamicum*, *Corynebacterium herculis*, *Corynebacterium lilium*, *Corynebacterium acetoacidophilum*, *Corynebacterium acetoglutamicum*,
- 5 *Corynebacterium acetophilum*, *Corynebacterium ammoniagenes*, *Corynebacterium fujikense*, *Corynebacterium nitrilophilus*, *Brevibacterium ammoniagenes*, *Brevibacterium butanicum*, *Brevibacterium divaricatum*, *Brevibacterium flavum*, *Brevibacterium healii*, *Brevibacterium ketoglutamicum*, *Brevibacterium ketosoreductum*, *Brevibacterium lactofermentum*, *Brevibacterium linens*,
- 10 *Brevibacterium paraffinolyticum*, and those strains set forth in Table 3.

30. The method of claim 25, wherein expression of the nucleic acid molecule from said vector results in modulation of production of said fine chemical.

- 15 31. The method of claim 25, wherein said fine chemical is selected from the group consisting of: organic acids, proteinogenic amino acids, nonproteinogenic amino acids, purine and pyrimidine bases, nucleosides, nucleotides, lipids, saturated and unsaturated fatty acids, diols, carbohydrates, aromatic compounds, vitamins, cofactors, polyketides, and enzymes.

- 20 32. The method of claim 25, wherein said fine chemical is an amino acid.

33. The method of claim 32, wherein said amino acid is drawn from the group consisting of: lysine, glutamate, glutamine, alanine, aspartate, glycine, serine, threonine,
- 25 methionine, cysteine, valine, leucine, isoleucine, arginine, proline, histidine, tyrosine, phenylalanine, and tryptophan.

34. A method for producing a fine chemical, comprising culturing a cell whose genomic DNA has been altered by the inclusion of a nucleic acid molecule of any one of
- 30 claims 1-9.

35. A method for diagnosing the presence or activity of *Corynebacterium diphtheriae* in a subject, comprising detecting the presence of one or more of the sequences set forth in Appendix A or Appendix B in the subject, provided that the sequences are not or are not encoded by any of the F-designated sequences set forth in Table 1, thereby diagnosing the presence or activity of *Corynebacterium diphtheriae* in the subject.

36. A host cell comprising a nucleic acid molecule selected from the group consisting of the nucleic acid molecules set forth in Appendix A, wherein the nucleic acid molecule is disrupted.

5

37. A host cell comprising a nucleic acid molecule selected from the group consisting of the nucleic acid molecules set forth in Appendix A, wherein the nucleic acid molecule comprises one or more nucleic acid modifications from the sequence set forth in Appendix A.

10

38. A host cell comprising a nucleic acid molecule selected from the group consisting of the nucleic acid molecules set forth in Appendix A, wherein the regulatory region of the nucleic acid molecule is modified relative to the wild-type regulatory region of the molecule.

15

002290-1-004960

**CORYNEBACTERIUM GLUTAMICUM GENES ENCODING
PHOSPHOENOLPYRUVATE: SUGAR PHOSPHOTRANSFERASE SYSTEM
PROTEINS**

5

Abstract of the Disclosure

Isolated nucleic acid molecules, designated PTS nucleic acid molecules, which encode novel PTS proteins from *Corynebacterium glutamicum* are described. The invention also provides antisense nucleic acid molecules, recombinant expression
10 vectors containing PTS nucleic acid molecules, and host cells into which the expression vectors have been introduced. The invention still further provides isolated PTS proteins, mutated PTS proteins, fusion proteins, antigenic peptides and methods for the improvement of production of a desired compound from *C. glutamicum* based on genetic engineering of PTS genes in this organism.

15

002290-1324960

DECLARATION, PETITION AND POWER OF ATTORNEY FOR PATENT APPLICATION

My residence, post office address and citizenship are as stated below next to my name,

**CORYNEBACTERIUM GLUTAMICUM GENES ENCODING
PHOSPHOENOLPYRUVATE: SUGAR PHOSPHOTRANSFERASE SYSTEM PROTEINS**

X is attached hereto.

was filed on _____ as Application Serial No. _____

and was amended on _____
(if applicable)

I acknowledge the duty to disclose to the Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

CLAIM OF BENEFIT OF EARLIER FOREIGN APPLICATION(S)

I hereby claim priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below, and have also identified below any foreign application(s) for patent or inventor's certificate filed by me on the same subject matter having a filing date before that of the application(s) from which priority is claimed.

Check one:

☐ no such applications have been filed.

☒ such applications have been filed as follows

**EARLIEST FOREIGN APPLICATION(S), IF ANY, FILED WITHIN 12 MONTHS
(6 MONTHS FOR DESIGN) PRIOR TO THIS U.S. APPLICATION**

Country	Application Number	Date of Filing (month,day,year)	Priority Claimed Under 35 USC 119
DE	19942095.5	09/03/99	<input checked="" type="checkbox"/> Yes No <input type="checkbox"/>
DE	19942097.1	09/03/99	<input checked="" type="checkbox"/> Yes No <input type="checkbox"/>
			<input type="checkbox"/> Yes No <input checked="" type="checkbox"/>
			<input type="checkbox"/> Yes No <input type="checkbox"/>
			<input type="checkbox"/> Yes No <input type="checkbox"/>

**ALL FOREIGN APPLICATION(S), IF ANY FILED MORE THAN 12 MONTHS
(6 MONTHS FOR DESIGN) PRIOR TO THIS U.S. APPLICATION**

CLAIM FOR BENEFIT OF U.S. PROVISIONAL APPLICATION(S)

I hereby claim the benefit under 35 U.S.C. §119(e) of any United States provisional application(s) listed below.

60/142,691
(Application Serial No.)

July 1, 1999
(Filing Date)

60/150,310
(Application Serial No.)

August 23, 1999
(Filing Date)

03604231.62703

CLAIM FOR BENEFIT OF EARLIER U.S./PCT APPLICATION(S)

I hereby claim the benefit under Title 35, United States Code, §120 of any earlier United States application(s) or PCT international application(s) designating the United States listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the earlier application(s) in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose to the Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56 which became available between the filing date(s) of the earlier application(s) and the national or PCT international filing date of this application. As to subject matter of this application which is common to my earlier application(s), if any, described below, I do not know and do not believe that the same was known or used by others in the United States or patented or described in a printed publication in any country before my invention thereof, or patented or described in a printed publication in any country or in public use or on sale in the United States more than one year prior to the date(s) of said earlier application(s), or first patented or caused to be patented or made the subject of an inventor's certificate by me or my legal representatives or assigns in a country foreign to the United States prior to the date(s) of said earlier application(s) on an application filed more than twelve months (six months if this application is for a design) before the filing of said earlier application(s); and I acknowledge that no application for patent or inventor's certificate on said subject matter has been filed by me or my representatives or assigns in any country foreign to the United States except those identified herein.

(Application Serial No.)

(Filing Date)

(Status)
(patented,pending,aband.)

(Application Serial No.)

(Filing Date)

(Status)
(patented,pending,aband.)

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorneys and/or agents to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

James E. Cockfield	Reg. No. 19,162	Megan E. Williams	Reg. No. 43,270
Thomas V. Smurzynski	Reg. No. 24,798	Nicholas P. Triano III	Reg. No. 36,397
Ralph A. Loren	Reg. No. 29,325	Peter C. Lauro	Reg. No. 32,360
Giulio A. DeConti, Jr.	Reg. No. 31,503	Timothy J. Douros	Reg. No. 41,716
Ann Lamport Hammitte	Reg. No. 34,858	DeAnn F. Smith	Reg. No. 36,683
Elizabeth A. Hanley	Reg. No. 33,505	William D. DeVaul	Reg. No. 42,483
Amy E. Mandragouras	Reg. No. 36,207	David J. Rikkers	Reg. No. 43,882
Anthony A. Laurentano	Reg. No. 38,220	Chi Suk Kim	Reg. No. 42,728
Jane E. Remillard	Reg. No. 38,872	Maria Laccotripe Zacharakis	Limited Recognition Under 37 C.F.R. § 10.9(b)
Jeremiah Lynch	Reg. No. 17,425		
Kevin J. Canning	Reg. No. 35,470	Debra J. Milasincic	Reg. No. P46,931
David A. Lane, Jr.	Reg. No. 39,261	David R. Burns	Reg. No. P46,590
Jeanne M. DiGiorgio	Reg. No. 41,710		

Send Correspondence to Giulio A. DeConti, Jr., Esq. at **Customer Number: 000959** whose address is:

Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109

Direct Telephone Calls to: (name and telephone number)

Giulio A. DeConti, Jr., Esq., (617) 227-7400

Wherefore I petition that letters patent be granted to me for the invention or discovery described and claimed in the attached specification and claims, and hereby subscribe my name to said specification and claims and to the foregoing declaration, power of attorney, and this petition.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or first inventor Markus Pompejus	
Inventor's signature	Date
Residence Wenjenstrasse 21, 67251 Freinsheim, Germany	
Citizenship Germany	
Post Office Address (if different)	

Full name of second inventor, if any Burkhard Kröger	
Inventor's signature	Date
Residence Im Waldhof 1, 67117 Limburgerhof, Germany	
Citizenship Germany	
Post Office Address (if different)	

Full name of third inventor, if any Hartwig Schröder	
Inventor's signature	Date
Residence Goethestr. 5, 69226 Nussloch, Germany	
Citizenship Germany	
Post Office Address (if different)	

Full name of fourth inventor, if any Oskar Zelder	
Inventor's signature	Date
Residence Rossmarktstr. 27, 67346 Speyer, Germany	
Citizenship Germany	
Post Office Address (if different)	

Full name of fifth inventor, if any Gregor Haberhauer	
Inventor's signature	Date
Residence Moselstr. 42, 67117 Limburgerhof, Germany	
Citizenship Germany	
Post Office Address (if different)	

TABLE 1: Genes Included in the Invention

PHOSPHOENOLPYRUVATE: SUGAR PHOSPHOTRANSFERASE SYSTEM

Nucleotide SEQ ID NO	Amino Acid SEQ ID NO	Identification Code	Contig	NT Start	NT Stop	Function
1	2	RX500315				PTS SYSTEM, SUCROSE-SPECIFIC IABC COMPONENT (EIABC-SCR), SUCROSE- PERMEASE IABC COMPONENT (PHOSPHOTRANSFERASE ENZYME IABC COMPONENT) (EC 2.7.1.69)
3	4	F RXA00315	GR00053	6537	5452	PTS SYSTEM, BETA-GLUCOSIDE-SPECIFIC IABC COMPONENT (EIABC-BGL, BETA-GLUCOSIDES- PERMEASE IABC COMPONENT) (PHOSPHOTRANSFERASE ENZYME II, ABC COMPONENT) (EC 2.7.1.69)
5	6	RXA01503	GR00424	10392	10640	PTS SYSTEM, BETA-GLUCOSIDES-SPECIFIC IABC COMPONENT (EIABC-BGL, BETA-GLUCOSIDES- PERMEASE IABC COMPONENT) (PHOSPHOTRANSFERASE ENZYME II, ABC COMPONENT) (EC 2.7.1.69)
7	8	RXN01299	VV0068	11954	9891	PTS SYSTEM, FRUCTOSE-SPECIFIC IIBC COMPONENT (EC 2.7.1.69)
9	10	F RXA01299	GR00375	6	446	PTS SYSTEM, FRUCTOSE-SPECIFIC IIBC COMPONENT (EC 2.7.1.69)
11	12	F RXA01863	GR00538	2154	2633	PTS SYSTEM, FRUCTOSE-SPECIFIC IIBC COMPONENT (EC 2.7.1.69)
13	14	F RXA01869	GR00540	77	631	PTS SYSTEM, FRUCTOSE-SPECIFIC IIBC COMPONENT (EC 2.7.1.69)
15	16	RXA00951	GR00261	564	172	PTS SYSTEM, FRUCTOSE-SPECIFIC IIBC COMPONENT (EC 2.7.1.69)
17	18	RXN01244	VV0068	14141	15844	PTS SYSTEM, MANNITOL (CRYPTIC), SPECIFIC IIA COMPONENT (EIIA-CMNTL) (MANNITOL (CRYPTIC)- PERMEASE IIA COMPONENT) (PHOSPHOTRANSFERASE ENZYME II, A COMPONENT) (EC 2.7.1.69)
19	20	F RXA01244	GR00359	4837	3329	PHOSPHOENOLPYRUVATE-PROTEIN PHOSPHOTRANSFERASE (EC 2.7.3.9)
21	22	RXA01300	GR00375	637	903	PHOSPHOENOLPYRUVATE-PROTEIN PHOSPHOTRANSFERASE (EC 2.7.3.9)
23	24	RXN03002	VV0236	1437	1644	PHOSPHOCARRIER PROTEIN HPR
25	26	RXC00653	VV0260	1634	1082	PTS SYSTEM, MANNITOL (CRYPTIC), SPECIFIC IIA COMPONENT (EIIA-CMNTL) (MANNITOL (CRYPTIC)-PERMEASE IIA COMPONENT) (PHOSPHOTRANSFERASE ENZYME II, A COMPONENT) (EC 2.7.1.69)
27	28	RXC03001				Membrane Spanning Protein involved in PTS system
29	30	RXN01943	VV0120	4326	6374	PTS SYSTEM, GLUCOSE-SPECIFIC IABC COMPONENT (EC 2.7.1.69)
31	32	F RXA02191	GR00642	3395	4633	PHOSPHOENOLPYRUVATE SUGAR PHOSPHOTRANSFERASE
33	34	F RXA01943	GR00557	3644	3540	cr gene, phosphotransferase system glucose-specific enzyme III

TABLE 2: GENES IDENTIFIED FROM GENBANK

Attorney Docket No.: BGI-122CP

GenBank™ Accession No.	Gene Name	Gene Function	Reference
A09073	ppg	Phosphoenol pyruvate carboxylase	Bachmann, B. et al. "DNA fragment coding for phosphoenolpyruvate carboxylase, recombinant DNA carrying said fragment, strains carrying the recombinant DNA, and method for producing L- amino acids using said strains," Patent: EP 0038940-A 3 03/21/90
A45579, A45581, A45583, A45585 A45587		Threonine dehydratase	Moedeke, B. et al. "Production of L- isoleucine by means of recombinant micro-organisms with deregulated threonine dehydratase," Patent: WO 95/9442-A 5 07/20/95
AB005132	murC, flsQ, flsZ		Kobayashi, M. et al. "Cloning, sequencing, and characterization of the flsZ gene from corynebacter bacteria," <i>Biochem. Biophys. Res. Commun.</i> , 236(2):383-388 (1997)
AB015023	murC, flsQ		Wach, M. et al. "A murC gene from Corynebacter bacteria," <i>Appl. Microbiol. Biotechnol.</i> , 51(2):223-228 (1999)
AB018530	dsrK		Kimura, E. et al. "Molecular cloning of a novel gene, dsrK, which rescues the detergent sensitivity of a mutant derived from <i>Brevibacterium lactofermentum</i> ," <i>Biosci. Biotechnol. Biochem.</i> , 60(10):1565-1570 (1996)
AB018531	dsrK1, dsrK2		
AB020624	mutI	D-glutamate racemase	
AB023377	tkt	transketolase	
AB024708	gluB, gluD	Glutamine 2-oxoglutarate aminotransferase	
AB025424	acn	large and small subunits	
AB027714	rep	aconitase	
AB027715	rep; aad	Replication protein	
AF005242	argC	Replication protein; aminoglycoside adenyltransferase	
AF005635	gluA	N-acetylglutamate-5-semialdehyde dehydrogenase	
AF030405	hisA	Glutamine synthetase	
AF030520	argG	cyclase	
AF031518	argE	Argininosuccinate synthetase	
AF036932	aroD	Ornithine carbamoyltransferase	
		3-dehydroquinate dehydratase	

09604231 062700

GenBank™ Accession No.	Gene Name	Gene Function	Reference
AF038548	pyc	Pyruvate carboxylase	
AF038651	dcfA;E; apt; rel	Dipeptide-binding protein, adenine phosphoribosyltransferase, GTP pyrophosphokinase	Wehner, L. et al. "The role of the Corynebacterium glutamicum rel gene in (p)ppGpp metabolism." <i>Microbiology</i> , 144:1853-1862 (1998)
AF041436	argR	Arginine repressor	
AF045998	impA	Imidazol monophosphate phosphatase	
AF048764	argH	Argininosuccinate lyase	
AF049897	argC; argJ; argB; argD; argI; argK; argG; argH	N-acetylglutarylphosphatase; ornithine acetyltransferase; N-acetylglutamate kinase; acetylornithine transaminase; ornithine carbamoyltransferase; arginine repressor; argininosuccinate synthase; argininosuccinate lyase	
AF050109	inhA	Enoyl-acyl carrier protein reductase	
AF050166	hscG	ATP phosphoribosyltransferase	
AF051846	hscA	Phosphoribosylformimino-5-amino-1-phosphoribosyl-4-imidazolecarboxamide isomerase	
AF052652	metA	Homoserine O-acetyltransferase	Park, S. et al. "Isolation and analysis of metA, a methionine biosynthetic gene encoding homoserine acetyltransferase in Corynebacterium glutamicum," <i>Mol. Cells</i> , 8(3):286-294 (1998)
AF053071	aroB	Dehydroquinate synthetase	
AF060558	hish	Glutamine amidotransferase	
AF086704	hishE	Phosphoribosyl-ATP-pyrophosphohydrolyase	
AF114233	aroA	5-enolpyruvylshikimate 3-phosphate synthase	
AF116184	pand	L-aspartate-alpha-decarboxylase precursor	Dusch, N. et al. "Expression of the Corynebacterium glutamicum pand gene encoding L-aspartate-alpha-decarboxylase leads to pantothinate overproduction in Escherichia coli." <i>Appl. Environ. Microbiol.</i> , 65(4):1530-1539 (1999)

09604231.062700

GenBank™ Accession No.	Gene Name	Gene Function	Reference
AF124518	aroJ, aroE	3-dehydroquinate: shikimate dehydrogenase	
AF124600	aroC, aroK, aroB, pepQ	Chorismate synthase; shikimate kinase; 3-dehydroquinate synthase; putative cytoplasmic peptidase	
AF145897	inhA		
AF145898	inhA		
AJ001436	ecfP	Transport of ectoine, glycine betaine, proline	Peter, H. et al. "Corynebacterium glutamicum is equipped with four secondary carriers for compatible solutes. Identification, sequencing, and characterization of the proline/ectoine uptake system, ProP, and the ectoine/proline/glycine betaine carrier, EcfP." <i>J. Bacteriol.</i> , 180(22):6005-6012 (1998)
AJ004934	dapD	Tetrahydrodipicolinate succinylase (incomplete)	Weinmann, A. et al. "Different modes of diaminoipinate synthesis and their role in cell wall integrity: A study with <i>Corynebacterium glutamicum</i> ," <i>J. Bacteriol.</i> , 180(12):3159-3165 (1998)
AJ007732	ppe, secC, amt, ocd, soxA	Phosphoenolpyruvate:carboxylase; γ -high affinity ammonium uptake protein; putative ornithine-cyclodextranase; sarcosine oxidase	Jakoby, M. et al. "Nitrogen regulation in <i>Corynebacterium glutamicum</i> : Isolation of genes involved in biochemical characterization of corresponding proteins." <i>FEMS Microbiol.</i> , 173(2):303-310 (1999)
AJ010319	ftsY, glnB, glnD, sfp, amtP	Involved in cell division; PII protein; uridylyltransferase (uridylyl-removing enzyme); signal recognition particle; low affinity ammonium uptake protein	
AJ132968	cat	Chloramphenicol acetyl transferase	
AJ224946	mgo	L-malate: quinone oxidoreductase	Molenaar, D. et al. "Biochemical and genetic characterization of the membrane-associated malate dehydrogenase (aceP) from <i>Corynebacterium glutamicum</i> ," <i>Eur. J. Biochem.</i> , 254(2):395-403 (1998)
AJ238250	nsh	NADH dehydrogenase	
AJ238703	porA	Porin	Lichtinger, T. et al. "Biochemical and biophysical characterization of the cell wall porin of <i>Corynebacterium glutamicum</i> : The channel is formed by a low molecular mass polypeptide." <i>Biochemistry</i> , 37(43):15024-15032 (1998)
DI7429		Transposable element IS31831	Veres, A.A. et al. "Isolation and characterization of IS31831, a transposable element from <i>Corynebacterium glutamicum</i> ," <i>Mol. Microbiol.</i> , 11(4):739-746 (1994)

09604231.062700

GenBank™ Accession No.	Gene Name	Gene Function	Reference
D84102	odhA	2-oxoglutarate dehydrogenase	Usuda, Y. et al. "Molecular cloning of the Corynebacterium glutamicum (Brevibacterium lactofermentum A112036) odhA gene encoding a novel type of 2-oxoglutarate dehydrogenase," <i>Microbiology</i> , 142:3347-3354 (1996)
E01358	hcdL, hkc	Homoserine dehydrogenase; homoserine kinase	Katsumata, R. et al. "Production of L-threonine and L-isoleucine," Patent: JP 1987232392-A 1 10/12/87
E01359		Upstream of the start codon of homoserine kinase gene	Katsumata, R. et al. "Production of L-threonine and L-isoleucine," Patent: JP 1987232392-A 2 10/12/87
E01375		Tryptophan operon	
E01376	trpL, trpE	Leader peptide; anthranilate synthase	Matsu, K. et al. "Tryptophan operon, peptide and protein coded thereby, utilization of tryptophan operon gene expression and production of tryptophan," Patent: JP 1987244382-A 1 10/24/87
E01377		Promoter and operator regions of tryptophan operon	Matsu, K. et al. "Tryptophan operon, peptide and protein coded thereby, utilization of tryptophan operon gene expression and production of tryptophan," Patent: JP 1987244382-A 1 10/24/87
E03937		Biotin-synthase	Hatakeyama, K. et al. "DNA fragment containing gene capable of coding biotin synthetase and its utilization," Patent: JP 1992278088-A 1 10/02/92
E04040		Diamino pteralgonic acid aminotransferase	Kohama, K. et al. "Gene coding diaminopteridic acid aminotransferase and deshydrobiotin synthetase and its utilization," Patent: JP 1992330284-A 1 11/18/92
E04041		Deshydrobiotin synthetase	Kohama, K. et al. "Gene coding diaminopteridic acid aminotransferase and deshydrobiotin synthetase and its utilization," Patent: JP 1992330284-A 1 11/18/92
E04307		Flavum aspartase	Kurusu, Y. et al. "Gene DNA coding aspartase and utilization thereof," Patent: JP 1993030977-A 1 02/09/93
E04376		Isochitric acid lyase	Katsumata, R. et al. "Gene manifestation controlling DNA," Patent: JP 1993056782-A 3 03/09/93
E04377		Isochitric acid lyase N-terminal fragment	Katsumata, R. et al. "Gene manifestation controlling DNA," Patent: JP 1993056782-A 3 03/09/93
E04484		Prephenate dehydratase	Sorouchi, N. et al. "Production of L-phenylalanine by fermentation," Patent: JP 1993076332-A 2 03/30/93
E05108		Aspartokinase	Fujino, N. et al. "Gene DNA coding Aspartokinase and its use," Patent: JP 1993184366-A 1 07/27/93
E05112		Dihydro-dipicolinate synthetase	Hatakeyama, K. et al. "Gene DNA coding dihydrodipicolinic acid synthetase and its use," Patent: JP 1993184371-A 1 07/27/93

09604231.062700

GenBank™ Accession No.	Gene Name	Gene Function	Reference
E05776		Diaminopimelic acid dehydrogenase	Kobayashi, M. et al. "Gene DNA coding Diaminopimelic acid dehydrogenase and its use," Patent: JP 1993284970-A 1 11/02/93
E05779		Threonine synthase	Kohama, K. et al. "Gene DNA coding threonine synthase and its use," Patent: JP 1993284972-A 1 11/02/93
E06110		Prephenate dehydratase	Kikuchi, T. et al. "Production of L-phenylalanine by fermentation method," Patent: JP 1993344881-A 1 12/27/93
E06111		Mutated Prephenate dehydratase	Kikuchi, T. et al. "Production of L-phenylalanine by fermentation method," Patent: JP 1993344881-A 1 12/27/93
E06146		Acetyl/droxy acid synthetase	Inui, M. et al. "Gene capable of coding Acetyl/droxy acid synthetase and its use," Patent: JP 199344893-A 1 12/27/93
E06825		Aspartokinase	Sugimoto, M. et al. "Mutant aspartokinase gene," patent: JP 1994062866-A 1 03/08/94
E06826		Mutated aspartokinase alpha subunit	Sugimoto, M. et al. "Mutant aspartokinase gene," patent: JP 1994062866-A 1 03/08/94
E06827		Mutated aspartokinase alpha subunit	Sugimoto, M. et al. "Mutant aspartokinase gene," patent: JP 1994062866-A 1 03/08/94
E07701	secY		Homo, N. et al. "Gene DNA participating in integration of membraneous protein to membrane," Patent: JP 1994169780-A 1 06/21/94
E08177		Aspartokinase	Sato, Y. et al. "Genetic DNA capable of coding Aspartokinase released from feedback inhibition and its utilization," Patent: JP 1994261766-A 1 09/20/94
E08178, E08179, E08180, E08181, E08182		Feedback inhibition-released Aspartokinase	Sato, Y. et al. "Genetic DNA capable of coding Aspartokinase released from feedback inhibition and its utilization," Patent: JP 1994261766-A 1 09/20/94
E08332		Acetyl/droxy-acid isomerase/reductase	Inui, M. et al. "Gene DNA coding acetyl/droxy acid isomerase/reductase," Patent: JP 1994277067-A 1 10/04/94
E08334	secE		Asai, Y. et al. "Gene DNA coding for translocation machinery of protein," Patent: JP 1994277073-A 1 10/04/94
E08643		FT aminotransferase and deshydrobiotin synthetase promoter region	Hatakeyama, K. et al. "DNA fragment having promoter function in Corynebacterium bacterium," Patent: JP 1995031476-A 1 02/03/95
E08646		Biotin synthetase	Hatakeyama, K. et al. "DNA fragment having promoter function in Corynebacterium bacterium," Patent: JP 1995031476-A 1 02/03/95

09604231.062700

GenBank™ Accession No.	Gene Name	Gene Function	Reference
E08649		Aspartase	Kohama, K. et al. "DNA fragment having promoter function in corynebacterium," Patent: JP 1995031478-A 1 02/03/95
E08900		Dihydrodipicolinate reductase	Madori, M. et al. "DNA fragment containing gene coding Dihydrodipicolinate acid reductase and utilization thereof," Patent: JP 1995075578-A 1 03/20/95
E08901		Diaminopimelic acid decarboxylase	Madori, M. et al. "DNA fragment containing gene coding Diaminopimelic acid decarboxylase and utilization thereof," Patent: JP 1995075579-A 1 03/20/95
E12594		Serine hydroxymethyltransferase	Hatakeyama, K. et al. "Production of L-tryptophan," Patent: JP 1997028391-A 1 02/04/97
E12760, E12759, E12758		transposase	Moriya, M. et al. "Amplification of gene using artificial transposon," Patent: JP 1997070291-A 03/18/97
E12764		Arginyl-tRNA synthetase; diaminopimelic acid decarboxylase	Moriya, M. et al. "Amplification of gene using artificial transposon," Patent: JP 1997070291-A 03/18/97
E12767		Dihydrodipicolinic acid synthetase	Moriya, M. et al. "Amplification of gene using artificial transposon," Patent: JP 1997070291-A 03/18/97
E12770		aspartokinase	Moriya, M. et al. "Amplification of gene using artificial transposon," Patent: JP 1997070291-A 03/18/97
E12773		Dihydrodipicolinic acid reductase	Moriya, M. et al. "Amplification of gene using artificial transposon," Patent: JP 1997070291-A 03/18/97
E13655		Glucose-6-phosphate dehydrogenase	Hatakeyama, K. et al. "Glucose-6-phosphate dehydrogenase and DNA capable of coding the same," Patent: JP 199724661-A 1 09/02/97
L01508	ilvA	Threonine dehydratase	Moeschel, B. et al. "Functional and structural analysis of the threonine dehydratase of <i>Corynebacterium glutamicum</i> ," <i>J. Bacteriol.</i> 174:8065-8072 (1992)
L07603	EC 4.2.1.15	3-deoxy-D-arabinotriphosphonate-7-phosphate synthase	Chen, C. et al. "The cloning and nucleotide sequence of <i>Corynebacterium glutamicum</i> 3-deoxy-D-arabinotriphosphonate-7-phosphate synthase gene," <i>FEBS Microbiol. Lett.</i> 107:223-230 (1993)
L09232	ilvB; ilvN; ilvC	Acetylhydroxy acid synthase large subunit; Acetylhydroxy acid synthase small subunit; Acetylhydroxy acid isomerase	Keilhauer, C. et al. "Isolation and structural analysis of the ilvB-ilvN-ilvC operon," <i>J. Bacteriol.</i> 175(17):5595-5603 (1993)

09604231.062709

GenBank™ Accession No.	Gene Name	Gene Function	Reference
L18874	psm	phosphoenolpyruvate sugar phosphotransferase	Fouet, A. et al. "Bacillus subtilis sucrose-specific enzyme II of the phosphotransferase system: expression in <i>Escherichia coli</i> and homology to enzymes II from enteric bacteria," <i>PNAS USA</i> , 84(24):8773-8777 (1987); Lee, J.K. et al. "Nucleotide sequence of the gene encoding the Corynebacterium glutamicum mannose enzyme II and analyses of the deduced protein sequence," <i>FEMS Microbiol. Lett.</i> , 119(1-2):137-145 (1994)
L27123	aceB	Malate synthase	Lee H-S. et al. "Molecular characterization of aceB, a gene encoding malate synthase in <i>Corynebacterium glutamicum</i> ," <i>J. Microbiol. Biotechnol.</i> , 4(4):256-263 (1994)
L27126		Pyruvate kinase	Jetten, M. S. et al. "Structural and functional analysis of pyruvate kinase from <i>Corynebacterium glutamicum</i> ," <i>Appl. Environ. Microbiol.</i> , 60(7):2501-2507 (1994)
L28760	aceA	Isocitrate lyase	Oguzta, J.A. et al. "Molecular cloning, DNA sequence analysis, and characterization of the <i>Corynebacterium diptheriae</i> dxrR from <i>Brevibacterium lactofermentum</i> ," <i>J. Bacteriol.</i> , 177(2):465-467 (1995)
L35906	dxr	Diphtheria toxin repressor	Follert, M. T. et al. "Molecular cloning and nucleotide sequence of the <i>Corynebacterium glutamicum</i> pheA gene," <i>J. Bacteriol.</i> , 167:695-702 (1986)
M13774		Prephenate dehydratase	Park, Y.-H. et al. "Phylogenetic analysis of the <i>coryneform</i> bacteria by 56 rRNA sequences," <i>J. Bacteriol.</i> , 169:1801-1806 (1987)
M16175	5S rRNA		Sano, K. et al. "Structure and function of the trp operon control regions of <i>Brevibacterium lactofermentum</i> , a glutamic-acid-producing bacterium," <i>Gene</i> , 52:191-200 (1987)
M16663	trpE	Anthranilate synthase, 5' end	Sano, K. et al. "Structure and function of the trp operon control regions of <i>Brevibacterium lactofermentum</i> , a glutamic-acid-producing bacterium," <i>Gene</i> , 52:191-200 (1987)
M16664	trpA	Tryptophan synthase, 3' end	O'Regan, M. et al. "Cloning and nucleotide sequence of the Phosphoenolpyruvate carboxylase-coding gene of <i>Corynebacterium glutamicum</i> ATCC13032," <i>Gene</i> , 77(2):237-251 (1989)
M25819		Phosphoenolpyruvate carboxylase	Kotler, C. et al. "Gram-positive bacteria with a high DNA G+C content are characterized by a common insertion within their 23S rRNA genes," <i>J. Gen. Microbiol.</i> , 138:1167-1175 (1992)
M85106		23S rRNA gene insertion sequence	

036041231.062700

GenBank™ Accession No.	Gene Name	Gene Function	Reference
M85107, M85108		23S rRNA gene insertion sequence	Roller, C. et al. "Gram-positive bacteria with a high DNA G+C content are characterized by a common insertion within their 23S rRNA genes." <i>J. Gen. Microbiol.</i> , 138:1167-1175 (1992)
M85931	acdD, bmqQ, ybhW	Beta C-S lyase; branched-chain amino acid uptake carrier; hypothetical protein ybhW	Kosol, I. et al. "The Corynebacterium glutamicum acdD gene encodes a C-S lyase with alpha, beta-elimination activity that degrades aminocyclohexane." <i>J. Bacteriol.</i> , 174(9):2968-2977 (1992); Jauch, A. et al. "Isolucine uptake in Corynebacterium glutamicum ATCC 13032 is directed by the bmqQ gene product." <i>Arch. Microbiol.</i> , 169(4):303-312 (1998)
S59299	trp	Leader gene (promoter)	Herry, D.M. et al. "Cloning of the trp gene cluster from a tryptophan-hyperproducing strain of Corynebacterium glutamicum: identification of a mutation in the trp leader sequence." <i>Appl. Environ. Microbiol.</i> , 59(3):791-799 (1993)
U11545	trpD	Anthranilate phosphoribosyltransferase	O'Gara, J.P. and Dunne, L.K. (1994) Complete nucleotide sequence of the Corynebacterium glutamicum ATCC 21850 pD gene." Thesis, Microbiology Department, University College Galway, Ireland.
U13922	cgIIIM; cgIIIR; cglIIIR	Putative type II 5-cytosine methylesterase; putative type II restriction endonuclease; putative type I or type III restriction endonuclease	Schäfer, A. et al. "Cloning and characterization of a DNA region encoding a stress-sensitive restriction system from Corynebacterium glutamicum ATCC 13032 and analysis of its role in intergeneric conjugation with Escherichia coli." <i>J. Bacteriol.</i> , 176(23):7309-7319 (1994); Schäfer, A. et al. "The Corynebacterium glutamicum cglIIIM gene encoding a 5-cytosine in an MCRBC-deficient Escherichia coli strain." <i>Gene</i> , 203(2):95-101 (1997)
U14965	recA		Ankri, S. et al. "Mutations in the Corynebacterium glutamicum proline biosynthetic pathway: A natural bypass of the proA step." <i>J. Bacteriol.</i> , 178(15):4412-4419 (1996)
U31224	ppx		Ankri, S. et al. "Mutations in the Corynebacterium glutamicum proline biosynthetic pathway: A natural bypass of the proA step." <i>J. Bacteriol.</i> , 178(15):4412-4419 (1996)
U31225	proC	L-proline: NAD(P)+5-oxido-reductase	Ankri, S. et al. "Mutations in the Corynebacterium glutamicum proline biosynthetic pathway: A natural bypass of the proA step." <i>J. Bacteriol.</i> , 178(15):4412-4419 (1996)
U31230	obg, proB, unkdh	γ-gamma glutamyl kinase; similar to D-isomer specific 2-hydroxyacid dehydrogenases	Ankri, S. et al. "Mutations in the Corynebacterium glutamicum proline biosynthetic pathway: A natural bypass of the proA step." <i>J. Bacteriol.</i> , 178(15):4412-4419 (1996)

GenBank™ Accession No.	Gene Name	Gene Function	Reference
U31281	bioB	Biotin synthase	SerchrisKil, I.G., "Two new members of the bio B superfamily: Cloning, sequencing and expression of bio B genes of <i>Methylobacillus flagellatum</i> and <i>Corynebacterium glutamicum</i> ," <i>Gene</i> , 175:15-22 (1996)
U35023	thrR, accBC	Thiosulfate sulfurtransferase, acyl CoA carboxylase	Jager, W. et al. "A <i>Corynebacterium glutamicum</i> gene encoding a two-domain protein similar to biotin carboxylases and biotin-carboxyl-carrier proteins," <i>Arch. Microbiol.</i> , 166(2):76-82 (1996)
U43535	cmr	Multidrug resistance protein	Jager, W. et al. "A <i>Corynebacterium glutamicum</i> gene conferring multidrug resistance in the heterologous host <i>Escherichia coli</i> ," <i>J. Bacteriol.</i> , 179(7):2449-2451 (1997)
U43536	cjbB	Heat shock ATP-binding protein	
U53587	aphA-3	3',5'-aminoglycoside phosphotransferase	
U89648		<i>Corynebacterium glutamicum</i> unidentified sequence involved in histidine biosynthesis, partial sequence	
X04960	trpA; trpB; trpC; trpD; trpE; trpG; trpL	Tryptophan operon	Matsu, K. et al. "Complete nucleotide and deduced amino acid sequences of the <i>Brevibacterium lactofermentum</i> tryptophan operon," <i>Nucleic Acids Res.</i> , 14(24):10113-10114 (1986)
X07563	lys A	DAP decarboxylase (meso-diaminopimelate decarboxylase, EC 4.1.1.20)	Yeh, P. et al. "Nucleic sequence of the <i>lysA</i> gene of <i>Corynebacterium glutamicum</i> and possible mechanisms for modulation of its expression," <i>Mol. Gen. Genet.</i> , 212(1):112-119 (1988)
X14234	EC 4.1.1.31	Phosphoenolpyruvate carboxylase	Ekman, B.J. et al. "The Phosphoenolpyruvate carboxylase gene of <i>Corynebacterium glutamicum</i> : Molecular cloning, nucleotide sequence, and expression," <i>Mol. Gen. Genet.</i> , 218(2):330-339 (1989); Lepoint, L. et al. "Sorghum Phosphoenolpyruvate carboxylase gene family: structure, function and molecular evolution," <i>Plant. Mol. Biol.</i> , 21 (3):487-502 (1993)
X17313	fda	Fructose-bisphosphate aldolase	Von der Osten, C.H. et al. "Molecular cloning, nucleotide sequence and fine-structural analysis of the <i>Corynebacterium glutamicum</i> fda gene: structural comparison of <i>C. glutamicum</i> fructose-1, 6-bisphosphate aldolase to class I and class II aldolases," <i>Mol. Microbiol.</i>
X53993	dapA	L-2, 3-dihydrodipicolinate synthetase (EC 4.2.1.52)	Bonmassé, S. et al. "Nucleic sequence of the dapA gene from <i>Corynebacterium glutamicum</i> ," <i>Nucleic Acids Res.</i> , 18(21):6421 (1990)

09604231.062700

GenBank™ Accession No.	Gene Name	Gene Function	Reference
X54223		AuB-related site	Chianfroto, N. et al. "DNA sequence homology between att B-related sites of Corynebacterium diptheriae, Corynebacterium ulcerans, Corynebacterium glutamicum, and the attP site of lambdacytophage," <i>FEMS Microbiol. Lett.</i> , 66:299-302 (1990)
X54740	argS, lysA	Arginyl-tRNA synthetase; Diaminopimelate decarboxylase	Mazel, T. et al. "Nucleotide sequence and organization of the upstream region of the Corynebacterium glutamicum lysA gene," <i>Mol. Microbiol.</i> , 4(11):1819-1830 (1990)
X55994	trpL, trpE	Purative leader peptide; anthranilate synthase component 1	Heery, D.M. et al. "Nucleotide sequence of the Corynebacterium glutamicum trpE gene," <i>Nucleic Acids Res.</i> , 18(23):7138 (1990)
X56037	thrC	Threonine synthase	Han, K. S. et al. "The molecular structure of the Corynebacterium glutamicum threonine synthase gene," <i>Mol. Microbiol.</i> , 4(10):1693-1702 (1990)
X56075	attB-related site	Attachment site	Cunetto, N. et al. "DNA sequence homology between att B-related sites of Corynebacterium diptheriae, Corynebacterium ulcerans, Corynebacterium glutamicum, and the attP site of lambdacytophage," <i>FEMS Microbiol. Lett.</i> , 66:299-302 (1990)
X57226	lysC-alpha; lysC-beta; asd	Aspartokinase-alpha subunit; aspartate beta semialdehyde dehydrogenase	Kalinowski, J. et al. "Genetic and biochemical analysis of the Aspartokinase from Corynebacterium glutamicum," <i>Mol. Microbiol.</i> , 5(5):1197-1204 (1991); Kalinowski, J. et al. "Aspartokinase genes lysC alpha and lysC beta overlap and are adjacent to the aspartate beta-semialdehyde dehydrogenase gene asd in Corynebacterium glutamicum," <i>Mol. Gen. Gene.</i> , 224(3):317-324 (1990)
X59403	gap;pgk; tpi	Glycerinaldehyde-3-phosphate; phosphoglycerate kinase; triosephosphate isomerase	Ekman, B.J. "Identification, sequence analysis, and expression of a Corynebacterium glutamicum gene cluster encoding the three glycolytic enzymes glyceraldehyde-3-phosphate dehydrogenase, 3-phosphoglycerate kinase, and triosephosphate isomerase," <i>J. Bacteriol.</i> , 174(19):6076-6086 (1992)
X59404	gdh	Glutamate dehydrogenase	Bormann, E.R. et al. "Molecular analysis of the Corynebacterium glutamicum gdh gene encoding glutamate dehydrogenase," <i>Mol. Microbiol.</i> , 6(3):317-326 (1992)
X60312	lysI	L-lysine permease	Seep-Feldhaus, A.H. et al. "Molecular analysis of the Corynebacterium glutamicum lysI gene involved in lysine uptake," <i>Mol. Microbiol.</i> , 5(12):2995-3005 (1991)

09604231.062700

GenBank™ Accession No.	Gene Name	Gene Function	Reference
X66078	copI	PSI protein	Jolliff, G. et al. "Cloning and nucleotide sequence of the copI gene encoding PSI, one of the two major secreted proteins of <i>Corynebacterium glutamicum</i> : The deduced N-terminal region of PSI is similar to the Mycobacterium antigen 85 complex," <i>Mol. Microbiol.</i> , 6(16):2349-2362 (1992)
X66112	glt	Citrate synthase	Ekman, B.J. et al. "Cloning sequence, expression and transcriptional analysis of the <i>Corynebacterium glutamicum</i> gltA gene encoding citrate synthase," <i>Microbiol.</i> , 140:1817-1828 (1994)
X66737	dapB	Dihydrodipicolinate reductase	Peyret, J.L. et al. "Characterization of the espB gene encoding PS2, an ordered surface-layer protein in <i>Corynebacterium glutamicum</i> ," <i>Mol. Microbiol.</i> , 9(1):97-109 (1993)
X69103	esp2	Surface layer protein PS2	Bonamy, C. et al. "Identification of IS1206, a <i>Corynebacterium glutamicum</i> IS3-related insertion sequence and phylogenetic analysis," <i>Mol. Microbiol.</i> , 14(3):571-581 (1994)
X69104		IS3 related insertion element	Patek, M. et al. "Leucine synthetase in <i>Corynebacterium glutamicum</i> : enzyme activities, structure of leuA, and effect of leuA inactivation on lysine synthesis," <i>Appl. Environ. Microbiol.</i> , 60(1):133-140 (1994)
X70959	leuA	Isopropylmalate synthase	Ekman, B.J. et al. "Cloning sequence analysis, expression, and inactivation of the <i>Corynebacterium glutamicum</i> leuA gene encoding isocitrate dehydrogenase and biochemical characterization of the enzyme," <i>J. Bacteriol.</i> , 177(3):774-782 (1995)
X71489	icd	Isocitrate dehydrogenase (NADP+)	
X72855	GDH A	Glutamate dehydrogenase (NADP+)	
X75083, X70584	mraA	5-methyltryptophan resistance	Heery, D.M. et al. "A sequence from a tryptophan-hyperproducing strain of <i>Corynebacterium glutamicum</i> encoding resistance to 5-methyltryptophan," <i>Biochem. Biophys. Res. Commun.</i> , 201(3):1255-1262 (1994)
X75085	recA		Fitzpatrick, R. et al. "Construction and characterization of recA mutant strains of <i>Corynebacterium glutamicum</i> and <i>Brevibacterium lactofermentum</i> ," <i>Appl. Microbiol. Biotechnol.</i> , 42(4):575-580 (1994)
X75504	aceA; thtX	Partial Isocitrate lyase; ?	Reischeld, D.J. et al. "Characterization of the isocitrate lyase gene from <i>Corynebacterium glutamicum</i> and biochemical analysis of the enzyme," <i>J. Bacteriol.</i> , 176(12):3474-3483 (1994)
X76875		ATPase beta-subunit	Ludwig, W. et al. "Phylogenetic relationships of bacteria based on comparative sequence analysis of elongation factor Tu and ATP-synthase beta-subunit genes," <i>Antonie Van Leeuwenhoek</i> , 64:285-305 (1993)

09604231.062700

GenBank TM Accession No.	Gene Name	Gene Function	Reference
X77034	tuf	Elongation factor Tu	Ludwig, W. et al. "Phylogenetic relationships of bacteria based on comparative sequence analysis of elongation factor Tu and ATP-synthase beta-subunit genes." <i>Antonie Van Leeuwenhoek</i> , 64:285-305 (1993)
X77384	recA		Billman-Jacobe, H. "Nucleotide sequence of a recA gene from <i>Corynebacterium glutamicum</i> ." <i>DNA Seq.</i> 4(6):403-404 (1994)
X78491	acb3	Malate synthase	Reinscheid, D.J. et al. "Malate synthase from <i>Corynebacterium glutamicum</i> pta-ack operon encoding phosphotransacetylase: sequence analysis." <i>Microbiology</i> , 140:3099-3108 (1994)
X80629	16S rDNA	16S ribosomal RNA	Ratney, F. A. et al. "Phylogenetic analysis of the genera <i>Rhodococcus</i> and <i>Norcardia</i> and evidence for the evolutionary origin of the genus <i>Norcardia</i> from within the radiation of <i>Rhodococcus</i> species." <i>Microbiol.</i> , 141:523-528 (1995)
X81191	gluA ₁ , gltB ₁ , gltC ₁ , gltD	Glutamate uptake system	Kronmeyer, W. et al. "Structure of the gluABCD cluster encoding the glutamate uptake system of <i>Corynebacterium glutamicum</i> ." <i>J. Bacteriol.</i> , 177(5):1152-1158 (1995)
X81379	dapE	Succinylidiaminopimelate desuccinylase	Wehrmann, A. et al. "Analysis of different DNA fragments of <i>Corynebacterium glutamicum</i> complementing dapE of <i>Escherichia coli</i> ." <i>Microbiology</i> , 40:3349-56 (1994)
X82061	16S rDNA	16S ribosomal RNA	Ratney, F. et al. "Phylogeny of the genus <i>Corynebacterium</i> deduced from analyses of small-subunit ribosomal DNA sequences." <i>Int. J. Syst. Bacteriol.</i> , 43(4):740-746 (1995)
X82928	asd; lysC	Aspartate-semialdehyde dehydrogenase; ?	Scherjiski, I. et al. "Multicopy suppression by asd gene and osmotic stress-dependent complementation by heterologous proA in proA mutants." <i>J. Bacteriol.</i> , 177(24):7255-7260 (1995)
X82929	proA	Gamma-glutamyl phosphate reductase	Scherjiski, I. et al. "Multicopy suppression by asd gene and osmotic stress-dependent complementation by heterologous proA in proA mutants." <i>J. Bacteriol.</i> , 177(24):7255-7260 (1995)
X84257	16S rDNA	16S ribosomal RNA	Pastcal, C. et al. "Phylogenetic analysis of the genus <i>Corynebacterium</i> based on 16S rRNA gene sequences." <i>Int. J. Syst. Bacteriol.</i> , 45(4):724-728 (1995)
X85965	arop; dapE	Aromatic amino acid permease; ?	Wehrmann, A. et al. "Functional analysis of sequences adjacent to dapE of <i>Corynebacterium glutamicum</i> reveals the presence of arp, which encodes the aromatic amino acid transporter." <i>J. Bacteriol.</i> , 177(20):5991-5995 (1995)

09604231.062700

GenBank™ Accession No.	Gene Name	Gene Function	Reference
X86157	argH; argC; argD; argF; argJ	Acetylglutamate kinase; N-acetyl-l-glutamyl-l-phosphate reductase; acetylornithine aminotransferase; ornithine carbamoyltransferase; glutamate N-acetyltransferase	Sakanyan, V. et al. "Genes and enzymes of the acetyl cycle of arginine biosynthesis in <i>Corynebacterium glutamicum</i> : enzyme evolution in the early steps of the arginine pathway," <i>Microbiology</i> , 142:99-108 (1996)
X89084	pta; ackA	Phosphate acetyltransferase; acetate kinase	Reinscheid, D.J. et al. "Cloning, sequence analysis, expression and inactivation of the <i>Corynebacterium glutamicum</i> pta-ack operon encoding phosphotransacetylase and acetate kinase," <i>Microbiology</i> , 143:503-513 (1999)
X89850	attB	Attachment site	Le Marrec, C. et al. "Genetic characterization of site-specific integration functions of phi AAU2 infecting " <i>Athrobacter aureus</i> C70," <i>J. Bacteriol.</i> , 178(7):1996-2004 (1996)
X90356		Promoter fragment F1	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90357		Promoter fragment F2	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90358		Promoter fragment F10	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90359		Promoter fragment F13	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90360		Promoter fragment F22	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90361		Promoter fragment F34	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90362		Promoter fragment F37	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)

09604231.062700

GenBank™ Accession No.	Gene Name	Gene Function	Reference
X90363		Promoter fragment F45	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90364		Promoter fragment F64	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90365		Promoter fragment F75	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90366		Promoter fragment PF101	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90367		Promoter fragment PF104	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90368		Promoter fragment PF109	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X93513	amt	Ammonium transport system	Stewe, R.M. et al. "Functional and genetic characterization of the (methy) ammonium uptake carrier of <i>Corynebacterium glutamicum</i> ," <i>J. Biol. Chem.</i> , 271(10):5398-5403 (1996)
X93514	betP	Glycine betaine transport system	Peter, H. et al. "Isolation, characterization, and expression of the <i>Corynebacterium glutamicum</i> betP gene, encoding the transport system for the compatible solute glycine betaine," <i>J. Bacteriol.</i> , 178(17):5229-5234 (1996)
X95649	orf4		Patek, M. et al. "Identification and transcriptional analysis of the dapA-ORF4-dapA-ORF4 operon of <i>Corynebacterium glutamicum</i> , encoding two enzymes involved in L-lysine synthesis," <i>Biochem. Lett.</i> , 19:1131-1137 (1997)
X96471	lysE, lysG	Lysine exporter protein; Lysine export regulator protein	Vrjlic, M. et al. "A new type of transporter with a new type of cellular function: L-lysine export from <i>Corynebacterium glutamicum</i> ," <i>Mol. Microbiol.</i> , 22(5):815-826 (1996)

09604231.062700

GenBank™ Accession No.	Gene Name	Gene Function	Reference
X96580	panB; panC; xyfB	3-methyl-2-oxobutanoate hydroxymethyltransferase; pantoate-beta-alanine ligase; xylokinase	Sahn, H. et al., "D-pantoic acid synthesis in <i>Corynebacterium glutamicum</i> and use of panB and genes encoding L-valine synthesis for D-pantoic acid overproduction," <i>Appl. Environ. Microbiol.</i> , 65(5):1973-1979 (1999)
X96662		Insertion sequence [S]207 and transposase	
X99289		Elongation factor P	Ramos, A. et al., "Cloning, sequencing and expression of the gene encoding elongation factor P in the amino acid producer <i>Brevibacterium lactofermentum</i> (<i>Corynebacterium glutamicum</i> ATCC 13869)," <i>Gene</i> , 198:217-222 (1997)
Y00140	thrB	Homoserine kinase	Matéos, L.M. et al., "Nucleotide sequence of the homoserine kinase (thrB) gene of the <i>Brevibacterium lactofermentum</i> ," <i>Nucleic Acids Res.</i> , 15(9):3922 (1987)
Y00151	ddh	Meso-diaminopimelate D-dehydrogenase (EC 1.4.1.16)	Ishino, S. et al., "Nucleotide sequence of the meso-diaminopimelate D-dehydrogenase gene from <i>Corynebacterium glutamicum</i> ," <i>Nucleic Acids Res.</i> , 15(9):3917 (1987)
Y00476	thrA	Homoserine dehydrogenase	Matéos, L.M. et al., "Nucleotide sequence of the homoserine dehydrogenase (thrA) gene of the <i>Brevibacterium lactofermentum</i> ," <i>Nucleic Acids Res.</i> , 15(24):10598 (1987)
Y00546	hom; thrH	Homoserine dehydrogenase; homoserine kinase	Peoples, O.P. et al., "Nucleotide sequence and fine structural analysis of the <i>Corynebacterium glutamicum</i> hom-thrB operon," <i>Mol. Microbiol.</i> , 2(1):65-72 (1988)
Y08964	murC; ftsQ/divD; ftsZ	UDP-N-acetylglutamate-amine ligase; division initiation protein or cell division protein; cell division system	Homrhub, M.P. et al., "Identification, characterization, and chromosomal organization of the ftsZ gene from <i>Brevibacterium lactofermentum</i> ," <i>Mol. Gen. Genet.</i> , 259(1):97-104 (1998)
Y09163	putP	High affinity proline transport system	Peter, H. et al., "Isolation of the putP gene of <i>Corynebacterium glutamicum</i> and characterization of a low-affinity uptake system for compatible solutes," <i>Arch. Microbiol.</i> , 168(2):143-151 (1997)
Y09548	pyc	Pyruvate carboxylase	Peters-Wendisch, P.G. et al., "Pyruvate carboxylase from <i>Corynebacterium glutamicum</i> : characterization, expression and inactivation of the pyc gene," <i>Microbiology</i> , 144(9):5427 (1998)
Y09578	leuB	3-isopropylmalate dehydrogenase	Patek, M. et al., "Analysis of the leuB gene from <i>Corynebacterium glutamicum</i> ," <i>Appl. Microbiol. Biotechnol.</i> , 50(1):42-47 (1998)
Y12472		Attachment site bacteriophage Phi-16	Moreau, S. et al., "Site-specific integration of coryneophage Phi-16: The construction of an integration vector," <i>Microbiol.</i> , 145:539-548 (1999)

09604233.062700

GenBank™ Accession No.	Gene Name	Gene Function	Reference
Y12537	proP	Proline/ectoine uptake system protein	Peter, H. et al. "Corynebacterium glutamicum is equipped with four secondary carriers for compatible solutes: Identification, sequencing, and characterization of the proline/ectoine uptake system, ProP, and the ectoine/proline/glycine betaine carrier, EcPp." <i>J. Bacteriol.</i> , 180(22):6005-6012 (1998)
Y13221	gluA	Glutamine synthetase I	Jakoby, M. et al. "Isolation of <i>Corynebacterium glutamicum</i> gluA gene encoding glutamine synthetase I." <i>FEMS Microbiol. Lett.</i> , 154(1):81-88 (1997)
Y16642	lpd	Dihydrodipicolinate dehydrogenase	
Y18059		Attachment site Corynephage 304L	Moreau, S. et al. "Analysis of the integration functions of ϕ h3:304L: An integrase module among corynephages." <i>Virology</i> , 255(1):150-159 (1999)
Z21501	argS; lysA	Arginyl-tRNA synthetase; diamminopimelate decarboxylase (partial)	Oguzta, J.A. et al. "A gene encoding arginyl-tRNA synthetase is located in the upstream region of the lysA gene in <i>Brevibacterium lactofermentum</i> : Regulation of argS-lysA cluster expression by arginine." <i>J. Bacteriol.</i> , 175(22):7356-7362 (1993)
Z21502	dapA; dapB	Dihydrodipicolinate synthase; dihydrodipicolinate reductase	Pisabarro, A. et al. "A cluster of three genes (dapA, orf2, and dapB) of <i>Brevibacterium lactofermentum</i> encodes dihydrodipicolinate reductase, and a third polypeptide of unknown function." <i>J. Bacteriol.</i> , 173(9):2743-2749 (1993)
Z29563	thtC	Threonine synthase	Malumbres, M. et al. "Analysis and expression of the thtC gene of the encoded threonine synthase." <i>Appl. Environ. Microbiol.</i> , 60(7):2209-2219 (1994)
Z46753	16S rDNA	Gene for 16S ribosomal RNA	
Z49822	sigA	SigA sigma factor	Oguzta, J.A. et al. "Multiple sigma factor genes in <i>Brevibacterium lactofermentum</i> : Characterization of sigA and sigB." <i>J. Bacteriol.</i> , 178(2):550-553 (1996)
Z49823	galE; dtxR	Catalytic activity UDP-galactose 4-epimerase; diphtheria toxin regulatory protein	Oguzta, J.A. et al. "The galE gene encoding the UDP-galactose 4-epimerase of <i>Brevibacterium lactofermentum</i> is coupled transcriptionally to the dtxR gene." <i>Gene</i> , 177:103-107 (1996)
Z49824	orf1; sigB	?; SigB sigma factor	Oguzta, J.A. et al. "Multiple sigma factor genes in <i>Brevibacterium lactofermentum</i> : Characterization of sigA and sigB." <i>J. Bacteriol.</i> , 178(2):550-553 (1996)
Z66534		Transposase	Correia, A. et al. "Cloning and characterization of an IS-like element present in the genome of <i>Brevibacterium lactofermentum</i> ATCC 13869." <i>Gene</i> , 170(1):91-94 (1996)

A sequence for this gene was published in the indicated reference. However, the sequence obtained by the inventors of the present application is significantly longer than the published version. It is believed that the published version relied on an incorrect start codon, and thus represents only a fragment of the actual coding region.

09604231.062700

TABLE 3: Corynebacterium and Brevibacterium Strains Which May be Used in the Practice of the Invention

Genus	Species	ATCC	FIRM	NRRL	DECT	NCIMB	CBS	NCIC	DSMZ
Brevibacterium	ammoniaenes	21054							
Brevibacterium	ammoniaenes	19350							
Brevibacterium	ammoniaenes	19351							
Brevibacterium	ammoniaenes	19352							
Brevibacterium	ammoniaenes	19353							
Brevibacterium	ammoniaenes	19354							
Brevibacterium	ammoniaenes	19355							
Brevibacterium	ammoniaenes	19356							
Brevibacterium	ammoniaenes	21055							
Brevibacterium	ammoniaenes	21077							
Brevibacterium	ammoniaenes	21553							
Brevibacterium	ammoniaenes	21580							
Brevibacterium	ammoniaenes	39101							
Brevibacterium	butanicum	21196							
Brevibacterium	divaricatum	21792	P928						
Brevibacterium	flavum	21474							
Brevibacterium	flavum	21129							
Brevibacterium	flavum	21518							
Brevibacterium	flavum			B11474					
Brevibacterium	flavum			B11472					
Brevibacterium	flavum	21127							
Brevibacterium	flavum	21128							
Brevibacterium	flavum	21427							
Brevibacterium	flavum	21475							
Brevibacterium	flavum	21517							
Brevibacterium	flavum	21528							
Brevibacterium	flavum	21529							

03604231.062700

Brevibacterium	flavum			B11477					
Brevibacterium	flavum			B11478					
Brevibacterium	flavum	21127							
Brevibacterium	flavum			B11474					
Brevibacterium	hawaii	15527							
Brevibacterium	ketoglutaricum	21004							
Brevibacterium	ketoglutaricum	21089							
Brevibacterium	ketoreducticum	21914							
Brevibacterium	lactofermentum			70					
Brevibacterium	lactofermentum			74					
Brevibacterium	lactofermentum			77					
Brevibacterium	lactofermentum	21798							
Brevibacterium	lactofermentum	21799							
Brevibacterium	lactofermentum	21800							
Brevibacterium	lactofermentum	21801							
Brevibacterium	lactofermentum			B11470					
Brevibacterium	lactofermentum			B11471					
Brevibacterium	lactofermentum	21086							
Brevibacterium	lactofermentum	21420							
Brevibacterium	lactofermentum	21086							
Brevibacterium	lactofermentum	31269							
Brevibacterium	lactofermentum	9174							
Brevibacterium	linens	19391							
Brevibacterium	linens	8377							
Brevibacterium	paraffinolyticum					11160			
Brevibacterium	spec.					717.73			
Brevibacterium	spec.	14604				717.73			
Brevibacterium	spec.	21860							
Brevibacterium	spec.	21864							
Brevibacterium	spec.	21865							

09604231.062700

Brevibacterium	spec.	21866							
Brevibacterium	spec.	19240							
Corynebacterium	acetoacidophilum	21476							
Corynebacterium	acetoacidophilum	13870							
Corynebacterium	aceto-glutamicum			B11473					
Corynebacterium	aceto-glutamicum			B11475					
Corynebacterium	aceto-glutamicum	15806							
Corynebacterium	aceto-glutamicum	21491							
Corynebacterium	aceto-glutamicum	31270							
Corynebacterium	acetophilum			B3671					
Corynebacterium	ammonia-ferens	6872							
Corynebacterium	ammonia-ferens	15511						2399	
Corynebacterium	fujikense	21496							
Corynebacterium	glutamicum	14067							
Corynebacterium	glutamicum	39137							
Corynebacterium	glutamicum	21254							
Corynebacterium	glutamicum	21255							
Corynebacterium	glutamicum	31830							
Corynebacterium	glutamicum	13032							
Corynebacterium	glutamicum	15455							
Corynebacterium	glutamicum	14305							
Corynebacterium	glutamicum	13058							
Corynebacterium	glutamicum	13059							
Corynebacterium	glutamicum	13060							
Corynebacterium	glutamicum	21492							
Corynebacterium	glutamicum	21513							
Corynebacterium	glutamicum	21526							
Corynebacterium	glutamicum	21543							
Corynebacterium	glutamicum	13287							
Corynebacterium	glutamicum	21851							
Corynebacterium	glutamicum	21253							

09604231.062700

Corynebacterium	glutamicum	21514																	
Corynebacterium	glutamicum	21516																	
Corynebacterium	glutamicum	21299																	
Corynebacterium	glutamicum	21300																	
Corynebacterium	glutamicum	39684																	
Corynebacterium	glutamicum	21488																	
Corynebacterium	glutamicum	21649																	
Corynebacterium	glutamicum	21650																	
Corynebacterium	glutamicum	19223																	
Corynebacterium	glutamicum	13869																	
Corynebacterium	glutamicum	21157																	
Corynebacterium	glutamicum	21158																	
Corynebacterium	glutamicum	21159																	
Corynebacterium	glutamicum	21355																	
Corynebacterium	glutamicum	31808																	
Corynebacterium	glutamicum	21674																	
Corynebacterium	glutamicum	21562																	
Corynebacterium	glutamicum	21563																	
Corynebacterium	glutamicum	21564																	
Corynebacterium	glutamicum	21565																	
Corynebacterium	glutamicum	21566																	
Corynebacterium	glutamicum	21567																	
Corynebacterium	glutamicum	21568																	
Corynebacterium	glutamicum	21569																	
Corynebacterium	glutamicum	21570																	
Corynebacterium	glutamicum	21571																	
Corynebacterium	glutamicum	21572																	
Corynebacterium	glutamicum	21573																	
Corynebacterium	glutamicum	21579																	
Corynebacterium	glutamicum	19049																	
Corynebacterium	glutamicum	19050																	

09604231.062700

Corynebacterium	glutamicum	19051							
Corynebacterium	glutamicum	19052							
Corynebacterium	glutamicum	19053							
Corynebacterium	glutamicum	19054							
Corynebacterium	glutamicum	19055							
Corynebacterium	glutamicum	19056							
Corynebacterium	glutamicum	19057							
Corynebacterium	glutamicum	19058							
Corynebacterium	glutamicum	19059							
Corynebacterium	glutamicum	19060							
Corynebacterium	glutamicum	19185							
Corynebacterium	glutamicum	13286							
Corynebacterium	glutamicum	21515							
Corynebacterium	glutamicum	21527							
Corynebacterium	glutamicum	21544							
Corynebacterium	glutamicum	21492							
Corynebacterium	glutamicum			B8183					
Corynebacterium	glutamicum			B8182					
Corynebacterium	glutamicum			B12416					
Corynebacterium	glutamicum			B12417					
Corynebacterium	glutamicum			B12418					
Corynebacterium	glutamicum			B11476					
Corynebacterium	glutamicum	21608							
Corynebacterium	lilium		P973						
Corynebacterium	nitrilophilus	21419					11594		
Corynebacterium	spec.		P4445						
Corynebacterium	spec.		P4446						
Corynebacterium	spec.	31088							
Corynebacterium	spec.	31089							
Corynebacterium	spec.	31090							
Corynebacterium	spec.	31090							

036044231_062700

Corynebacterium spec.	31090						
Corynebacterium spec.	15954						2045
Corynebacterium spec.	21857						
Corynebacterium spec.	21862						
Corynebacterium spec.	21863						

ATCC: American Type Culture Collection, Rockville, MD, USA

FERM: Fermentation Research Institute, Chiba, Japan

NRRL: ARS Culture Collection, Northern Regional Research Laboratory, Peoria, IL, USA

CECT: Coleccion Espanola de Cultivos Tipo, Valencia, Spain

NCIMB: National Collection of Industrial and Marine Bacteria Ltd., Aberdeen, UK

CBS: Centraalbureau voor Schimmelmcultures, Baarn, NL

NCTC: National Collection of Type Cultures, London, UK

DSMZ: Deutsche Sammlung von Mikroorganismen und Zellkulturen, Braunschweig, Germany

For reference see Sugaawara, H. et al. (1993) World directory of collections of cultures of microorganisms: Bacteria, fungi and yeasts (4th edn). World federation for culture collections world data center on microorganisms, Saitama, Japan.

TABLE 4: ALIGNMENT RESULTS

ID #	length NT	Genbank Hit	Length	Accession	Name of Genbank Hit	Source of Genbank Hit	% homology (GAP)	Date of Deposit
na00315	1527	GB_BA1.AB007125 GB_M1.DEC49D2	4078 17361	AB007125 U64861	<i>Serratia marcescens</i> salk1 gene for surface layer protein, complete cds, isolate 8000 <i>Caenorhabditis elegans</i> osm-2 Caenorhabditis elegans clone 732G3 *** SEQUENCING IN PROGRESS *** 9 unrelated clones.	<i>Serratia marcescens</i> <i>Caenorhabditis elegans</i> <i>Caenorhabditis elegans</i>	40.366 36.007 36.428	26-MAR-1998 28-JUL-96 23-Feb-99
na01503	3172	GB_M1GC.A0368732 GB_M1GC.A0305019	156453 183352	AC036732 AC030519	Human sapiens BAC clone G5250A.6 from 7p21-p22, complete sequence. RP0211-15(GS12) RP0211-11 Homo sapiens genomic clone RP0211-11- 15(G) genomic survey sequence.	Human sapiens Human sapiens	39.137 43.137	21-MAY-1999
na01298	2187	GB_GSS512.A0293040 GB_GSS512.A0293040 GB_GSS512.A0293040	690 542 542	AC039040 AC029423 AC029423	HP3_308_281_C10_17C C1 Approved Human Sperm Library D Homo sapiens genomic clone Hs308-281-C10-17C-Rover genomic survey sequence Hs308-281-C10-17C-11 NH ₂ B1AP ₂ _M_S2 Mus musculus cDNA clone U44k-B1 B1-salm-03-D-U-3 mRNA sequence	Human sapiens Mus musculus	37.643 41.475	3-Aug-99 18-Sep-99
na01298	2187	GB_EST18.AM047296	614	AM047296	Rattus norvegicus mRNA for Bmi-1-class homeodomain transcription factor, complete cds	Rattus norvegicus	41.031	2-Sep-98
na01298	2187	GB_EST18.AM047296	614	AB004056	Rattus norvegicus mRNA for Bmi-1-class homeodomain transcription factor, complete cds	Rattus norvegicus	40.717	2-Sep-98
na00851	4176	GB_P01.A030456 GB_P01.A030456 GB_P01.A030456	1581 1581 1581	AB004056 AB004056 AB004056	Streptococcus coelicolor strain J21 Mouse hypoxanthine phosphoribosyl transferase (hprt) cDNA, replication protein, glycoprotein, DNA replication protein, FLICE-inhibitory protein and V-protein genes, complete cds, and tegument protein gene, partial cds	Streptococcus coelicolor Mus musculus	34.513 40.097	5-Aug-99 04-DEC-1999
na01244	1827	GB_BA1.AFAPBHI GB_P03.HS835E13	4501 7805	M85036 AL050326	<i>Alcaligenes eutrophus</i> npten H (gntH) and protein I (gntI) genes, complete cds Human DNA sequences from clone 830E13 on chromosome 20 Contains ESTs, EST16162 Normalized rat lung Banno Soares Rattus sp. cDNA clone RUCF56 3' end, mRNA sequence. Homo sapiens deoxyribonucleic acid kinase gene, complete cds Sequence 1 from patient US 5622351 Sequence 2 from patient US 5622351	<i>Alcaligenes eutrophus</i> Homo sapiens	45.624 37.303	26-Apr-93 23-Nov-99
na01300	390	GB_P03.HS835E13 GB_P03.HS835E13 GB_P03.HS835E13	2674 2674 2674	AL102027 L38674 L38674	Human sapiens deoxyribonucleic acid kinase gene, complete cds Sequence 1 from patient US 5622351 Sequence 2 from patient US 5622351	Human sapiens	37.644 37.644 37.644	20-Jan-99 11-Aug-98 13-MAY-1997
na01843	2172	GB_BA1.BRLP1SG GB_BA1.BRLP1SG GB_BA1.BRLP1SG	2856 3163 2841	AF045481 AF045481 AF045481	<i>Corynebacterium ammoniagenes</i> glucose permease (glsG) gene, complete cds	<i>Corynebacterium ammoniagenes</i>	53.658	29-Jul-98

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re the application of: Markus Pompejus *et al.*

Serial No.: Not Yet Assigned

Filed: Herewith

For: "*Corynebacterium Glutamicum Genes Encoding Phosphoenolpyruvate: Sugar Phosphotransferase System Proteins*"

Attorney Docket No.: BGI-122CP

Group Art Unit: Not Assigned

Examiner: Not Assigned

Assistant Commissioner for Patents


BOX SEQUENCE LISTING

Washington, DC 20231

TRANSMITTAL LETTER FOR DISKETTE OF SEQUENCE LISTING

Dear Sir:


Enclosed is a diskette which contains a computer readable form of the Sequence Listing for the patent application filed herewith. The Sequence Listing complies with the requirements of 37 C.F.R. §1.821. The material on this diskette is identical in substance to the paper copy of the Sequence Listing appearing on pages 1-47 which is submitted herewith, as required by 37 CFR §1.821(f). The computer readable form of the Sequence Listing contained on the enclosed diskette is understood to comply with the requirements of §1.824(d).

"Express Mail" mailing label number	EL 373 207 242 US
Date of Deposit	June 27, 2000
I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to: Box Sequence Listing, Assistant Commissioner For Patents, Washington, D.C. 20231	
Signature	
Please Print Name of Person Signing	
Larry Taylor	

LAHIVE & COCKFIELD, LLP

Attorneys at Law

By:


Elizabeth A. Hanley
Registration No. 33,505
28 State Street
Boston, MA 02109

Dated: June 27, 2000

SEQUENCE LISTING

<110> Pompejus, Markus
 Kroger, Burkhard
 Schroder, Hartwig
 Zelder, Oskar
 Haberhauer, Gregor
 <120> CORYNEBACTERIUM GLUTAMICUM GENES ENCODING
 PHOSPHOENOLPYRUVATE:SUGAR PHOSPHOTRANSFERASE
 SYSTEM PROTEINS
 <130> BGI-122CP
 <140>
 <141>
 <160> 34

 <210> 1
 <211> 1527
 <212> DNA
 <213> Corynebacterium glutamicum

 <220>
 <221> CDS
 <222> (101)..(1504)
 <223> RXS00315

 <400> 1
 ctcatggcat ctgcgccgtt cgcgttcttg ccagtggttg ttggtttcac cgcaaccaag 60
 cgtttcggcg gcaatgagtt cctgggcgcc gcgtattggt atg gcg atg gtg ttc 115
 Met Ala Met Val Phe
 1 5

 cag agc ttg gtg aac gcc tac gac gtg gcc gcc acc atg gct gcg gcc 163
 Pro Ser Leu Val Asn Gly Tyr Asp Val Ala Ala Thr Met Ala Ala Gly
 10 15 20

 gaa atg cca atg tgg tcc ctg ttt ggt tta gat gtt gcc caa gcc ggt 211
 Glu Met Pro Met Trp Ser Leu Phe Gly Leu Asp Val Ala Gln Ala Gly
 25 30 35

 tac cag gcc acc gtg ctt cct gtg ctg gtg gtt tct tgg att ctg gca 259
 Tyr Gln Gly Thr Val Leu Pro Val Leu Val Val Ser Trp Ile Leu Ala
 40 45 50

 acg atc gag aag ttc ctg cac aag cga ctc aag gcc act gca gac ttc 307
 Thr Ile Glu Lys Phe Leu His Lys Arg Leu Lys Gly Thr Ala Asp Phe
 55 60 65

 ctg atc act cca gtg ctg acg ttg ctg ctc acc gga ttc ctt aca ttc 355
 Leu Ile Thr Pro Val Leu Thr Leu Leu Leu Thr Gly Phe Leu Thr Phe
 70 75 80 85

 atc gcc att gcc cca gca atg cgc tgg gtg gcc gat gtg ctg gca cac 403
 Ile Ala Ile Gly Pro Ala Met Arg Trp Val Gly Asp Val Leu Ala His
 90 95 100

 ggt cta cag gga ctt tat gat ttc ggt ggt cca gtc gcc ggt ctg ctc 451
 Gly Leu Gln Gly Leu Tyr Asp Phe Gly Gly Pro Val Gly Gly Leu Leu
 105 110 115

 ttc ggt ctg gtc tac tca cca atc gtc atc act ggt ctg cac cag tcc 499

03604221-152700

Phe Gly Leu Val Tyr Ser Pro Ile Val Ile Thr Gly Leu His Gln Ser
 120 125 130
 ttc ccg cca att gag ctg gag ctg ttt aac cag ggt gga tcc ttc atc 547
 Phe Pro Pro Ile Glu Leu Glu Leu Phe Asn Gln Gly Gly Ser Phe Ile
 135 140 145
 ttc gca acg gca tct atg gct aat atc gcc cag ggt gcg gca tgt ttg 595
 Phe Ala Thr Ala Ser Met Ala Asn Ile Ala Gln Gly Ala Ala Cys Leu
 150 155 160 165
 gca gtg ttc ttc ctg gcg aag agt gaa aag ctc aag ggc ctt gca ggt 643
 Ala Val Phe Phe Leu Ala Lys Ser Glu Lys Leu Lys Gly Leu Ala Gly
 170 175 180
 gct tca ggt gtc tcc gct gtt ctt ggt att acg gag cct gcg atc ttc 691
 Ala Ser Gly Val Ser Ala Val Leu Gly Ile Thr Glu Pro Ala Ile Phe
 185 190 195
 ggt gtg aac ctt cgc ctg cgc tgg ccg ttc ttc atc ggt atc ggt acc 739
 Gly Val Asn Leu Arg Leu Arg Trp Pro Phe Phe Ile Gly Ile Gly Thr
 200 205 210
 gca gct atc ggt ggc gct ttg att gca ctc ttt aat atc aag gca gtt 787
 Ala Ala Ile Gly Gly Ala Leu Ile Ala Leu Phe Asn Ile Lys Ala Val
 215 220 225
 gcg ttg ggc gct gca ggt ttc ttg ggt gtt gtt tct att gat gct cca 835
 Ala Leu Gly Ala Ala Gly Phe Leu Gly Val Val Ser Ile Asp Ala Pro
 230 235 240 245
 gat atg gtc atg ttc ttg gtg tgt gca gtt gtt acc ttc ttc atc gca 883
 Asp Met Val Met Phe Leu Val Cys Ala Val Thr Phe Phe Ile Ala
 250 255 260
 ttc ggc gca gcg att gct tat ggc ctt tac ttg gtt cgc cgc aac ggc 931
 Phe Gly Ala Ala Ile Ala Tyr Gly Leu Tyr Leu Val Arg Arg Asn Gly
 265 270 275
 agc att gat cca gat gca acc gct gct cca gtg cct gca gga acg acc 979
 Ser Ile Asp Pro Asp Ala Thr Ala Ala Pro Val Pro Ala Gly Thr Thr
 280 285 290
 aaa gcc gaa gca gaa gca ccc gca gaa ttt tca aac gat tcc acc atc
 1027
 Lys Ala Glu Ala Glu Ala Pro Ala Glu Phe Ser Asn Asp Ser Thr Ile
 295 300 305
 atc cag gca cct ttg acc ggt gaa gct att gca ctg agc agc gtc agc
 1075
 Ile Gln Ala Pro Leu Thr Gly Glu Ala Ile Ala Leu Ser Ser Val Ser
 310 315 320 325
 gat gcc atg ttt gcc agc gga aag ctt ggc tgc ggc gtt gcc atc gtc
 1123
 Asp Ala Met Phe Ala Ser Gly Lys Leu Gly Ser Gly Val Ala Ile Val
 330 335 340
 cca acc aag ggg cag tta gtt tct ccg gtg agt gga aag att gtg gtg
 1171
 Pro Thr Lys Gly Gln Leu Val Ser Pro Val Ser Gly Lys Ile Val Val

00004231.002700

```

345              350              355
gca ttc cca tct ggc cat gct ttc gca gtt cgc acc aag gct gag gat
1219
Ala Phe Pro Ser Gly His Ala Phe Ala Val Arg Thr Lys Ala Glu Asp
360              365              370

ggt tcc aat gtg gat atc ttg atg cac att ggt ttc gac aca gta aac
1267
Gly Ser Asn Val Asp Ile Leu Met His Ile Gly Phe Asp Thr Val Asn
375              380              385

ctc aac ggc acg cac ttt aac ccg ctg aag aag cag ggc gat gaa gtc
1315
Leu Asn Gly Thr His Phe Asn Pro Leu Lys Lys Gln Gly Asp Glu Val
390              395              400              405

aaa gca ggg gag ctg ctg tgt gaa ttc gat att gat gcc att aag gct
1363
Lys Ala Gly Glu Leu Leu Cys Glu Phe Asp Ile Asp Ala Ile Lys Ala
410              415              420

gca ggt tat gag gta acc acg ccg att gtt gtt tog aat tac aag aaa
1411
Ala Gly Tyr Glu Val Thr Thr Pro Ile Val Val Ser Asn Tyr Lys Lys
425              430              435

acc gga cct gta aac act tac ggt ttg ggc gaa att gaa gcg gga gcc
1459
Thr Gly Pro Val Asn Thr Tyr Gly Leu Gly Glu Ile Glu Ala Gly Ala
440              445              450

aac ctg ctc aac gtc gca aag aaa gaa gcg gtg cca gca aca cca
1504
Asn Leu Leu Asn Val Ala Lys Lys Glu Ala Val Pro Ala Thr Pro
455              460              465

taagttgaaa ccttgagtgt tcg
1527

<210> 2
<211> 468
<212> PRT
<213> Corynebacterium glutamicum

<400> 2
Met Ala Met Val Phe Pro Ser Leu Val Asn Gly Tyr Asp Val Ala Ala
1              5              10              15

Thr Met Ala Ala Gly Glu Met Pro Met Trp Ser Leu Phe Gly Leu Asp
20              25              30

Val Ala Gln Ala Gly Tyr Gln Gly Thr Val Leu Pro Val Leu Val Val
35              40              45

Ser Trp Ile Leu Ala Thr Ile Glu Lys Phe Leu His Lys Arg Leu Lys
50              55              60

Gly Thr Ala Asp Phe Leu Ile Thr Pro Val Leu Thr Leu Leu Thr
65              70              75              80

```

Gly	Phe	Leu	Thr	Phe 85	Ile	Ala	Ile	Gly	Pro 90	Ala	Met	Arg	Trp	Val	Gly
Asp	Val	Leu	Ala 100	His	Gly	Leu	Gln	Gly 105	Leu	Tyr	Asp	Phe	Gly 110	Gly	Pro
Val	Gly	Gly 115	Leu	Leu	Phe	Gly	Leu 120	Val	Tyr	Ser	Pro	Ile 125	Val	Ile	Thr
Gly	Leu	His	Gln	Ser	Phe 135	Pro	Pro	Ile	Glu	Leu	Glu 140	Leu	Phe	Asn	Gln
Gly	Gly	Ser	Phe	Ile	Phe 150	Ala	Thr	Ala	Ser	Met 155	Ala	Asn	Ile	Ala	Gln 160
Gly	Ala	Ala	Cys	Leu	Ala 165	Val	Phe	Phe	Leu 170	Ala	Lys	Ser	Glu	Lys	Leu 175
Lys	Gly	Leu	Ala 180	Gly	Ala	Ser	Gly	Val 185	Ser	Ala	Val	Leu	Gly 190	Ile	Thr
Glu	Pro	Ala 195	Ile	Phe	Gly	Val	Asn 200	Leu	Arg	Leu	Arg	Trp 205	Pro	Phe	Phe
Ile	Gly 210	Ile	Gly	Thr	Ala 215	Ala	Ile	Gly	Gly	Ala	Leu 220	Ile	Ala	Leu	Phe
Asn 225	Ile	Lys	Ala	Val	Ala 230	Leu	Gly	Ala	Ala	Gly 235	Phe	Leu	Gly	Val	Val 240
Ser	Ile	Asp	Ala 245	Pro	Asp	Met	Val	Met	Phe 250	Leu	Val	Cys	Ala	Val 255	Val
Thr	Phe	Phe 260	Ile	Ala	Phe	Gly	Ala	Ala 265	Ile	Ala	Tyr	Gly	Leu 270	Tyr	Leu
Val	Arg 275	Arg	Asn	Gly	Ser	Ile	Asp 280	Pro	Asp	Ala	Thr	Ala 285	Ala	Pro	Val
Pro	Ala 290	Gly	Thr	Thr	Lys	Ala 295	Glu	Ala	Glu	Ala	Pro 300	Ala	Glu	Phe	Ser
Asn 305	Asp	Ser	Thr	Ile	Ile 310	Gln	Ala	Pro	Leu	Thr 315	Gly	Glu	Ala	Ile	Ala 320
Leu	Ser	Ser	Val	Ser 325	Asp	Ala	Met	Phe	Ala 330	Ser	Gly	Lys	Leu	Gly 335	Ser
Gly	Val	Ala 340	Ile	Val	Pro	Thr	Lys	Gly 345	Gln	Leu	Val	Ser	Pro 350	Val	Ser
Gly	Lys 355	Ile	Val	Val	Ala	Phe	Pro 360	Ser	Gly	His	Ala 365	Phe	Ala	Val	Arg
Thr	Lys 370	Ala	Glu	Asp	Gly 375	Ser	Asn	Val	Asp	Ile	Leu 380	Met	His	Ile	Gly
Phe 385	Asp	Thr	Val	Asn	Leu 390	Asn	Gly	Thr	His	Phe 395	Asn	Pro	Leu	Lys	Lys 400

QUESTIONS

QUESTIONS

QUESTIONS

QUESTIONS

130	135	140	
ttg gtg tgt gca gtt gtt acc ttc ttc atc gca ttc ggc gca gcg att Leu Val Cys Ala Val Val Thr Phe Phe Ile Ala Phe Gly Ala Ala Ile 145 150 155 160			480
gct tat ggc ctt tac ttg gtt cgc cgc aac ggc agc att gat cca gat Ala Tyr Gly Leu Tyr Leu Val Arg Arg Asn Gly Ser Ile Asp Pro Asp 165 170 175			528
gca acc gct gct cca gtg cct gca gga acg acc aaa gcc gaa gca gaa Ala Thr Ala Ala Pro Val Pro Ala Gly Thr Thr Lys Ala Glu Ala Glu 180 185 190			576
gca ccc gca gaa ttt tca aac gat tcc acc atc atc cag gca cct ttg Ala Pro Ala Glu Phe Ser Asn Asp Ser Thr Ile Ile Gln Ala Pro Leu 195 200 205			624
acc ggt gaa gct att gca ctg agc agc gtc agc gat gcc atg ttt gcc Thr Gly Glu Ala Ile Ala Leu Ser Ser Val Ser Asp Ala Met Phe Ala 210 215 220			672
agc gga aag ctt ggc tcg ggc gtt gcc atc gtc cca acc aag ggg cag Ser Gly Lys Leu Gly Ser Gly Val Ala Ile Val Pro Thr Lys Gly Gln 225 230 235 240			720
tta gtt tct ccg gtg agt gga aag att gtg gtg gca ttc cca tct ggc Leu Val Ser Pro Val Ser Gly Lys Ile Val Val Ala Phe Pro Ser Gly 245 250 255			768
cat gct ttc gca gtt cgc acc aag gct gag gat ggt tcc aat gtg gat His Ala Phe Ala Val Arg Thr Lys Ala Glu Asp Gly Ser Asn Val Asp 260 265 270			816
atc ttg atg cac att ggt ttc gac aca gta aac ctc aac ggc acg cac Ile Leu Met His Ile Gly Phe Asp Thr Val Asn Leu Asn Gly Thr His 275 280 285			864
ttt aac ccg ctg aag aag cag ggc gat gaa gtc aaa gca ggg gag ctg Phe Asn Pro Leu Lys Lys Gln Gly Asp Glu Val Lys Ala Gly Glu Leu 290 295 300			912
ctg tgt gaa ttc gat att gat gcc att aag gct gca ggt tat gag gta Leu Cys Glu Phe Asp Ile Asp Ala Ile Lys Ala Ala Gly Tyr Glu Val 305 310 315 320			960
acc acg ccg att gtt gtt tcg aat tac aag aaa acc gga cct gta aac 1008 Thr Thr Pro Ile Val Val Ser Asn Tyr Lys Lys Thr Gly Pro Val Asn 325 330 335			
act tac ggt ttg ggc gaa att gaa gcg gga gcc aac ctg ctc aac gtc 1056 Thr Tyr Gly Leu Gly Glu Ile Glu Ala Gly Ala Asn Leu Leu Asn Val 340 345 350			
gca aag aaa gaa gcg gtg cca gca aca cca taagttagaa ccttgagtgt 1106 Ala Lys Lys Glu Ala Val Pro Ala Thr Pro 355 360			

400> 4	Tyr	Asp	Phe	Gly	Gly	Pro	Val	Gly	Gly	Leu	Leu	Phe	Gly	Leu	Val	Tyr
1				5						10					15	
Ser	Pro	Ile	Val	Ile	Thr	Gly	Leu	His	Gln	Ser	Phe	Pro	Pro	Ile	Glu	
			20					25						30		
Leu	Glu	Leu	Phe	Asn	Gln	Gly	Gly	Ser	Phe	Ile	Phe	Ala	Thr	Ala	Ser	
		35					40					45				
Met	Ala	Asn	Ile	Ala	Gln	Gly	Ala	Ala	Cys	Leu	Ala	Val	Phe	Phe	Leu	
50						55					60					
Ala	Lys	Ser	Glu	Lys	Leu	Lys	Gly	Leu	Ala	Gly	Ala	Ser	Gly	Val	Ser	
65					70					75					80	
Ala	Val	Leu	Gly	Ile	Thr	Glu	Pro	Ala	Ile	Phe	Gly	Val	Asn	Leu	Arg	
			85					90						95		
Leu	Arg	Trp	Pro	Phe	Phe	Ile	Gly	Ile	Gly	Thr	Ala	Ala	Ile	Gly	Gly	
			100					105					110			
Ala	Leu	Ile	Ala	Leu	Phe	Asn	Ile	Lys	Ala	Val	Ala	Leu	Gly	Ala	Ala	
		115					120				125					
Gly	Phe	Leu	Gly	Val	Val	Ser	Ile	Asp	Ala	Pro	Asp	Met	Val	Met	Phe	
130						135					140					
Leu	Val	Cys	Ala	Val	Val	Thr	Phe	Phe	Ile	Ala	Phe	Gly	Ala	Ala	Ile	
145					150					155					160	
Ala	Tyr	Gly	Leu	Tyr	Leu	Val	Arg	Arg	Asn	Gly	Ser	Ile	Asp	Pro	Asp	
				165				170						175		
Ala	Thr	Ala	Ala	Pro	Val	Pro	Ala	Gly	Thr	Thr	Lys	Ala	Glu	Ala	Glu	
			180					185					190			
Ala	Pro	Ala	Glu	Phe	Ser	Asn	Asp	Ser	Thr	Ile	Ile	Gln	Ala	Pro	Leu	
		195					200					205				
Thr	Gly	Glu	Ala	Ile	Ala	Leu	Ser	Ser	Val	Ser	Asp	Ala	Met	Phe	Ala	
	210					215					220					
Ser	Gly	Lys	Leu	Gly	Ser	Gly	Val	Ala	Ile	Val	Pro	Thr	Lys	Gly	Gln	
225					230					235					240	
Leu	Val	Ser	Pro	Val	Ser	Gly	Lys	Ile	Val	Val	Ala	Phe	Pro	Ser	Gly	
				245				250						255		
His	Ala	Phe	Ala	Val	Arg	Thr	Lys	Ala	Glu	Asp	Gly	Ser	Asn	Val	Asp	
		260						265					270			

```

Ile Leu Met His Ile Gly Phe Asp Thr Val Asn Leu Asn Gly Thr His
  275                                285

Phe Asn Pro Leu Lys Lys Gln Gly Asp Glu Val Lys Ala Gly Glu Leu
  290                                300

Leu Cys Glu Phe Asp Ile Asp Ala Ile Lys Ala Ala Gly Tyr Glu Val
  305                                315                                320

Thr Thr Pro Ile Val Val Ser Asn Tyr Lys Lys Thr Gly Pro Val Asn
  325                                330                                335

Thr Tyr Gly Leu Gly Glu Ile Glu Ala Gly Ala Asn Leu Leu Asn Val
  340                                345                                350

Ala Lys Lys Glu Ala Val Pro Ala Thr Pro
  355                                360

```

```

<210> 5
<211> 372
<212> DNA
<213> Corynebacterium glutamicum

```

```

<220>
<221> CDS
<222> (101)..(349)
<223> RXA01503

<400> 5
gtatcctctcaa aggcctttcta gctgttgacg ctcgacagcga ctcgggtggat acgacatcca 60

cgacctatca aattcttttat gctgcaggcg atgccttttc atg ttc ttg gca gtc 115
Met Phe Leu Ala Val
1 5

att ttg gcg att act gcg gct cgt aaa ttc ggt gcc aat gtc ttt aca 163
Ile Leu Ala Ile Thr Ala Ala Arg Lys Phe Gly Ala Asn Val Phe Thr
10 15 20

tca gtc gca ctc gct ggt gca ttg ctg cag aca cag ctt cag gca gta 211
Ser Val Ala Leu Ala Gly Ala Leu Leu His Thr Gln Leu Gln Ala Val
25 30 35

acc gtg ttg gtt gac ggt gaa ctc cag tcg atg act ctg gtg gct ttc 259
Thr Val Leu Val Asp Gly Glu Leu Gln Ser Met Thr Leu Val Ala Phe
40 45 50

caa aag gct ggt aat gac gtc acc ttc ctg ggc att cca gtg gtg ctg 307
Gln Lys Ala Gly Asn Asp Val Thr Phe Leu Gly Ile Pro Val Val Leu
55 60 65

cag ttg gcg ttg cat gta gcg agt ttg atg aag ttg tcg cga 349
Gln Leu Ala Leu His Val Ala Ser Leu Met Lys Leu Ser Arg
70 75 80

taagaggagg ggcgtgtcgg tct 372

```

```

<210> 6
<211> 83

```

<212> PRT

<213> *Corynebacterium glutamicum*

<400> 6

Met Phe Leu Ala Val Ile Leu Ala Ile Thr Ala Ala Arg Lys Phe Gly
 1 5 10 15

Ala Asn Val Phe Thr Ser Val Ala Leu Ala Gly Ala Leu Leu His Thr
 20 25 30

Gln Leu Gln Ala Val Thr Val Leu Val Asp Gly Glu Leu Gln Ser Met
 35 40 45

Thr Leu Val Ala Phe Gln Lys Ala Gly Asn Asp Val Thr Phe Leu Gly
 50 55 60

Ile Pro Val Val Leu Gln Leu Ala Leu His Val Ala Ser Leu Met Lys
 65 70 75 80

Leu Ser Arg

<210> 7

<211> 2187

<212> DNA

<213> *Corynebacterium glutamicum*

<220>

<221> CDS

<222> (101)..(2164)

<223> RXN01299

<400> 7

cgactcgggc gtctcttccct ggcactacca ttccctcgtec tgaccaactc gccacagctg 60

gtgcaacggt cacccaagtc aaaggattga aagaatcagc atg aat agc gta aat 115
 Met Asn Ser Val Asn
 1 5

aat tcc tcg ctt gtc cgg ctg gat gtc gat ttc ggc gac tcc acc acg 163
 Asn Ser Ser Leu Val Arg Leu Asp Val Asp Phe Gly Asp Ser Thr Thr
 10 15 20

gat gtc atc aac aac ctt gcc act gtt att ttc gac gct ggc cga gct 211
 Asp Val Ile Asn Asn Leu Ala Thr Val Ile Phe Asp Ala Gly Arg Ala
 25 30 35

tcc tcc gcc gac gcc ctt gcc aaa gac cgg ctg gat cgt gaa gca aag 259
 Ser Ser Ala Asp Ala Leu Ala Lys Asp Ala Leu Asp Arg Glu Ala Lys
 40 45 50

tcc ggc acc ggc gtt cct ggt caa gtt gct atc ccc cac tgc cgt tcc 307
 Ser Gly Thr Gly Val Pro Gly Gln Val Ala Ile Pro His Cys Arg Ser
 55 60 65

gaa gcc gta tct gtc cct acc ttg ggc ttt gct cgc ctg agc aag ggt 355
 Glu Ala Val Ser Val Pro Thr Leu Gly Phe Ala Arg Leu Ser Lys Gly
 70 75 80 85

gtg gac ttc agc gga cct gat ggc gat gcc aac ttg gtg ttc ctc att 403

Val	Asp	Phe	Ser	Gly	Pro	Asp	Gly	Asp	Ala	Asn	Leu	Val	Phe	Leu	Ile	
				90					95					100		
gca	gca	cct	gct	ggc	ggc	ggc	aaa	gag	cac	ctg	aag	atc	ctg	tcc	aag	451
Ala	Ala	Pro	Ala	Gly	Gly	Gly	Lys	Glu	His	Leu	Lys	Ile	Leu	Ser	Lys	
			105					110					115			
ctt	gct	cgc	tcc	ttg	gtg	aag	aag	gat	ttc	atc	aag	gct	ctg	cag	gaa	499
Leu	Ala	Arg	Ser	Leu	Val	Lys	Lys	Asp	Phe	Ile	Lys	Ala	Leu	Gln	Glu	
			120					125					130			
gcc	acc	acc	gag	cag	gaa	atc	gtc	gac	gtt	gtc	gat	gcc	gtg	ctc	aac	547
Ala	Thr	Thr	Glu	Gln	Glu	Ile	Val	Asp	Val	Val	Asp	Ala	Val	Leu	Asn	
			135				140					145				
cca	gca	cca	aaa	acc	acc	gag	cca	gct	gca	gct	ccg	gct	gcg	gcg	gcg	595
Pro	Ala	Pro	Lys	Thr	Thr	Glu	Pro	Ala	Ala	Ala	Pro	Ala	Ala	Ala	Ala	
						155					160				165	
gtt	gct	gag	agt	ggg	gcg	gcg	tcg	aca	agc	gtt	act	cgt	atc	gtg	gca	643
Val	Ala	Glu	Ser	Gly	Ala	Ala	Ser	Thr	Ser	Val	Thr	Arg	Ile	Val	Ala	
				170						175				180		
atc	acc	gca	tgc	cca	acc	ggt	atc	gca	cac	acc	tac	atg	gct	gcg	gat	691
Ile	Thr	Ala	Cys	Pro	Thr	Gly	Ile	Ala	His	Thr	Tyr	Met	Ala	Ala	Asp	
				185				190					195			
tcc	ctg	acg	caa	aac	gcg	gaa	ggc	cgc	gat	gat	gtg	gaa	ctc	gtt	gtg	739
Ser	Leu	Thr	Gln	Asn	Ala	Glu	Gly	Arg	Asp	Asp	Val	Glu	Leu	Val	Val	
			200				205					210				
gag	act	cag	ggc	tct	tcc	gct	gtc	acc	cca	gtc	gat	ccg	aag	atc	atc	787
Glu	Thr	Gln	Gly	Ser	Ser	Ala	Val	Thr	Pro	Val	Asp	Pro	Lys	Ile	Ile	
			215				220					225				
gaa	gct	gcc	gac	gcc	gtc	atc	ttc	gcc	acc	gac	gtg	gga	gtt	aaa	gac	835
Glu	Ala	Ala	Asp	Ala	Val	Ile	Phe	Ala	Thr	Asp	Val	Gly	Val	Lys	Asp	
			230			235				240				245		
cgc	gag	cgt	ttc	gct	ggc	aag	cca	gtc	att	gaa	tcc	ggc	gtc	aag	cgc	883
Arg	Glu	Arg	Phe	Ala	Gly	Lys	Pro	Val	Ile	Glu	Ser	Gly	Val	Lys	Arg	
				250					255				260			
gcg	atc	aat	gag	cca	gcc	aag	atg	atc	gac	gag	gcc	atc	gca	gcc	tcc	931
Ala	Ile	Asn	Glu	Pro	Ala	Lys	Met	Ile	Asp	Glu	Ala	Ile	Ala	Ala	Ser	
				265				270					275			
aag	aac	cca	aac	gcc	cgc	aag	gtt	tcc	ggt	tcc	ggt	gtc	gcg	gca	tct	979
Lys	Asn	Pro	Asn	Ala	Arg	Lys	Val	Ser	Gly	Ser	Gly	Val	Ala	Ala	Ser	
			280				285					290				
gct	gaa	acc	acc	ggc	gag	aag	ctc	ggc	tgg	ggc	aag	cgc	atc	cag	cag	
			1027													
Ala	Glu	Thr	Thr	Gly	Glu	Lys	Leu	Gly	Trp	Gly	Lys	Arg	Ile	Gln	Gln	
			295				300					305				
gca	gtc	atg	acc	ggc	gtg	tcc	tac	atg	gtt	cca	ttc	gta	gct	gcc	ggc	
			1075													
Ala	Val	Met	Thr	Gly	Val	Ser	Tyr	Met	Val	Pro	Phe	Val	Ala	Ala	Gly	
			310			315				320					325	

ggc ctc ctg ttg gct ctc ggc ttc gca ttc ggt gga tac gac atg gcg
 1123
 Gly Leu Leu Leu Ala Leu Gly Phe Ala Phe Gly Gly Tyr Asp Met Ala
 330 335 340

aac ggc tgg gca gca atc gcc acc cag ttc tct ctg acc aac ctg cca
 1171
 Asn Gly Trp Gln Ala Ile Ala Thr Gln Phe Ser Leu Thr Asn Leu Pro
 345 350 355

ggc aac acc gtc gat gtt gac ggc gtg gcc atg acc ttc gag cgt tca
 1219
 Gly Asn Thr Val Asp Val Asp Gly Val Ala Met Thr Phe Glu Arg Ser
 360 365 370

ggc ttc ctg ttg tac ttc ggc gca gtc ctg ttc gcc acc ggc caa gca
 1267
 Gly Phe Leu Leu Tyr Phe Gly Ala Val Leu Phe Ala Thr Gly Gln Ala
 375 380 385

gcc atg ggc ttc atc gtg gca gcc ctg tct ggc tac acc gca tac gca
 1315
 Ala Met Gly Phe Ile Val Ala Ala Leu Ser Gly Tyr Thr Ala Tyr Ala
 390 395 400 405

ctt gct gga cgc cca ggc atc gcg ccg ggc ttc gtc ggt ggc gcc atc
 1363
 Leu Ala Gly Arg Pro Gly Ile Ala Pro Gly Phe Val Gly Gly Ala Ile
 410 415 420

tcc gtc acc atc ggc gct ggc ttc att ggt ggt ctg gtt acc ggt atc
 1411
 Ser Val Thr Ile Gly Ala Gly Phe Ile Gly Gly Leu Val Thr Gly Ile
 425 430 435

ttg gct ggt ctc att gcc ctg tgg att ggc tcc tgg aag gtg cca cgc
 1459
 Leu Ala Gly Leu Ile Ala Leu Trp Ile Gly Ser Trp Lys Val Pro Arg
 440 445 450

gtg gtg cag tca ctg atg cct gtg gtc atc atc ccg cta ctt acc tca
 1507
 Val Val Gln Ser Leu Met Pro Val Val Ile Ile Pro Leu Leu Thr Ser
 455 460 465

gtg gtt gtt ggt ctc gtc atg tac ctc ctg ctg ggt cgc cca ctc gca
 1555
 Val Val Val Gly Leu Val Met Tyr Leu Leu Leu Gly Arg Pro Leu Ala
 470 475 480 485

tcc atc atg act ggt ttg cag gac tgg cta tcg tca atg tcc gga agc
 1603
 Ser Ile Met Thr Gly Leu Gln Asp Trp Leu Ser Ser Met Ser Gly Ser
 490 495 500

tcc gcc atc ttg ctg ggt atc atc ttg ggc ctc atg atg tgt ttc gac
 1651
 Ser Ala Ile Leu Leu Gly Ile Ile Leu Gly Leu Met Met Cys Phe Asp
 505 510 515

ctc ggc gga cca gta aac aag gca gcc tac ctc ttt ggt acc gca ggc
 1699
 Leu Gly Gly Pro Val Asn Lys Ala Ala Tyr Leu Phe Gly Thr Ala Gly
 520 525 530

ctg tct acc ggc gac caa gct tcc atg gaa atc atg gcc gcg atc atg
 1747
 Leu Ser Thr Gly Asp Gln Ala Ser Met Glu Ile Met Ala Ala Ile Met
 535 540 545

gca gct ggc atg gtc cca cca atc gcg ttg tcc att gct acc ctg ctg
 1795
 Ala Ala Gly Met Val Pro Pro Ile Ala Leu Ser Ile Ala Thr Leu Leu
 550 555 560 565

gcg aag aag ctg ttc acc cca gca gag caa gaa aac ggc aag tct tcc
 1843
 Arg Lys Lys Leu Phe Thr Pro Ala Glu Gln Glu Asn Gly Lys Ser Ser
 570 575 580

tgg ctg ctt ggc ctg gca ttc gtc tcc gaa ggt gcc atc cca ttc gcc
 1891
 Trp Leu Leu Gly Leu Ala Phe Val Ser Glu Gly Ala Ile Pro Phe Ala
 585 590 595

gca gct gac cca ttc cgt gtg atc cca gca atg atg gct ggc ggt gca
 1939
 Ala Ala Asp Pro Phe Arg Val Ile Pro Ala Met Met Ala Gly Gly Ala
 600 605 610

acc act ggt gca atc tcc atg gca ctg ggc gtc ggc tct cgg gct cca
 1987
 Thr Thr Gly Ala Ile Ser Met Ala Leu Gly Val Gly Ser Arg Ala Pro
 615 620 625

cac ggc ggt atc ttc gtg gtc tgg gca atc gaa cca tgg tgg ggc tgg
 2035
 His Gly Gly Ile Phe Val Val Trp Ala Ile Glu Pro Trp Trp Gly Trp
 630 635 640 645

ctc atc gca ctt gca gca ggc acc atc gtg tcc acc atc gtt gtc atc
 2083
 Leu Ile Ala Leu Ala Ala Gly Thr Ile Val Ser Thr Ile Val Val Ile
 650 655 660

gca ctg aag cag ttc tgg cca aac aag gcc gtc gct gca gaa gtc gcg
 2131
 Ala Leu Lys Gln Phe Trp Pro Asn Lys Ala Val Ala Ala Glu Val Ala
 665 670 675

aag caa gaa gca caa caa gca gct gta aac gca taatcggacc ttgaccgat
 2184
 Lys Gln Glu Ala Gln Gln Ala Ala Val Asn Ala
 680 685

gtc
 2187

<210> 8

<211> 688

<212> PRT

<213> *Corynebacterium glutamicum*

<400> 8

```

Met Asn Ser Val Asn Asn Ser Ser Leu Val Arg Leu Asp Val Asp Phe
  1             5             10             15

Gly Asp Ser Thr Thr Asp Val Ile Asn Asn Leu Ala Thr Val Ile Phe
          20             25             30

Asp Ala Gly Arg Ala Ser Ser Ala Asp Ala Leu Ala Lys Asp Ala Leu
          35             40             45

Asp Arg Glu Ala Lys Ser Gly Thr Gly Val Pro Gly Gln Val Ala Ile
          50             55             60

Pro His Cys Arg Ser Glu Ala Val Ser Val Pro Thr Leu Gly Phe Ala
          65             70             75             80

Arg Leu Ser Lys Gly Val Asp Phe Ser Gly Pro Asp Gly Asp Ala Asn
          85             90             95

Leu Val Phe Leu Ile Ala Ala Pro Ala Gly Gly Gly Lys Glu His Leu
          100            105            110

Lys Ile Leu Ser Lys Leu Ala Arg Ser Leu Val Lys Lys Asp Phe Ile
          115            120            125

Lys Ala Leu Gln Glu Ala Thr Thr Glu Gln Glu Ile Val Asp Val Val
          130            135            140

Asp Ala Val Leu Asn Pro Ala Pro Lys Thr Thr Glu Pro Ala Ala Ala
          145            150            155            160

Pro Ala Ala Ala Ala Val Ala Glu Ser Gly Ala Ala Ser Thr Ser Val
          165            170            175

Thr Arg Ile Val Ala Ile Thr Ala Cys Pro Thr Gly Ile Ala His Thr
          180            185            190

Tyr Met Ala Ala Asp Ser Leu Thr Gln Asn Ala Glu Gly Arg Asp Asp
          195            200            205

Val Glu Leu Val Val Glu Thr Gln Gly Ser Ser Ala Val Thr Pro Val
          210            215            220

Asp Pro Lys Ile Ile Glu Ala Ala Asp Ala Val Ile Phe Ala Thr Asp
          225            230            235            240

Val Gly Val Lys Asp Arg Glu Arg Phe Ala Gly Lys Pro Val Ile Glu
          245            250            255

Ser Gly Val Lys Arg Ala Ile Asn Glu Pro Ala Lys Met Ile Asp Glu
          260            265            270

Ala Ile Ala Ala Ser Lys Asn Pro Asn Ala Arg Lys Val Ser Gly Ser
          275            280            285

Gly Val Ala Ala Ser Ala Glu Thr Thr Gly Glu Lys Leu Gly Trp Gly
          290            295            300

```

Year	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099
1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	

625					630					635				640
Pro	Trp	Trp	Gly	Trp	Leu	Ile	Ala	Leu	Ala	Ala	Gly	Thr	Ile	Val
			645						650					Ser
Thr	Ile	Val	Val	Ile	Ala	Leu	Lys	Gln	Phe	Trp	Pro	Asn	Lys	Ala
			660					665					670	Val
Ala	Ala	Glu	Val	Ala	Lys	Gln	Glu	Ala	Gln	Gln	Ala	Ala	Val	Asn
		675					680				685			Ala

```
<210> 9
<211> 464
<212> DNA
<213> Corynebacterium glutamicum
```

```
<220>  
<221> CDS  
<222> (1)..(441)  
<223> FRXA01299
```

400> 9																
atg	gaa	atc	atg	gcc	gcg	atc	atg	gca	gct	ggc	atg	gtc	cca	cca	atc	48
Met	Glu	Ile	Met	Ala	Ala	Ile	Met	Ala	Ala	Gly	Met	Val	Pro	Pro	Ile	
1				5					10				15			
gcg ttg tcc att gct acc ctg ctg cgc aag aag ctg ttc acc cca gca															96	
Ala	Leu	Ser	Ile	Ala	Thr	Leu	Leu	Arg	Lys	Lys	Leu	Phe	Thr	Pro	Ala	
			20					25					30			
gag caa gaa aac ggc aag tct tcc tgg ctg ctt ggc ctg gca ttc gtc															144	
Glu	Gln	Glu	Asn	Gly	Lys	Ser	Ser	Trp	Leu	Leu	Gly	Leu	Ala	Phe	Val	
			35				40					45				
tcc gaa ggt gcc atc cca ttc gcc gca gct gac cca ttc cgt gtg atc															192	
Ser	Glu	Gly	Ala	Ile	Pro	Phe	Ala	Ala	Ala	Asp	Pro	Phe	Arg	Val	Ile	
	50					55					60					
cca gca atg atg gct gcc ggt gca acc act ggt gca atc tcc atg gca															240	
Pro	Ala	Met	Met	Ala	Gly	Gly	Ala	Thr	Thr	Gly	Ala	Ile	Ser	Met	Ala	
	65				70					75				80		
ctg gcc gtc gcc tct cgg gct cca cac gcc ggt atc ttc gtg gtc tgg															288	
Leu	Gly	Val	Gly	Ser	Arg	Ala	Pro	His	Gly	Gly	Ile	Phe	Val	Val	Trp	
				85					90					95		
gca atc gaa cca tgg tgg gcc tgg ctc atc gca ctt gca gca gcc acc															336	
Ala	Ile	Glu	Pro	Trp	Trp	Gly	Trp	Leu	Ile	Ala	Leu	Ala	Ala	Gly	Thr	
			100					105					110			
atc gtg tcc acc atc gtt gtc atc gca ctg aag cag ttc tgg cca aac															384	
Ile	Val	Ser	Thr	Ile	Val	Val	Ile	Ala	Leu	Lys	Gln	Phe	Trp	Pro	Asn	
		115					120					125				
aag gcc gtc gct gca gaa gtc gcg aag caa gaa gca caa caa gca gct															432	
Lys	Ala	Val	Ala	Ala	Glu	Val	Ala	Lys	Gln	Glu	Ala	Gln	Gln	Ala	Ala	
	130					135				140						

gta aac gca taatcggacc ttgaccgat gtc
Val Asn Ala
145

464

<210> 10
<211> 147
<212> PRT
<213> Corynebacterium glutamicum

<400> 10
Met Glu Ile Met Ala Ala Ile Met Ala Ala Gly Met Val Pro Pro Ile
1 5 10 15

Ala Leu Ser Ile Ala Thr Leu Leu Arg Lys Lys Leu Phe Thr Pro Ala
20 25 30

Glu Gln Glu Asn Gly Lys Ser Ser Trp Leu Leu Gly Leu Ala Phe Val
35 40 45

Ser Glu Gly Ala Ile Pro Phe Ala Ala Ala Asp Pro Phe Arg Val Ile
50 55 60

Pro Ala Met Met Ala Gly Gly Ala Thr Thr Gly Ala Ile Ser Met Ala
65 70 75 80

Leu Gly Val Gly Ser Arg Ala Pro His Gly Gly Ile Phe Val Val Trp
85 90 95

Ala Ile Glu Pro Trp Trp Gly Trp Leu Ile Ala Leu Ala Ala Gly Thr
100 105 110

Ile Val Ser Thr Ile Val Val Ile Ala Leu Lys Gln Phe Trp Pro Asn
115 120 125

Lys Ala Val Ala Ala Glu Val Ala Lys Gln Glu Ala Gln Gln Ala Ala
130 135 140

Val Asn Ala
145

<210> 11
<211> 580
<212> DNA
<213> Corynebacterium glutamicum

<220>
<221> CDS
<222> (101)..(580)
<223> FRXA01883

<400> 11
cgactgggc gtctcttctt ggcactacca ttectcgcc tgaccaactc gccacagctg 60

gtgcaacggt caccgaagtc aaaggattga aagaatcagc atg aat agc gta aat 115
Met Asn Ser Val Asn
1 5

aat tcc tcg ctt gtc cgg ctg gat gtc gat ttc ggc gac tcc acc acg 163

Asn Ser Ser Leu Val Arg Leu Asp Val Asp Phe Gly Asp Ser Thr Thr
 10 15 20
 gat gtc atc aac aac ctt gcc act gtt att ttc gac gct ggc cga gct 211
 Asp Val Ile Asn Asn Leu Ala Thr Val Ile Phe Asp Ala Gly Arg Ala
 25 30 35
 tcc tcc gcc gac gcc ctt gcc aaa gac gcg ctg gat cgt gaa gca aag 259
 Ser Ser Ala Asp Ala Leu Ala Lys Asp Ala Leu Asp Arg Glu Ala Lys
 40 45 50
 tcc ggc acc ggc gtt cct ggt caa gtt gct atc ccc cac tgc cgt tcc 307
 Ser Gly Thr Gly Val Pro Gly Gln Val Ala Ile Pro His Cys Arg Ser
 55 60 65
 gaa gcc gta tct gtc cct acc ttg ggc ttt gct cgc ctg agc aag ggt 355
 Glu Ala Val Ser Val Pro Thr Leu Gly Phe Ala Arg Leu Ser Lys Gly
 70 75 80 85
 gtg gac ttc agc gga cct gat ggc gat gcc aac ttg gtg ttc ctc att 403
 Val Asp Phe Ser Gly Pro Asp Gly Asp Ala Asn Leu Val Phe Leu Ile
 90 95 100
 gca gca cct gct ggc ggc ggc aaa gag cac ctg aag atc ctg tcc aag 451
 Ala Ala Pro Ala Gly Gly Gly Lys Glu His Leu Lys Ile Leu Ser Lys
 105 110 115
 ctt gct cgc tcc ttg gtg aag aag gat ttc atc aag gct ctg cag gaa 499
 Leu Ala Arg Ser Leu Val Lys Lys Asp Phe Ile Lys Ala Leu Gln Glu
 120 125 130
 gcc acc acc gag cag gaa atc gtc gac gtt gtc gat gcc gtg ctc aac 547
 Ala Thr Thr Glu Gln Glu Ile Val Asp Val Val Asp Ala Val Leu Asn
 135 140 145
 cca gca cca aaa aac cac cga gcc agc tgc agc 580
 Pro Ala Pro Lys Asn His Arg Ala Ser Cys Ser
 150 155 160

<210> 12

<211> 160

<212> PRT

<213> Corynebacterium glutamicum

<400> 12

Met Asn Ser Val Asn Asn Ser Ser Leu Val Arg Leu Asp Val Asp Phe
 1 5 10 15
 Gly Asp Ser Thr Thr Asp Val Ile Asn Asn Leu Ala Thr Val Ile Phe
 20 25 30
 Asp Ala Gly Arg Ala Ser Ser Ala Asp Ala Leu Ala Lys Asp Ala Leu
 35 40 45
 Asp Arg Glu Ala Lys Ser Gly Thr Gly Val Pro Gly Gln Val Ala Ile
 50 55 60
 Pro His Cys Arg Ser Glu Ala Val Ser Val Pro Thr Leu Gly Phe Ala
 65 70 75 80

```

Arg Leu Ser Lys Gly Val Asp Phe Ser Gly Pro Asp Gly Asp Ala Asn
      85                      90                      95
Leu Val Phe Leu Ile Ala Ala Pro Ala Gly Gly Gly Lys Glu His Leu
      100                    105                    110
Lys Ile Leu Ser Lys Leu Ala Arg Ser Leu Val Lys Lys Asp Phe Ile
      115                    120                    125
Lys Ala Leu Gln Glu Ala Thr Thr Glu Gln Glu Ile Val Asp Val Val
      130                    135                    140
Asp Ala Val Leu Asn Pro Ala Pro Lys Asn His Arg Ala Ser Cys Ser
      145                    150                    155                    160

```

```

<210> 13
<211> 631
<212> DNA
<213> Corynebacterium glutamicum

```

```

<220>
<221> CDS
<222> (77)..(631)
<223> FRXA01889

```

```

<400> 13
acgcagccag ctgcagctcc ggctgcggcg gcgcgttggt aagagtgggg cggcgtcgac 60

aagcgttact cgtatcgtg gca atc acc gca tgc cca acc ggt atc gca cac 112
      Val Ala Ile Thr Ala Cys Pro Thr Gly Ile Ala His
      1                      5                      10

acc tac atg gct gcg gat tcc ctg acg caa aac gcg gaa ggc cgc gat 160
Thr Tyr Met Ala Ala Asp Ser Leu Thr Gln Asn Ala Glu Gly Arg Asp
      15                      20                      25

gat gtg gaa ctc gtt gtg gag act cag ggc tct tcc gct gtc acc cca 208
Asp Val Glu Leu Val Val Glu Thr Gln Gly Ser Ser Ala Val Thr Pro
      30                      35                      40

gtc gat ccg aag atc atc gaa gct gcc gac gcc gtc atc ttc gcc acc 256
Val Asp Pro Lys Ile Ile Glu Ala Ala Asp Ala Val Ile Phe Ala Thr
      45                      50                      55                      60

gac gtg gga gtt aaa gac cgc gag cgt ttc gct ggc aag cca gtc att 304
Asp Val Gly Val Lys Asp Arg Glu Arg Phe Ala Gly Lys Pro Val Ile
      65                      70                      75

gaa tcc ggc gtc aag cgc gcg atc aat gag cca gcc aag atg atc gac 352
Glu Ser Gly Val Lys Arg Ala Ile Asn Glu Pro Ala Lys Met Ile Asp
      80                      85                      90

gag gcc atc gca gcc tcc aag aac cca aac gcc cgc aag gtt tcc ggt 400
Glu Ala Ile Ala Ala Ser Lys Asn Pro Asn Ala Arg Lys Val Ser Gly
      95                      100                    105

tcc ggt gtc gcg gca tct gct gaa acc acc ggc gag aag ctc ggc tgg 448

```

Ser	Gly	Val	Ala	Ala	Ser	Ala	Glu	Thr	Thr	Gly	Glu	Lys	Leu	Gly	Trp	
	110					115					120					
ggc	aag	cgc	atc	cag	cag	gca	gtc	atg	acc	ggc	gtg	tcc	tac	atg	gtt	496
Gly	Lys	Arg	Ile	Gln	Gln	Ala	Val	Met	Thr	Gly	Val	Ser	Tyr	Met	Val	
	125				130					135				140		
cca	ttc	gta	gct	gcc	ggc	ggc	ctc	ctg	tgt	gct	ctc	ggc	ttc	gca	ttc	544
Pro	Phe	Val	Ala	Ala	Gly	Gly	Leu	Leu	Leu	Ala	Leu	Gly	Phe	Ala	Phe	
				145				150					155			
ggt	gga	tac	gac	atg	gcg	aac	ggc	tgg	caa	gca	atc	gcc	acc	cag	ttc	592
Gly	Gly	Tyr	Asp	Met	Ala	Asn	Gly	Trp	Gln	Ala	Ile	Ala	Thr	Gln	Phe	
			160					165				170				
tct	ctg	acc	aac	ctg	cca	ggc	aac	acc	gtc	gat	gtt	gac				631
Ser	Leu	Thr	Asn	Leu	Pro	Gly	Asn	Thr	Val	Asp	Val	Asp				
		175					180				185					

<210> 14

<211> 18

<212> PRT

<213> *Corynebacterium glutamicum*

<400> 14

Val Ala Ile Thr Ala Cys Pro Thr Gly Ile Ala His Thr Tyr Met Ala
1 5 10 15

Ala Asp Ser Leu Thr Gln Asn Ala Glu Gly Arg Asp Asp Val Glu Leu
20 25 30

Val Val Glu Thr Gln Gly Ser Ser Ala Val Thr Pro Val Asp Pro Lys
35 40 45

Ile Ile Glu Ala Ala Asp Ala Val Ile Phe Ala Thr Asp Val Gly Val
50 55 60

Lys Asp Arg Glu Arg Phe Ala Gly Lys Pro Val Ile Glu Ser Gly Val
65 70 75 80

Lys Arg Ala Ile Asn Glu Pro Ala Lys Met Ile Asp Glu Ala Ile Ala
85 90 95

Ala Ser Lys Asn Pro Asn Ala Arg Lys Val Ser Gly Ser Gly Val Ala
100 105 110

Ala Ser Ala Glu Thr Thr Gly Glu Lys Leu Gly Trp Gly Lys Arg Ile
115 120 125

Gln Gln Ala Val Met Thr Gly Val Ser Tyr Met Val Pro Phe Val Ala
130 135 140

Ala Gly Gly Leu Leu Leu Ala Leu Gly Phe Ala Phe Gly Gly Tyr Asp
145 150 155 160

Met Ala Asn Gly Trp Gln Ala Ile Ala Thr Gln Phe Ser Leu Thr Asn
165 170 175

Leu Pro Gly Asn Thr Val Asp Val Asp
180 185

```
<210> 15
<211> 416
<212> DNA
<213> Corynebacterium glutamicum
```

```
<220>  
<221> CDS  
<222> (1)..(393)  
<223> RXA00951
```

400> 15																
atc	caa	gca	atc	tta	gag	aag	gca	gca	gcg	ccg	gcg	aag	cag	aag	gct	48
Ile	Gln	Ala	Ile	Leu	Glu	Lys	Ala	Ala	Ala	Pro	Ala	Lys	Gln	Lys	Ala	
1				5					10					15		
cct	gct	gtg	gct	cct	gct	gta	aca	ccc	act	gac	gct	cct	gca	gcc	tca	96
Pro	Ala	Val	Ala	Pro	Ala	Val	Thr	Pro	Thr	Asp	Ala	Pro	Ala	Ala	Ser	
			20					25					30			
gtc	caa	tcc	aaa	acc	cac	gac	aag	atc	ctc	acc	gtc	tgt	ggc	aac	ggc	144
Val	Gln	Ser	Lys	Thr	His	Asp	Lys	Ile	Leu	Thr	Val	Cys	Gly	Asn	Gly	
			35				40					45				
ttg	ggt	acc	tcc	ctc	ttc	ctc	aaa	aac	acc	ctt	gag	caa	gtt	ttc	gac	192
Leu	Gly	Thr	Ser	Leu	Phe	Leu	Lys	Asn	Thr	Leu	Glu	Gln	Val	Phe	Asp	
	50					55				60						
acc	tggt	ggt	tggt	ggt	cca	tac	atg	acg	gtg	gag	gca	acc	gac	act	atc	240
Thr	Trp	Gly	Trp	Gly	Pro	Tyr	Met	Thr	Val	Glu	Ala	Thr	Asp	Thr	Ile	
65					70				75					80		
tcc	gcc	aag	ggc	aaa	gcc	aag	gaa	gct	gat	ctc	atc	glu	acc	tct	ggt	288
Ser	Ala	Lys	Gly	Lys	Ala	Lys	Glu	Ala	Asp	Leu	Ile	Met	Thr	Ser	Gly	
				85					90				95			
gaa	atc	gcc	cgc	acg	ttg	ggt	gat	gtt	gga	atc	ccg	gtt	cac	gtg	atc	336
Glu	Ile	Ala	Arg	Thr	Leu	Gly	Asp	Val	Gly	Ile	Pro	Val	His	Val	Ile	
			100				105						110			
aat	gac	ttc	acg	agc	acc	gat	gaa	atc	gat	gct	gcg	ctt	cgt	gaa	cgc	384
Asn	Asp	Phe	Thr	Ser	Thr	Asp	Glu	Ile	Asp	Ala	Ala	Lys	Arg	Glu	Arg	
		115				120						125				
tac	gac	atc	taactacttt	aaaaggacga	aaa											416
Tyr	Asp	Ile														
130																

```
<210> 16
<211> 131
<212> PRT
<213> Corynebacterium glutamicum
```

<400> 16
 Ile Gln Ala Ile Leu Glu Lys Ala Ala Ala Pro Ala Lys Gln Lys Ala
 1 5 10 15
 Pro Ala Val Ala Pro Ala Val Thr Pro Thr Asp Ala Pro Ala Ala Ser
 20 25 30

Val Gln Ser Lys Thr His Asp Lys Ile Leu Thr Val Cys Gly Asn Gly
 35 40 45

Leu Gly Thr Ser Leu Phe Leu Lys Asn Thr Leu Glu Gln Val Phe Asp
 50 55 60

Thr Trp Gly Trp Gly Pro Tyr Met Thr Val Glu Ala Thr Asp Thr Ile
 65 70 75 80

Ser Ala Lys Gly Lys Ala Lys Glu Ala Asp Leu Ile Met Thr Ser Gly
 85 90 95

Glu Ile Ala Arg Thr Leu Gly Asp Val Gly Ile Pro Val His Val Ile
 100 105 110

Asn Asp Phe Thr Ser Thr Asp Glu Ile Asp Ala Ala Leu Arg Glu Arg
 115 120 125

Tyr Asp Ile
 130

<210> 17
 <211> 1827
 <212> DNA
 <213> Corynebacterium glutamicum

<220>
 <221> CDS
 <222> (101)..(1804)
 <223> RXN01244

<400> 17
 gatattgtgtt tgtttgtcaa tatccaaatg tttgaatagt tgcacaactg ttggttttgt 60

ggtgatcttg aggaaattaa ctcaatgatt gtgaggatgg gtg gct act gtg gct 115
 Val Ala Thr Val Ala
 1 5

gat gtg aat caa gac act gta ctg aag ggc acc ggc gtt gtc ggt gga 163
 Asp Val Asn Gln Asp Thr Val Leu Lys Gly Thr Gly Val Val Gly Gly
 10 15 20

gtc cgt tat gca agc gcg gtg tgg att acc cca cgc ccc gaa cta ccc 211
 Val Arg Tyr Ala Ser Ala Val Trp Ile Thr Pro Arg Pro Glu Leu Pro
 25 30 35

caa gca ggc gaa gtc gtc gcc gaa gaa aac cgt gaa gca gag cag gag 259
 Gln Ala Gly Glu Val Val Ala Glu Glu Asn Arg Glu Ala Glu Gln Glu
 40 45 50

cgt ttc gac gcc gct gca gcc aca gtc tct tct cgt ttg ctt gag cgc 307
 Arg Phe Asp Ala Ala Ala Thr Val Ser Ser Arg Leu Leu Glu Arg
 55 60 65

tcc gaa gct gct gaa gga cca gca gct gag gtg ctt aaa gct act gct 355
 Ser Glu Ala Ala Glu Gly Pro Ala Ala Glu Val Leu Lys Ala Thr Ala
 70 75 80 85

ggc atg gtc aat gac cgt ggc tgg cgt aag gct gtc atc aag ggt gtc 403

C:\BGI\BGI-122CP\BGI-122CP-21-1827.DNA

Gly Met Val Asn Asp Arg Gly Trp Arg Lys Ala Val Ile Lys Gly Val	
90 95 100	
aag ggt ggt cac cct gcg gaa tac gcc gtg gtt gca gca aca acc aag	451
Lys Gly Gly His Pro Ala Glu Tyr Ala Val Val Ala Ala Thr Thr Lys	
105 110 115	
ttc atc tcc atg ttc gaa gcc gca ggc ctg atc gcg gag cgc acc	499
Phe Ile Ser Met Phe Glu Ala Ala Gly Gly Leu Ile Ala Glu Arg Thr	
120 125 130	
aca gac ttg cgc gac atc cgc gac cgc gtc atc gca gaa ctt cgt ggc	547
Thr Asp Leu Arg Asp Ile Arg Asp Arg Val Ile Ala Glu Leu Arg Gly	
135 140 145	
gat gaa gag cca ggt ctg cca gct gtt tcc gga cag gtc att ctc ttt	595
Asp Glu Glu Pro Gly Leu Pro Ala Val Ser Gly Gln Val Ile Leu Phe	
150 155 160 165	
gca gat gac ctc tcc cca gca gac acc gcg gca cta gac aca gat ctc	643
Ala Asp Asp Leu Ser Pro Ala Asp Thr Ala Ala Leu Asp Thr Asp Leu	
170 175 180	
ttt gtg gga ctt gtc act gag ctg ggt ggc cca acg agc cac acc gcg	691
Phe Val Gly Leu Val Thr Glu Leu Gly Gly Pro Thr Ser His Thr Ala	
185 190 195	
atc atc gca cgc cag ctc aac gtg cct tgc atc gtc gca tcc ggc gcc	739
Ile Ile Ala Arg Gln Leu Asn Val Pro Cys Ile Val Ala Ser Gly Ala	
200 205 210	
ggc atc aag gac atc aag tcc ggc gaa aag gtg ctt atc gac ggc agc	787
Gly Ile Lys Asp Ile Lys Ser Gly Glu Lys Val Leu Ile Asp Gly Ser	
215 220 225	
ctc ggc acc att gac cgc aac gcg gac gaa gct gaa gca acc aag ctc	835
Leu Gly Thr Ile Asp Arg Asn Ala Asp Glu Ala Glu Ala Thr Lys Leu	
230 235 240 245	
gtc tcc gag tcc ctc gag cgc gct gct cgc atc gcc gag tgg aag ggt	883
Val Ser Glu Ser Leu Glu Arg Ala Ala Arg Ile Ala Glu Trp Lys Gly	
250 255 260	
cct gca caa acc aag gac ggc tac cgc gtt cag ctg ttg gcc aac gtc	931
Pro Ala Gln Thr Lys Asp Gly Tyr Arg Val Gln Leu Leu Ala Asn Val	
265 270 275	
caa gac ggc aac tct gca cag cag gct gca cag acc gaa gca gaa ggc	979
Gln Asp Gly Asn Ser Ala Gln Gln Ala Ala Gln Thr Glu Ala Glu Gly	
280 285 290	
atc ggc ctg ttc cgc acc gaa ctg tgc ttc ctt tcc gcc acc gaa gag	
1027 Ile Gly Leu Phe Arg Thr Glu Leu Cys Phe Leu Ser Ala Thr Glu Glu	
295 300 305	
cca agc gtt gat gag cag gct gcg gtc tac tca aag gtg ctt gaa gca	
1075 Pro Ser Val Asp Glu Gln Ala Ala Val Tyr Ser Lys Val Leu Glu Ala	
310 315 320 325	

BGI-122CP

ttc cca gag tcc aag gtc gtt gtc cgc tcc ctc gac gca ggt tct gac
 1123
 Phe Pro Glu Ser Lys Val Val Val Arg Ser Leu Asp Ala Gly Ser Asp
 330 335 340

aag cca gtt cca ttc gca tgc atg gct gat gag atg aac cca gca ctg
 1171
 Lys Pro Val Pro Phe Ala Ser Met Ala Asp Glu Met Asn Pro Ala Leu
 345 350 355

ggt gtt cgt ggc ctg cgt atc gca cgt gga cag gtt gat ctg ctg act
 1219
 Gly Val Arg Gly Leu Arg Ile Ala Arg Gly Gln Val Asp Leu Leu Thr
 360 365 370

cgc cag ctc gac gca att gcg aag gcc agc gaa gaa ctc ggc cgt ggc
 1267
 Arg Gln Leu Asp Ala Ile Ala Lys Ala Ser Glu Glu Leu Gly Arg Gly
 375 380 385

gac gac gcc cca acc tgg gtt atg gct cca atg gtg gct acc gct tat
 1315
 Asp Asp Ala Pro Thr Trp Val Met Ala Pro Met Val Ala Thr Ala Tyr
 390 395 400 405

gaa gca aag tgg ttt gct gac atg tgc cgt gag cgt ggc cta atc gcc
 1363
 Glu Ala Lys Trp Phe Ala Asp Met Cys Arg Glu Arg Gly Leu Ile Ala
 410 415 420

ggc gcc atg atc gaa gtt cca gca gca tcc ctg atg gca gac aag atc
 1411
 Gly Ala Met Ile Glu Val Pro Ala Ala Ser Leu Met Ala Asp Lys Ile
 425 430 435

atg cct cac ctg gac ttt gtt tcc atc ggt acc aac gac ctg acc cag
 1459
 Met Pro His Leu Asp Phe Val Ser Ile Gly Thr Asn Asp Leu Thr Gln
 440 445 450

tac acc atg gca gcg gac cgc atg tct cct gag ctt gcc tac ctg acc
 1507
 Tyr Thr Met Ala Ala Asp Arg Met Ser Pro Glu Leu Ala Tyr Leu Thr
 455 460 465

gat cct tgg cag cca gca gtc ctg cgc ctg atc aag cac acc tgt gac
 1555
 Asp Pro Trp Gln Pro Ala Val Leu Arg Leu Ile Lys His Thr Cys Asp
 470 475 480 485

gaa ggt gct cgc ttt aac acc cgg gtc ggt gtt tgt ggt gaa gca gca
 1603
 Glu Gly Ala Arg Phe Asn Thr Pro Val Gly Val Cys Gly Glu Ala Ala
 490 495 500

gca gac cca ctg ttg gca act gtc ctc acc ggt ctt ggc gtg aac tcc
 1651
 Ala Asp Pro Leu Leu Ala Thr Val Leu Thr Gly Leu Gly Val Asn Ser
 505 510 515

00604231.062700

ctg tcc gca gca tcc act gct ctc gca gca gtc ggt gca aag ctg tca
 1699
 Leu Ser Ala Ala Ser Thr Ala Leu Ala Ala Val Gly Ala Lys Leu Ser
 520 525 530
 gag gtc acc ctg gaa acc tgt aag aag gca gca gaa gca gca ctt gac
 1747
 Glu Val Thr Leu Glu Thr Cys Lys Lys Ala Ala Glu Ala Ala Leu Asp
 535 540 545
 gct gaa ggt gca act gaa gca cgc gat gct gta cgc gca gtg atc gac
 1795
 Ala Glu Gly Ala Thr Glu Ala Arg Asp Ala Val Arg Ala Val Ile Asp
 550 555 560 565
 gca gca gtc taaaccactg ttgagctaaa aag
 1827
 Ala Ala Val

```
<210> 18
<211> 568
<212> PRT
<213> Corynebacterium glutamicum
```

400> 18															
Val	Ala	Thr	Val	Ala	Asp	Val	Asn	Gln	Asp	Thr	Val	Leu	Lys	Gly	Thr
1				5				10						15	
Gly	Val	Val	Gly	Gly	Val	Arg	Tyr	Ala	Ser	Ala	Val	Trp	Ile	Thr	Pro
			20					25					30		
Arg	Pro	Glu	Leu	Pro	Gln	Ala	Gly	Glu	Val	Val	Ala	Glu	Glu	Asn	Arg
		35				40						45			
Glu	Ala	Glu	Gln	Glu	Arg	Phe	Asp	Ala	Ala	Ala	Ala	Thr	Val	Ser	Ser
	50					55					60				
Arg	Leu	Leu	Glu	Arg	Ser	Glu	Ala	Ala	Glu	Gly	Pro	Ala	Ala	Glu	Val
65					70					75				80	
Leu	Lys	Ala	Thr	Ala	Gly	Met	Val	Asn	Asp	Arg	Gly	Trp	Arg	Lys	Ala
				85					90					95	
Val	Ile	Lys	Gly	Val	Lys	Gly	Gly	His	Pro	Ala	Glu	Tyr	Ala	Val	Val
			100					105					110		
Ala	Ala	Thr	Thr	Lys	Phe	Ile	Ser	Met	Phe	Glu	Ala	Ala	Gly	Gly	Leu
		115					120					125			
Ile	Ala	Glu	Arg	Thr	Thr	Asp	Leu	Arg	Asp	Ile	Arg	Asp	Arg	Val	Ile
130						135					140				
Ala	Glu	Leu	Arg	Gly	Asp	Glu	Glu	Pro	Gly	Leu	Pro	Ala	Val	Ser	Gly
145					150					155				160	
Gln	Val	Ile	Leu	Phe	Ala	Asp	Asp	Leu	Ser	Pro	Ala	Asp	Thr	Ala	Ala
				165					170					175	
Leu	Asp	Thr	Asp	Leu	Phe	Val	Gly	Leu	Val	Thr	Glu	Leu	Gly	Gly	Pro

[illegible]

Leu Gly Val Asn Ser Leu Ser Ala Ala Ser Thr Ala Leu Ala Ala Val
515 520 525

Gly Ala Lys Leu Ser Glu Val Thr Leu Glu Thr Cys Lys Lys Ala Ala
530 535 540

Glu Ala Ala Leu Asp Ala Glu Gly Ala Thr Glu Ala Arg Asp Ala Val
545 550 555 560

Arg Ala Val Ile Asp Ala Ala Val
565

<210> 19

<211> 1629

<212> DNA

<213> Corynebacterium glutamicum

<220>

<221> CDS

<222> (98)..(1606)

<223> FRXA01244

<400> 19

agatgtcgcgatttctcgagga agaagttaac gccgaagaaa accgtgaatc agagcaggag 60

cgttcgcagc ccgtgcgacg cacagttctt tcttcgtttg ctt gag cgc tcc gaa 115
Leu Leu Glu Arg Ser Glu
1 5

gct gct gaa gga cca gca gct gag gtg ctt aaa gct act gct ggc atg 163
Ala Ala Glu Gly Pro Ala Ala Glu Val Leu Lys Ala Thr Ala Gly Met
10 15 20

gtc aat gac cgt ggc tgg cgt aag gct gtc atc aag ggt gtc aag ggt 211
Val Asn Asp Arg Gly Trp Arg Lys Ala Val Ile Lys Gly Val Lys Gly
25 30 35

ggt cac cct gcg gaa tac gcc gtg gtt gca gca aca acc aag ttc atc 259
Gly His Pro Ala Glu Tyr Ala Val Val Ala Ala Thr Thr Lys Phe Ile
40 45 50

tcc atg ttc gaa gcc gca ggc ggc ctg atc gcg gag cgc acc aca gac 307
Ser Met Phe Glu Ala Ala Gly Gly Leu Ile Ala Glu Arg Thr Thr Asp
55 60 65 70

ttg cgc gac atc cgc gac cgc gtc atc gca gaa ctt cgt ggc gat gaa 355
Leu Arg Asp Ile Arg Asp Arg Val Ile Ala Glu Leu Arg Gly Asp Glu
75 80 85

gag cca ggt ctg cca gct gtt tcc gga cag gtc att ctc ttt gca gat 403
Glu Pro Gly Leu Pro Ala Val Ser Gly Gln Val Ile Leu Phe Ala Asp
90 95 100

gac ctc tcc cca gca gac acc gcg gca cta gac aca gat ctc ttt gtg 451
Asp Leu Ser Pro Ala Asp Thr Ala Ala Leu Asp Thr Asp Leu Phe Val
105 110 115

gga ctt gtc act gag ctg ggt ggc cca acg agc cac acc gcg atc atc 499
Gly Leu Val Thr Glu Leu Gly Gly Pro Thr Ser His Thr Ala Ile Ile

00604231-002700

120	125	130	
gca cgc cag ctc aac gtg cct tgc atc gtc gca tcc ggc gcc ggc atc Ala Arg Gln Leu Asn Val Pro Cys Ile Val Ala Ser Gly Ala Gly Ile 135 140 145 150			547
aag gac atc aag tcc ggc gaa aag gtg ctt atc gac ggc agc ctc ggc Lys Asp Ile Lys Ser Gly Glu Lys Val Leu Ile Asp Gly Ser Leu Gly 155 160 165			595
acc att gac cgc aac gcg gac gaa gct gaa gca acc aag ctc gtc tcc Thr Ile Asp Arg Asn Ala Asp Glu Ala Glu Ala Thr Lys Leu Val Ser 170 175 180			643
gag tcc ctc gag cgc gct gct cgc atc gcc gag tgg aag ggt cct gca Glu Ser Leu Glu Arg Ala Ala Arg Ile Ala Glu Trp Lys Gly Pro Ala 185 190 195			691
caa acc aag gac ggc tac cgc gtt cag ctg ttg gcc aac gtc caa gac Gln Thr Lys Asp Gly Tyr Arg Val Gln Leu Leu Ala Asn Val Gln Asp 200 205 210			739
ggc aac tct gca cag cag gct gca cag acc gaa gca gaa ggc atc ggc Gly Asn Ser Ala Gln Gln Ala Ala Gln Thr Glu Ala Glu Gly Ile Gly 215 220 225 230			787
ctg ttc cgc acc gaa ctg tgc ttc ctt tcc gcc acc gaa gag cca agc Leu Phe Arg Thr Glu Leu Cys Phe Leu Ser Ala Thr Glu Glu Pro Ser 235 240 245			835
gtt gat gag cag gct gcg gtc tac tca aag gtg ctt gaa gca ttc cca Val Asp Glu Gln Ala Ala Val Tyr Ser Lys Val Leu Glu Ala Phe Pro 250 255 260			883
gag tcc aag gtc gtt gtc cgc tcc ctc gac gca ggt tct gac aag cca Glu Ser Lys Val Val Val Arg Ser Leu Asp Ala Gly Ser Asp Lys Pro 265 270 275			931
gtt cca ttc gca tcg atg gct gat gag atg aac cca gca ctg ggt gtt Val Pro Phe Ala Ser Met Ala Asp Glu Met Asn Pro Ala Leu Gly Val 280 285 290			979
cgt ggc ctg cgt atc gca cgt gga cag gtt gat ctg ctg act cgc cag 1027 Arg Gly Leu Arg Ile Ala Arg Gly Gln Val Asp Leu Leu Thr Arg Gln 295 300 305 310			
ctc gac gca att gcg aag gcc agc gaa gaa ctc ggc cgt ggc gac gac 1075 Leu Asp Ala Ile Ala Lys Ala Ser Glu Glu Leu Gly Arg Gly Asp Asp 315 320 325			
gcc cca acc tgg gtt atg gct cca atg gtg gct acc gct tat gaa gca 1123 Ala Pro Thr Trp Val Met Ala Pro Met Val Ala Thr Ala Tyr Glu Ala 330 335 340			
aag tgg ttt gct gac atg tgc cgt gag cgt ggc cta atc gcc ggc gcc 1171 Lys Trp Phe Ala Asp Met Cys Arg Glu Arg Gly Leu Ile Ala Gly Ala 345 350 355			

BGI-122CP-062700

atg atc gaa gtt cca gca gca tcc ctg atg gca gac aag atc atg cct
1219

Met Ile Glu Val Pro Ala Ala Ser Leu Met Ala Asp Lys Ile Met Pro
360 365 370

cac ctg gac ttt gtt tcc atc ggt acc aac gac ctg acc cag tac acc
1267

His Leu Asp Phe Val Ser Ile Gly Thr Asn Asp Leu Thr Gln Tyr Thr
375 380 385 390

atg gca gcg gac cgc atg tct cct gag ctt gcc tac ctg acc gat cct
1315

Met Ala Ala Asp Arg Met Ser Pro Glu Leu Ala Tyr Leu Thr Asp Pro
395 400 405

tggt cag cca gca gtc ctg cgc ctg atc aag cac acc tgt gac gaa ggt
1363

Trp Gln Pro Ala Val Leu Arg Leu Ile Lys His Thr Cys Asp Glu Gly
410 415 420

gct cgc ttt aac acc ccg gtc ggt gtt tgt ggt gaa gca gca gca gac
1411

Ala Arg Phe Asn Thr Pro Val Gly Val Cys Gly Glu Ala Ala Ala Asp
425 430 435

cca ctg ttg gca act gtc ctc acc ggt ctt gcc gtg aac tcc ctg tcc
1459

Pro Leu Leu Ala Thr Val Leu Thr Gly Leu Gly Val Asn Ser Leu Ser
440 445 450

gca gca tcc act gct ctc gca gca gtc ggt gca aag ctg tca gag gtc
1507

Ala Ala Ser Thr Ala Leu Ala Ala Val Gly Ala Lys Leu Ser Glu Val
455 460 465 470

acc ctg gaa acc tgt aag aag gca gca gaa gca gca ctt gac gct gaa
1555

Thr Leu Glu Thr Cys Lys Lys Ala Ala Glu Ala Ala Leu Asp Ala Glu
475 480 485

ggt gca act gaa gca cgc gat gct gta cgc gca gtg atc gac gca gca
1603

Gly Ala Thr Glu Ala Arg Asp Ala Val Arg Ala Val Ile Asp Ala Ala
490 495 500

gtc taaaccactg ttgagctaaa aag
1629

Val

<210> 20

<211> 503

<212> PRT

<213> Corynebacterium glutamicum

<400> 20

Leu Leu Glu Arg Ser Glu Ala Ala Glu Gly Pro Ala Ala Glu Val Leu
1 5 10 15

00001-00700

Lys Ala Thr Ala Gly Met Val Asn Asp Arg Gly Trp Arg Lys Ala Val
 20 25 30
 Ile Lys Gly Val Lys Gly Gly His Pro Ala Glu Tyr Ala Val Val Ala
 35 40 45
 Ala Thr Thr Lys Phe Ile Ser Met Phe Glu Ala Ala Gly Gly Leu Ile
 50 55 60
 Ala Glu Arg Thr Thr Asp Leu Arg Asp Ile Arg Asp Arg Val Ile Ala
 65 70 75 80
 Glu Leu Arg Gly Asp Glu Glu Pro Gly Leu Pro Ala Val Ser Gly Gln
 85 90 95
 Val Ile Leu Phe Ala Asp Asp Leu Ser Pro Ala Asp Thr Ala Ala Leu
 100 105 110
 Asp Thr Asp Leu Phe Val Gly Leu Val Thr Glu Leu Gly Gly Pro Thr
 115 120 125
 Ser His Thr Ala Ile Ile Ala Arg Gln Leu Asn Val Pro Cys Ile Val
 130 135 140
 Ala Ser Gly Ala Gly Ile Lys Asp Ile Lys Ser Gly Glu Lys Val Leu
 145 150 155 160
 Ile Asp Gly Ser Leu Gly Thr Ile Asp Arg Asn Ala Asp Glu Ala Glu
 165 170 175
 Ala Thr Lys Leu Val Ser Glu Ser Leu Glu Arg Ala Ala Arg Ile Ala
 180 185 190
 Glu Trp Lys Gly Pro Ala Gln Thr Lys Asp Gly Tyr Arg Val Gln Leu
 195 200 205
 Leu Ala Asn Val Gln Asp Gly Asn Ser Ala Gln Gln Ala Ala Gln Thr
 210 215 220
 Glu Ala Glu Gly Ile Gly Leu Phe Arg Thr Glu Leu Cys Phe Leu Ser
 225 230 235 240
 Ala Thr Glu Glu Pro Ser Val Asp Glu Gln Ala Ala Val Tyr Ser Lys
 245 250 255
 Val Leu Glu Ala Phe Pro Glu Ser Lys Val Val Val Arg Ser Leu Asp
 260 265 270
 Ala Gly Ser Asp Lys Pro Val Pro Phe Ala Ser Met Ala Asp Glu Met
 275 280 285
 Asn Pro Ala Leu Gly Val Arg Gly Leu Arg Ile Ala Arg Gly Gln Val
 290 295 300
 Asp Leu Leu Thr Arg Gln Leu Asp Ala Ile Ala Lys Ala Ser Glu Glu
 305 310 315 320
 Leu Gly Arg Gly Asp Asp Ala Pro Thr Trp Val Met Ala Pro Met Val
 325 330 335
 Ala Thr Ala Tyr Glu Ala Lys Trp Phe Ala Asp Met Cys Arg Glu Arg

002280.122400

		340						345							350				
Gly	Leu	Ile	Ala	Gly	Ala	Met	Ile	Glu	Val	Pro	Ala	Ala	Ser	Leu	Met				
		355					360						365						
Ala	Asp	Lys	Ile	Met	Pro	His	Leu	Asp	Phe	Val	Ser	Ile	Gly	Thr	Asn				
		370				375					380								
Asp	Leu	Thr	Gln	Tyr	Thr	Met	Ala	Ala	Asp	Arg	Met	Ser	Pro	Glu	Leu				
		385			390					395					400				
Ala	Tyr	Leu	Thr	Asp	Pro	Trp	Gln	Pro	Ala	Val	Leu	Arg	Leu	Ile	Lys				
				405					410					415					
His	Thr	Cys	Asp	Glu	Gly	Ala	Arg	Phe	Asn	Thr	Pro	Val	Gly	Val	Cys				
			420					425					430						
Gly	Glu	Ala	Ala	Ala	Asp	Pro	Leu	Leu	Ala	Thr	Val	Leu	Thr	Gly	Leu				
		435					440					445							
Gly	Val	Asn	Ser	Leu	Ser	Ala	Ala	Ser	Thr	Ala	Leu	Ala	Ala	Val	Gly				
		450				455					460								
Ala	Lys	Leu	Ser	Glu	Val	Thr	Leu	Glu	Thr	Cys	Lys	Lys	Ala	Ala	Glu				
		465			470					475					480				
Ala	Ala	Leu	Asp	Ala	Glu	Gly	Ala	Thr	Glu	Ala	Arg	Asp	Ala	Val	Arg				
			485						490					495					
Ala	Val	Ile	Asp	Ala	Ala	Val													
			500																

<210> 21
<211> 390
<212> DNA
<213> Corynebacterium glutamicum

<220>
<221> CDS
<222> (101)..(367)
<223> RXA01300

<400> 21
gatcgacatt aaatcccctc ctttggggggt tttaactaac aaatcgtgc gccctaatacc 60
gttcggatta acgccgtagc aacacgaaag gacaccttcc atg gct tcc aag act 115
Met Ala Ser Lys Thr
1 5
gta acc gtc ggt tcc tcc gtt ggc ctg cac gca cgt cca gca tcc atc 163
Val Thr Val Gly Ser Ser Val Gly Leu His Ala Arg Pro Ala Ser Ile
10 15 20
atc gct gaa gcg gct gct gag tac gac gac gaa atc ttg ctg acc ctg 211
Ile Ala Glu Ala Ala Ala Glu Tyr Asp Asp Glu Ile Leu Leu Thr Leu
25 30 35
gtt ggc tcc gat gat gac gaa gag acc gac gcg tcc tct tcc ctc atg 259
Val Gly Ser Asp Asp Asp Glu Glu Thr Asp Ala Ser Ser Ser Leu Met
40 45 50

[illegible]

```
<210> 22
<211> 89
<212> PRT
<213> Corynebacterium glutamicum
```

```

400> 22
Met Ala Ser Lys Thr Val Thr Val Gly Ser Ser Val Gly Leu His Ala
  1          5          10          15
Arg Pro Ala Ser Ile Ile Ala Glu Ala Ala Ala Glu Tyr Asp Asp Glu
          20          25          30
Ile Leu Leu Thr Leu Val Gly Ser Asp Asp Asp Glu Glu Thr Asp Ala
  35          40          45
Ser Ser Ser Leu Met Ile Met Ala Leu Gly Ala Glu His Gly Asn Glu
  50          55          60
Val Thr Val Thr Ser Asp Asn Ala Glu Ala Val Glu Lys Ile Ala Ala
  65          70          75          80
Leu Ile Ala Gln Asp Leu Asp Ala Glu
          85

```

```
<210> 23
<211> 508
<212> DNA
<213> Corynebacterium glutamicum
```

```
<220>
<221> CDS
<222> (101)..(508)
<223> RXN03002
```

```

400> 23
ggaacttcga ggtgtcttcg tggggcgctac ggagatctag caagtgtgac tttatgtttg 60
accctatcgc aatcaacatg cagtgaatta acatctactt atg ttt gta ctc aaa 115
                                         Met Phe Val Leu Lys
                                         1                               5

gat ctg cta aag gca gaa cgc ata gaa ctc gac cgc acg gtc acc gat 163
Asp Leu Leu Lys Ala Glu Arg Ile Glu Leu Asp Arg Thr Val Thr Asp
                        10                               15                               20

tgg cgt gaa qgc atc cgc qcc qca qgt gta ctc cta gaa aag aca aac 211

```

Trp	Arg	Glu	Gly	Ile	Arg	Ala	Ala	Gly	Val	Leu	Leu	Glu	Lys	Thr	Asn	
			25					30					35			
agc	att	gat	tcc	gcc	tac	acc	gat	gcc	atg	atc	gcc	agc	gtg	gaa	gaa	259
Ser	Ile	Asp		Ser	Ala	Tyr	Thr	Asp	Ala	Met	Ile	Ala	Ser	Val	Glu	
		40					45					50				
aaa	ggc	ccc	tac	att	gtg	gtc	gct	cca	ggt	ttc	gct	ttc	gcg	cac	gcc	307
Lys	Gly	Pro	Tyr	Ile	Val	Val	Ala	Pro	Gly	Phe	Ala	Phe	Ala	His	Ala	
	55				60					65						
cgc	ccc	agc	aga	gca	gtc	cgc	gag	acc	gct	atg	tcg	tggtg	gcg	cgc	ctg	355
Arg	Pro	Ser	Arg	Ala	Val	Arg	Glu	Thr	Ala	Met	Ser	Trp	Val	Arg	Leu	
70					75					80					85	
gcc	tcc	cct	gtt	tcc	ttc	ggt	cac	agt	aag	aat	gat	ccc	ctc	aat	ctc	403
Ala	Ser	Pro	Val	Ser	Phe	Gly	His	Ser	Lys	Asn	Asp	Pro	Leu	Asn	Leu	
				90					95					100		
atc	gtt	gct	ctc	gct	gcc	aaa	gat	gcc	acc	gca	cat	acc	caa	gcg	atg	451
Ile	Val	Ala	Leu	Ala	Ala	Lys	Asp	Ala	Thr	Ala	His	Thr	Gln	Ala	Met	
		105						110					115			
gcg	gca	ttg	gct	aaa	gct	tta	gga	aaa	tac	cga	aag	gat	ctc	gac	gag	499
Ala	Ala	Leu	Ala	Lys	Ala	Leu	Gly	Lys	Tyr	Arg	Lys	Asp	Leu	Asp	Glu	
		120				125						130				
gca	caa	agt														508
Ala	Gln	Ser														
		135														
<210> 24																
<211> 136																
<212> PRT																
<213> Corynebacterium glutamicum																
<400> 24																
Met	Phe	Val	Leu	Lys	Asp	Leu	Leu	Lys	Ala	Glu	Arg	Ile	Glu	Leu	Asp	
1				5					10					15		
Arg	Thr	Val	Thr	Asp	Trp	Arg	Glu	Gly	Ile	Arg	Ala	Ala	Gly	Val	Leu	
		20						25					30			
Leu	Glu	Lys	Thr	Asn	Ser	Ile	Asp	Ser	Ala	Tyr	Thr	Asp	Ala	Met	Ile	
		35					40					45				
Ala	Ser	Val	Glu	Glu	Lys	Gly	Pro	Tyr	Ile	Val	Val	Ala	Pro	Gly	Phe	
	50					55					60					
Ala	Phe	Ala	His	Ala	Arg	Pro	Ser	Arg	Ala	Val	Arg	Glu	Thr	Ala	Met	
	65				70					75					80	
Ser	Trp	Val	Arg	Leu	Ala	Ser	Pro	Val	Ser	Phe	Gly	His	Ser	Lys	Asn	
			85					90						95		
Asp	Pro	Leu	Asn	Leu	Ile	Val	Ala	Leu	Ala	Ala	Lys	Asp	Ala	Thr	Ala	
		100						105					110			
His	Thr	Gln	Ala	Met	Ala	Ala	Leu	Ala	Lys	Ala	Leu	Gly	Lys	Tyr	Arg	
		115				120					125					

Lys Asp Leu Asp Glu Ala Gln Ser
130 135

<210> 25
<211> 789
<212> DNA
<213> *Corynebacterium glutamicum*

<220>
<221> CDS
<222> (14)..(766)
<223> RXC00953

<400> 25
cttgcatgcc ccaatg gcg cca cca acg gta ggc aac tac atc atg cag tcc 52
Met Ala Pro Pro Thr Val Gly Asn Tyr Ile Met Gln Ser
1 5 10

ttc act caa ggt ctg cag ttc ggc gtt gca gtt gcc gtg att ctc ttt 100
Phe Thr Gln Gly Leu Gln Phe Gly Val Ala Val Ala Val Ile Leu Phe
15 20 25

ggg gtc cgc acc att ctt ggt gaa ctg gtc ccc gca ttc caa ggt att 148
Gly Val Arg Thr Ile Leu Gly Glu Leu Val Pro Ala Phe Gln Gly Ile
30 35 40 45

gct gcg aag gtt gtt ccc gga gct atc ccc gca ttg gat gca cgg atc 196
Ala Ala Lys Val Val Pro Gly Ala Ile Pro Ala Leu Asp Ala Pro Ile
50 55 60

gtg ttc ccc tac gcg cag aac gcc gtt ctc att ggt ttc ttg tct tcc 244
Val Phe Pro Tyr Ala Gln Asn Ala Val Leu Ile Gly Phe Leu Ser Arg
65 70 75

ttc gtc ggt gcc ttg gtt gcc ctg act gtt ctt gca tcg tgg ctg aac 292
Phe Val Gly Gly Leu Val Gly Leu Thr Val Leu Ala Ser Trp Leu Asn
80 85 90

cca gct ttt ggt gtc gcg ttg att ctg cct ggt ttg gtc ccc cac ttc 340
Pro Ala Phe Gly Val Ala Leu Ile Leu Pro Gly Leu Val Pro His Phe
95 100 105

ttc act ggt gcc gcg gcg gcc gtt tac ggt aat gcc acg ggt ggt cgt 388
Phe Thr Gly Gly Ala Ala Gly Val Tyr Gly Asn Ala Thr Gly Gly Arg
110 115 120 125

cga gga gca gta ttt gcc gcc ttt gcc aac ggt ctt ctg att acc ttc 436
Arg Gly Ala Val Phe Gly Ala Phe Ala Asn Gly Leu Leu Ile Thr Phe
130 135 140

ctc cct gct ttc ctg ctt ggt gtg ctt ggt tcc ttc ggg tca gag aac 484
Leu Pro Ala Phe Leu Leu Gly Val Leu Gly Ser Phe Gly Ser Glu Asn
145 150 155

acc act ttc ggt gat gcg gac ttt ggt tgg ttc gga atc gtt gtt ggt 532
Thr Thr Phe Gly Asp Ala Asp Phe Gly Trp Phe Gly Ile Val Val Gly
160 165 170

tct gca gcc aag gtg gaa ggt gct ggc ggg ctc atc ttg ttg ctc atc 580

002299-12240960

[illegible]

```

<210> 26
<211> 251
<212> PRT
<213> Corynebacterium glutamicum

<400> 26
Met Ala Pro Pro Thr Val Gly Asn Tyr Ile Met Gln Ser Phe Thr Gln
  1          5          10          15
Gly Leu Gln Phe Gly Val Ala Val Ala Val Ile Leu Phe Gly Val Arg
          20          25          30
Thr Ile Leu Gly Glu Leu Val Pro Ala Phe Gln Gly Ile Ala Ala Lys
          35          40          45
Val Val Pro Gly Ala Ile Pro Ala Leu Asp Ala Pro Ile Val Phe Pro
          50          55          60
Tyr Ala Gln Asn Ala Val Leu Ile Gly Phe Leu Ser Ser Phe Val Gly
  65          70          75          80
Gly Leu Val Gly Leu Thr Val Leu Ala Ser Trp Leu Asn Pro Ala Phe
          85          90          95
Gly Val Ala Leu Ile Leu Pro Gly Leu Val Pro His Phe Phe Thr Gly
          100          105          110
Gly Ala Ala Gly Val Tyr Gly Asn Ala Thr Gly Gly Arg Arg Gly Ala
          115          120          125
Val Phe Gly Ala Phe Ala Asn Gly Leu Leu Ile Thr Phe Leu Pro Ala
          130          135          140
Phe Leu Leu Gly Val Leu Gly Ser Phe Gly Ser Glu Asn Thr Thr Phe
          145          150          155          160
Gly Asp Ala Asp Phe Gly Trp Phe Gly Ile Val Val Gly Ser Ala Ala
          165          170          175

```

Lys Val Glu Gly Ala Gly Gly Leu Ile Leu Leu Leu Ile Ile Ala Ala
 180 185 190
 Val Leu Leu Gly Gly Ala Met Val Phe Gln Lys Arg Val Val Asn Gly
 195 200 205
 His Trp Asp Pro Ala Pro Asn Arg Glu Arg Val Glu Lys Ala Glu Ala
 210 215 220
 Asp Ala Thr Pro Thr Ala Gly Ala Arg Thr Tyr Pro Lys Ile Ala Pro
 225 230 235 240
 Pro Ala Gly Ala Pro Thr Pro Pro Ala Arg Ser
 245 250

<210> 27
 <211> 553
 <212> DNA
 <213> *Corynebacterium glutamicum*

<220>
 <221> CDS
 <222> (101)..(553)
 <223> RXC03001

<400> 27
 cccggttcac gtgatcaatg acttcacgag caccgatgaa atcgatgctg cgcttcgtga 60
 acgctacgac atctaactac tttaaaagga cgaaaaatatt atg gac tgg tta acc 115
 Met Asp Trp Leu Thr
 1 5
 att cct ott ttc ctc gtt aat gaa atc ctt gcg gtt ccg gct ttc ctc 163
 Ile Pro Leu Phe Leu Val Asn Glu Ile Leu Ala Val Pro Ala Phe Leu
 10 15 20
 atc ggt atc atc acc gcc gtg gga ttg ggt gcc atg ggg cgt tcc gtc 211
 Ile Gly Ile Ile Thr Ala Val Gly Leu Gly Ala Met Gly Arg Ser Val
 25 30 35
 ggt cag gtt atc ggt gga gca atc aaa gca acg ttg ggc ttt ttg ctc 259
 Gly Gln Val Ile Gly Gly Ala Ile Lys Ala Thr Leu Gly Phe Leu Leu
 40 45 50
 att ggt gcg ggt gcc acg ttg gtc act gcc tcc ctg gag cca ctg ggt 307
 Ile Gly Ala Gly Ala Thr Leu Val Thr Ala Ser Leu Glu Pro Leu Gly
 55 60 65
 gcg atg atc atg ggt gcc aca gcc atg cgt ggt gtt gtc cca acg aat 355
 Ala Met Ile Met Gly Ala Thr Gly Met Arg Gly Val Val Pro Thr Asn
 70 75 80 85
 gaa gcc atc gcc gga atc gca cag gct gaa tac ggc gcg cag gtg gcg 403
 Glu Ala Ile Ala Gly Ile Ala Gln Ala Glu Tyr Gly Ala Gln Val Ala
 90 95 100
 tgg ctg atg att ctg gcc ttc gcc atc tct ttg gtg ttg gct cgt ttc 451
 Trp Leu Met Ile Leu Gly Phe Ala Ile Ser Leu Val Leu Ala Arg Phe
 105 110 115

002390 1 553700

acc aac ctg cgt tat gtc ttg ctc aac gga cac cac gtg ctg ttg atg 499
Thr Asn Leu Arg Tyr Val Leu Leu Asn Gly His His Val Leu Leu Met
120 125 130

tgc acc atg ctc acc atg gtc ttg gcc acc gga aga gtt gat gcg tgg 547
Cys Thr Met Leu Thr Met Val Leu Ala Thr Gly Arg Val Asp Ala Trp
135 140 145

atc ttc 553
Ile Phe
150

<210> 28

<211> 15

<212> PRT

<213> Corynebacterium glutamicum

<400> 28

Met Asp Trp Leu Thr Ile Pro Leu Phe Leu Val Asn Glu Ile Leu Ala
1 5 10 15

Val Pro Ala Phe Leu Ile Gly Ile Ile Thr Ala Val Gly Leu Gly Ala
20 25 30

Met Gly Arg Ser Val Gly Gln Val Ile Gly Gly Ala Ile Lys Ala Thr
35 40 45

Leu Gly Phe Leu Leu Ile Gly Ala Gly Ala Thr Leu Val Thr Ala Ser
50 55 60

Leu Glu Pro Leu Gly Ala Met Ile Met Gly Ala Thr Gly Met Arg Gly
65 70 75 80

Val Val Pro Thr Asn Glu Ala Ile Ala Gly Ile Ala Gln Ala Glu Tyr
85 90 95

Gly Ala Gln Val Ala Trp Leu Met Ile Leu Gly Phe Ala Ile Ser Leu
100 105 110

Val Leu Ala Arg Phe Thr Asn Leu Arg Tyr Val Leu Leu Asn Gly His
115 120 125

His Val Leu Leu Met Cys Thr Met Leu Thr Met Val Leu Ala Thr Gly
130 135 140

Arg Val Asp Ala Trp Ile Phe
145 150

<210> 29

<211> 2172

<212> DNA

<213> *Corynebacterium glutamicum*

 $\langle 220 \rangle$

<221> CDS

<222> (101) . . (2149)

<223> RXN01943

<400> 29

ccgattctttt ttccgcccacaa ttccgtaacgg cgcctcctctt aagtgggacaa gaaagtctct 60

tgccccgcggg agacagaccc tacgtttaga aagggttgac atg gcg tcc aaa ctg 115
Met Ala Ser Lys Leu
1 5

acg acg aca tcg caa cat att ctg gaa aac ctt ggt gga cca gac aat 163
Thr Thr Thr Ser Gln His Ile Leu Glu Asn Leu Gly Gly Pro Asp Asn
10 15 20

att act tcg atg act cac tgt gcg act cgc ctt cgc ttc caa gtg aag 211
Ile Thr Ser Met Thr His Cys Ala Thr Arg Leu Arg Phe Gln Val Lys
25 30 35

gat caa tcc att gtt gat caa caa gaa att gac tcc gac cca tca gtt 259
Asp Gln Ser Ile Val Asp Gln Gln Glu Ile Asp Ser Asp Pro Ser Val
40 45 50

ctt ggc gta gta ccc caa gga tcc acc ggt atg cag gtg gtg atg ggt 307
Leu Gly Val Val Pro Gln Gly Ser Thr Gly Met Gln Val Val Met Gly
55 60 65

gga tct gtt gca aac tat tac caa gaa atc ctc aaa ctt gat gga atg 355
Gly Ser Val Ala Asn Tyr Tyr Gln Glu Ile Leu Lys Leu Asp Gly Met
70 75 80 85

aag cac ttc gcc gac ggt gaa gct aca gag agt tca tcc aag aag gaa 403
Lys His Phe Ala Asp Gly Glu Ala Thr Glu Ser Ser Ser Lys Lys Glu
90 95 100

tac ggc gga gtc cgt ggc aag tac tcg tgg att gac tac gcc ttc gag 451
Tyr Gly Gly Val Arg Gly Lys Tyr Ser Trp Ile Asp Tyr Ala Phe Glu
105 110 115

ttc ttg tct gat act ttc cga cca atc ctg tgg gcc ctg ctt ggt gcc 499
Phe Leu Ser Asp Thr Phe Arg Pro Ile Leu Trp Ala Leu Leu Gly Ala
120 125 130

tca ctg att att acc ttg ttg gtt ctt gcg gat act ttc ggt ttg caa 547
Ser Leu Ile Ile Thr Leu Val Leu Ala Asp Thr Phe Gly Leu Gln
135 140 145

gac ttc cgc gct cca atg gat gag cag cct gat act tat gta ttc ctg 595
Asp Phe Arg Ala Pro Met Asp Glu Gln Pro Asp Thr Tyr Val Phe Leu
150 155 160 165

cac tcc atg tgg cgc tcg gtc ttc tac ttc ctg cca att atg gtt ggt 643
His Ser Met Trp Arg Ser Val Phe Tyr Phe Leu Pro Ile Met Val Gly
170 175 180

gcc acc gca gct cga aag ctc ggc gca aac gag tgg att ggt gca gct 691
Ala Thr Ala Ala Arg Lys Leu Gly Ala Asn Glu Trp Ile Gly Ala Ala
185 190 195

att cca gcc gca ctt ctt act cca gaa ttc ttg gca ctg ggt tct gcc 739
Ile Pro Ala Ala Leu Leu Thr Pro Glu Phe Leu Ala Leu Gly Ser Ala
200 205 210

ggc gat acc gtc aca gtc ttt ggc ctg cca atg gtt ctg aat gac tac 787
Gly Asp Thr Val Thr Val Phe Gly Leu Pro Met Val Leu Asn Asp Tyr
215 220 225

0022700.14204060

tcc gga cag gta ttc cca ccg ctg att gca gca att ggt ctg tac tgg 835
 Ser Gly Gln Val Phe Pro Pro Leu Ile Ala Ala Ile Gly Leu Tyr Trp
 230 235 240 245

gtg gaa aag gga ctg aag aag atc atc cct gaa gca gtc caa atg gtg 883
 Val Glu Lys Gly Leu Lys Lys Ile Ile Pro Glu Ala Val Gln Met Val
 250 255 260

ttc gtc cca ttc ttc tcc ctg ctg att atg atc cca gcg acc gca ttc 931
 Phe Val Pro Phe Phe Ser Leu Leu Ile Met Ile Pro Ala Thr Ala Phe
 265 270 275

ctg ctt gga cct ttc ggc atc ggt gtt ggt aac gga att tcc aac ctg 979
 Leu Leu Gly Pro Phe Gly Ile Gly Val Gly Asn Gly Ile Ser Asn Leu
 280 285 290

ctt gaa gcg att aac aac ttc agc cca ttt att ctt tcc atc gtt atc
 1027
 Leu Glu Ala Ile Asn Asn Phe Ser Pro Phe Ile Leu Ser Ile Val Ile
 295 300 305

cca ttg ctc tac cca ttc ttg gtt cca ctt gga ttg cac tgg cca cta
 1075
 Pro Leu Leu Tyr Pro Phe Leu Val Pro Leu Gly Leu His Trp Pro Leu
 310 315 320 325

aac gcc atc atg atc cag aac atc aac acc ctg ggt tac gac ttc att
 1123
 Asn Ala Ile Met Ile Gln Asn Ile Asn Thr Leu Gly Tyr Asp Phe Ile
 330 335 340

cag gga cca atg ggt gcc tgg aac ttc gcc tgc ttc ggc ctg gtc acc
 1171
 Gln Gly Pro Met Gly Ala Trp Asn Phe Ala Cys Phe Gly Leu Val Thr
 345 350 355

ggc gtg ttc ttg ctc tcc att aag gaa cga aac aag gcc atg cgt cag
 1219
 Gly Val Phe Leu Leu Ser Ile Lys Glu Arg Asn Lys Ala Met Arg Gln
 360 365 370

gtt tcc ctg ggt ggc atg ttg gct ggt ttg ctc ggc ggc att tcc gag
 1267
 Val Ser Leu Gly Gly Met Leu Ala Gly Leu Leu Gly Gly Ile Ser Glu
 375 380 385

cct tcc ctc tac ggt gtt ctg ctc cga ttc aag aag acc tac ttc cgc
 1315
 Pro Ser Leu Tyr Gly Val Leu Leu Arg Phe Lys Lys Thr Tyr Phe Arg
 390 395 400 405

ctc ctg ccg ggt tgt ttg gca ggc ggt atc gtg atg ggc atc ttc gac
 1363
 Leu Leu Pro Gly Cys Leu Ala Gly Gly Ile Val Met Gly Ile Phe Asp
 410 415 420

atc aag gcg tac gct ttc gtg ttc acc tcc ttg ctt acc atc cca gca
 1411
 Ile Lys Ala Tyr Ala Phe Val Phe Thr Ser Leu Leu Thr Ile Pro Ala
 425 430 435

atg gac cca tgg ttg ggc tac acc att ggt atc gca gtt gca ttc ttc
1459

Met Asp Pro Trp Leu Gly Tyr Thr Ile Gly Ile Ala Val Ala Phe Phe
440 445 450

gtt tcc atg ttc ctt gtt ctc gca ctg gac tac cgt tcc aac gaa gag
1507

Val Ser Met Phe Leu Val Leu Ala Leu Asp Tyr Arg Ser Asn Glu Glu
455 460 465

cgc gat gag gca cgt gca aag gtt gct gct gac aag cag gca gaa gaa
1555

Arg Asp Glu Ala Arg Ala Lys Val Ala Ala Asp Lys Gln Ala Glu Glu
470 475 480 485

gat ctg aag gca gaa gct aat gca act cct gca gct cca gta gct gct
1603

Asp Leu Lys Ala Glu Ala Asn Ala Thr Pro Ala Ala Pro Val Ala Ala
490 495 500

gca ggt gcg gga gcc ggt gca ggt gca gga gcc gct gct ggc gct gca
1651

Ala Gly Ala Gly Ala Gly Ala Gly Ala Ala Ala Gly Ala Ala
505 510 515

acc gcc gtg gca gct aag ccg aag ctg gcc gct ggg gaa gta gtg gac
1699

Thr Ala Val Ala Ala Lys Pro Lys Leu Ala Ala Gly Glu Val Val Asp
520 525 530

att gtt tcc cca ctc gaa ggc aag gca att cca ctt tct gaa gta cct
1747

Ile Val Ser Pro Leu Glu Gly Lys Ala Ile Pro Leu Ser Glu Val Pro
535 540 545

gac cca atc ttt gca gca ggc aag ctt gga cca ggc att gca atc caa
1795

Asp Pro Ile Phe Ala Ala Gly Lys Leu Gly Pro Gly Ile Ala Ile Gln
550 555 560 565

cca act gga aac acc gtt gtt gct cca gca gac gct act gtc atc ctt
1843

Pro Thr Gly Asn Thr Val Val Ala Pro Ala Asp Ala Thr Val Ile Leu
570 575 580

gtc cag aaa tct gga cac gca gtg gca ttg cgc tta gat agc gga gtt
1891

Val Gln Lys Ser Gly His Ala Val Ala Leu Arg Leu Asp Ser Gly Val
585 590 595

gaa atc ctt gtc cac gtt gga ttg gac acc gtg caa ttg ggc ggc gaa
1939

Glu Ile Leu Val His Val Gly Leu Asp Thr Val Gln Leu Gly Gly Glu
600 605 610

ggc ttc acc gtt cac gtt gag cgc agg cag caa gtc aag gcg ggg gat
1987

Gly Phe Thr Val His Val Glu Arg Arg Gln Gln Val Lys Ala Gly Asp
615 620 625

002290-1449990

cca ctg atc act ttt gac gct gac ttc att cga tcc aag gat cta cct
 2035
 Pro Leu Ile Thr Phe Asp Ala Asp Phe Ile Arg Ser Lys Asp Leu Pro
 630 635 640 645

ttg atc acc cca gtt gtg gtg tct aac gcc gcg aaa ttc ggt gaa att
 2083
 Leu Ile Thr Pro Val Val Val Ser Asn Ala Ala Lys Phe Gly Glu Ile
 650 655 660

gaa ggt att cct gca gat cag gca aat tct tcc acg act gtg atc aag
 2131
 Glu Gly Ile Pro Ala Asp Gln Ala Asn Ser Ser Thr Thr Val Ile Lys
 665 670 675

gtc aac ggc aag aac gag taacctggga tccatgttgc gca
 2172
 Val Asn Gly Lys Asn Glu
 680

<210> 30
 <211> 683
 <212> PRT
 <213> Corynebacterium glutamicum

<400> 30
 Met Ala Ser Lys Leu Thr Thr Thr Ser Gln His Ile Leu Glu Asn Leu
 1 5 10 15

Gly Gly Pro Asp Asn Ile Thr Ser Met Thr His Cys Ala Thr Arg Leu
 20 25 30

Arg Phe Gln Val Lys Asp Gln Ser Ile Val Asp Gln Gln Glu Ile Asp
 35 40 45

Ser Asp Pro Ser Val Leu Gly Val Val Pro Gln Gly Ser Thr Gly Met
 50 55 60

Gln Val Val Met Gly Gly Ser Val Ala Asn Tyr Tyr Gln Glu Ile Leu
 65 70 75 80

Lys Leu Asp Gly Met Lys His Phe Ala Asp Gly Glu Ala Thr Glu Ser
 85 90 95

Ser Ser Lys Lys Glu Tyr Gly Gly Val Arg Gly Lys Tyr Ser Trp Ile
 100 105 110

Asp Tyr Ala Phe Glu Phe Leu Ser Asp Thr Phe Arg Pro Ile Leu Trp
 115 120 125

Ala Leu Leu Gly Ala Ser Leu Ile Ile Thr Leu Leu Val Leu Ala Asp
 130 135 140

Thr Phe Gly Leu Gln Asp Phe Arg Ala Pro Met Asp Glu Gln Pro Asp
 145 150 155 160

Thr Tyr Val Phe Leu His Ser Met Trp Arg Ser Val Phe Tyr Phe Leu
 165 170 175

Pro Ile Met Val Gly Ala Thr Ala Ala Arg Lys Leu Gly Ala Asn Glu

002250.1324096

180										185					190				
Trp	Ile	Gly	Ala	Ala	Ile	Pro	Ala	Ala	Leu	Leu	Thr	Pro	Glu	Phe	Leu				
195						200					205								
Ala	Leu	Gly	Ser	Ala	Gly	Asp	Thr	Val	Thr	Val	Phe	Gly	Leu	Pro	Met				
210						215					220								
Val	Leu	Asn	Asp	Tyr	Ser	Gly	Gln	Val	Phe	Pro	Pro	Leu	Ile	Ala	Ala				
225						230					235				240				
Ile	Gly	Leu	Tyr	Trp	Val	Glu	Lys	Gly	Leu	Lys	Lys	Ile	Ile	Pro	Glu				
245						250					255								
Ala	Val	Gln	Met	Val	Phe	Val	Pro	Phe	Phe	Ser	Leu	Leu	Ile	Met	Ile				
260						265					270								
Pro	Ala	Thr	Ala	Phe	Leu	Leu	Gly	Pro	Phe	Gly	Ile	Gly	Val	Gly	Asn				
275						280					285								
Gly	Ile	Ser	Asn	Leu	Leu	Glu	Ala	Ile	Asn	Asn	Phe	Ser	Pro	Phe	Ile				
290						295					300								
Leu	Ser	Ile	Val	Ile	Pro	Leu	Leu	Tyr	Pro	Phe	Leu	Val	Pro	Leu	Gly				
305						310					315				320				
Leu	His	Trp	Pro	Leu	Asn	Ala	Ile	Met	Ile	Gln	Asn	Ile	Asn	Thr	Leu				
325						330					335								
Gly	Tyr	Asp	Phe	Ile	Gln	Gly	Pro	Met	Gly	Ala	Trp	Asn	Phe	Ala	Cys				
340						345					350								
Phe	Gly	Leu	Val	Thr	Gly	Val	Phe	Leu	Leu	Ser	Ile	Lys	Glu	Arg	Asn				
355						360					365								
Lys	Ala	Met	Arg	Gln	Val	Ser	Leu	Gly	Gly	Met	Leu	Ala	Gly	Leu	Leu				
370						375					380								
Gly	Gly	Ile	Ser	Glu	Pro	Ser	Leu	Tyr	Gly	Val	Leu	Leu	Arg	Phe	Lys				
385						390					395				400				
Lys	Thr	Tyr	Phe	Arg	Leu	Leu	Pro	Gly	Cys	Leu	Ala	Gly	Gly	Ile	Val				
405						410					415								
Met	Gly	Ile	Phe	Asp	Ile	Lys	Ala	Tyr	Ala	Phe	Val	Phe	Thr	Ser	Leu				
420						425					430								
Leu	Thr	Ile	Pro	Ala	Met	Asp	Pro	Trp	Leu	Gly	Tyr	Thr	Ile	Gly	Ile				
435						440					445								
Ala	Val	Ala	Phe	Phe	Val	Ser	Met	Phe	Leu	Val	Leu	Ala	Leu	Asp	Tyr				
450						455					460								
Arg	Ser	Asn	Glu	Glu	Arg	Asp	Glu	Ala	Arg	Ala	Lys	Val	Ala	Ala	Asp				
465						470					475				480				
Lys	Gln	Ala	Glu	Glu	Asp	Leu	Lys	Ala	Glu	Ala	Asn	Ala	Thr	Pro	Ala				
485						490					495								
Ala	Pro	Val	Ala	Ala	Ala	Gly	Ala	Gly	Ala	Gly	Ala	Gly	Ala	Gly	Ala				
500						505					510								

09604231-062700

Ala Ala Gly Ala Ala Thr Ala Val Ala Ala Lys Pro Lys Leu Ala Ala
515 520 525

Gly Glu Val Val Asp Ile Val Ser Pro Leu Glu Gly Lys Ala Ile Pro
530 535 540

Leu Ser Glu Val Pro Asp Pro Ile Phe Ala Ala Gly Lys Leu Gly Pro
545 550 555 560

Gly Ile Ala Ile Gln Pro Thr Gly Asn Thr Val Val Ala Pro Ala Asp
565 570 575

Ala Thr Val Ile Leu Val Gln Lys Ser Gly His Ala Val Ala Leu Arg
580 585 590

Leu Asp Ser Gly Val Glu Ile Leu Val His Val Gly Leu Asp Thr Val
595 600 605

Gln Leu Gly Gly Glu Gly Phe Thr Val His Val Glu Arg Arg Gln Gln
610 615 620

Val Lys Ala Gly Asp Pro Leu Ile Thr Phe Asp Ala Asp Phe Ile Arg
625 630 635 640

Ser Lys Asp Leu Pro Leu Ile Thr Pro Val Val Val Ser Asn Ala Ala
645 650 655

Lys Phe Gly Glu Ile Glu Gly Ile Pro Ala Asp Gln Ala Asn Ser Ser
660 665 670

Thr Thr Val Ile Lys Val Asn Gly Lys Asn Glu
675 680

<210> 31
<211> 1339
<212> DNA
<213> Corynebacterium glutamicum

<220>
<221> CDS
<222> (101)..(1339)
<223> FRXA02191

<400> 31
ccgattcttt ttgcgcccaa ttogtaacgg cgtactctt aagtggacaa gaaagtctct 60
tgcccgcggg agacagaccc tacgtttaga aaggtttgac atg gcg tcc aaa ctg 115
Met Ala Ser Lys Leu
1 5
acg acg aca tcg caa cat att ctg gaa aac ctt ggt gga cca gac aat 163
Thr Thr Thr Ser Gln His Ile Leu Glu Asn Leu Gly Gly Pro Asp Asn
10 15 20
att act tcg atg act cac tgt gcg act cgc ctt cgc ttc caa gtg aag 211
Ile Thr Ser Met Thr His Cys Ala Thr Arg Leu Arg Phe Gln Val Lys
25 30 35
gat caa tcc att gtt gat caa caa gaa att gac tcc gac cca tca gtt 259

00004224.162760

Asp	Gln	Ser 40	Ile	Val	Asp	Gln	Gln 45	Glu	Ile	Asp	Ser	Asp 50	Pro	Ser	Val	
ctt Leu	ggc Gly	gta Val	gta Val	ccc Pro	caa Gln	gga Gly	tcc Ser	acc Thr	ggg Gly	atg Met	cag Met	gtg Val	gtg Val	atg Met	ggg Gly	307
gga Gly	tct Ser	ggt Val	gca Ala	aac Asn	tat Tyr	tac Tyr	caa Gln	gaa Glu	atc Ile	ctc Leu	aaa Lys	ctt Leu	gat Asp	gga Gly	atg Met	355
aag Lys	cac His	ttc Phe	gcc Ala	gac Asp	ggg Gly	gaa Glu	gct Ala	aca Thr	gag Glu	agt Ser	tca Ser	tcc Ser	aag Lys	aag Lys	gaa Glu	403
tac Tyr	ggc Gly	gga Gly	gtc Val	cgt Arg	ggc Gly	aag Lys	tac Tyr	tcg Ser	tgg Trp	att Ile	gac Asp	tac Tyr	gcc Ala	ttc Phe	gag Glu	451
ttc Phe	ttg Leu	tct Ser	gat Asp	act Thr	ttc Phe	cga Arg	cca Pro	atc Ile	ctg Leu	tgg Trp	gcc Ala	ctg Leu	ctt Leu	ggg Gly	gcc Ala	499
tca Ser	ctg Leu	att Ile	att Ile	acc Thr	ttg Leu	ttg Leu	gtt Val	ctt Leu	gcg Ala	gat Asp	act Thr	ttc Phe	ggg Gly	ttg Leu	caa Gln	547
gac Asp	ttc Phe	cgc Arg	gct Ala	cca Pro	atg Met	gat Asp	gag Glu	cag Gln	cct Pro	gat Asp	act Thr	tat Tyr	gta Val	ttc Phe	ctg Leu	595
cac His	tcc Ser	atg Met	tgg Trp	cgc Arg	tcg Ser	gtc Val	ttc Phe	tac Tyr	ttc Phe	ctg Leu	cca Gln	att Ile	atg Met	gtt Val	ggg Gly	643
gcc Ala	acc Thr	gca Ala	gct Ala	cga Arg	aag Lys	ctc Leu	ggc Gly	gca Ala	aac Asn	gag Glu	tgg Trp	att Ile	ggg Gly	gca Ala	gct Ala	691
att Ile	cca Pro	gcc Ala	gca Ala	ctt Leu	ctt Leu	act Thr	cca Pro	gaa Glu	ttc Phe	ttg Leu	gca Ala	ctg Leu	ggg Gly	tct Leu	gcc Ala	739
ggc Gly	gat Asp	acc Thr	gtc Val	aca Thr	gtc Val	ttt Phe	ggc Gly	ctg Leu	cca Pro	atg Met	gtt Val	ctg Leu	aat Asn	gac Asp	tac Tyr	787
tcc Ser	gga Gly	cag Gln	gta Val	ttc Phe	cca Pro	cgc Pro	ctg Leu	att Ile	gca Ala	gca Ala	att Ile	ggg Gly	ctg Leu	tac Tyr	tgg Trp	835
gtg Val	gaa Glu	aag Lys	gga Gly	ctg Leu	aag Lys	aag Lys	atc Ile	atc Ile	cct Pro	gaa Glu	gca Ala	gtc Val	caa Gln	atg Met	gtg Val	883
ttc Phe	gtc Val	cca Pro	ttc Phe	ttc Phe	ttc Ser	ctg Leu	ctg Leu	att Ile	atg Met	atc Ile	cca Pro	gcg Ala	acc Thr	gca Ala	ttc Phe	931
ctg Leu	ctt Leu	gga Gly	cct Pro	ttc Phe	ggc Gly	atc Ile	ggg Gly	gtt Val	ggg Gly	aac Asn	gga Gly	att Ile	tcc Asn	aac Asn	ctg Leu	979

280 285 290

ctt gaa gcg att aac aac ttc agc cca ttt att ctt tcc atc gtt atc
1027
Leu Glu Ala Ile Asn Asn Phe Ser Pro Phe Ile Leu Ser Ile Val Ile
295 300 305

cca ttg ctc tac cca ttc ttg gtt cca ctt gga ttg cac tgg cca cta
1075
Pro Leu Leu Tyr Pro Phe Leu Val Pro Leu Gly Leu His Trp Pro Leu
310 315 320 325

aac gcc atc atg atc cag aac atc aac acc ctg ggt tac gac ttc att
1123
Asn Ala Ile Met Ile Gln Asn Ile Asn Thr Leu Gly Tyr Asp Phe Ile
330 335 340

cag gga cca atg ggt gcc tgg aac ttc gcc tgc ttc ggc ctg gtc acc
1171
Gln Gly Pro Met Gly Ala Trp Asn Phe Ala Cys Phe Gly Leu Val Thr
345 350 355

ggc gtg ttc ttg ctc tcc att aag gaa cga aac aag gcc atg cgt cag
1219
Gly Val Phe Leu Leu Ser Ile Lys Glu Arg Asn Lys Ala Met Arg Gln
360 365 370

gtt tcc ctg ggt gcc atg ttg gct ggt ttg ctc ggc gcc att tcc gag
1267
Val Ser Leu Gly Gly Met Leu Ala Gly Leu Leu Gly Gly Ile Ser Glu
375 380 385

cct tcc ctc tac ggt gtt ctg ctc cga ttc aag aag acc tac ttc cgc
1315
Pro Ser Leu Tyr Gly Val Leu Leu Arg Phe Lys Lys Thr Tyr Phe Arg
390 395 400 405

ctc ctg ccg ggt tgt ttg gca gca
1339
Leu Leu Pro Gly Cys Leu Ala Ala
410

<210> 32
<211> 413
<212> PRT
<213> *Corynebacterium glutamicum*

<400> 32
Met Ala Ser Lys Leu Thr Thr Thr Ser Gln His Ile Leu Glu Asn Leu
1 5 10 15
Gly Gly Pro Asp Asn Ile Thr Ser Met Thr His Cys Ala Thr Arg Leu
20 25 30
Arg Phe Gln Val Lys Asp Gln Ser Ile Val Asp Gln Gln Glu Ile Asp
35 40 45
Ser Asp Pro Ser Val Leu Gly Val Val Pro Gln Gly Ser Thr Gly Met
50 55 60

Gln Val Val Met Gly Gly Ser Val Ala Asn Tyr Tyr Gln Glu Ile Leu
 65 70 75 80
 Lys Leu Asp Gly Met Lys His Phe Ala Asp Gly Glu Ala Thr Glu Ser
 85 90 95
 Ser Ser Lys Lys Glu Tyr Gly Gly Val Arg Gly Lys Tyr Ser Trp Ile
 100 105 110
 Asp Tyr Ala Phe Glu Phe Leu Ser Asp Thr Phe Arg Pro Ile Leu Trp
 115 120 125
 Ala Leu Leu Gly Ala Ser Leu Ile Ile Thr Leu Leu Val Leu Ala Asp
 130 135 140
 Thr Phe Gly Leu Gln Asp Phe Arg Ala Pro Met Asp Glu Gln Pro Asp
 145 150 155 160
 Thr Tyr Val Phe Leu His Ser Met Trp Arg Ser Val Phe Tyr Phe Leu
 165 170 175
 Pro Ile Met Val Gly Ala Thr Ala Ala Arg Lys Leu Gly Ala Asn Glu
 180 185 190
 Trp Ile Gly Ala Ala Ile Pro Ala Ala Leu Leu Thr Pro Glu Phe Leu
 195 200 205
 Ala Leu Gly Ser Ala Gly Asp Thr Val Thr Val Phe Gly Leu Pro Met
 210 215 220
 Val Leu Asn Asp Tyr Ser Gly Gln Val Phe Pro Pro Leu Ile Ala Ala
 225 230 235 240
 Ile Gly Leu Tyr Trp Val Glu Lys Gly Leu Lys Lys Ile Ile Pro Glu
 245 250 255
 Ala Val Gln Met Val Phe Val Pro Phe Phe Ser Leu Leu Ile Met Ile
 260 265 270
 Pro Ala Thr Ala Phe Leu Leu Gly Pro Phe Gly Ile Gly Val Gly Asn
 275 280 285
 Gly Ile Ser Asn Leu Leu Glu Ala Ile Asn Asn Phe Ser Pro Phe Ile
 290 295 300
 Leu Ser Ile Val Ile Pro Leu Leu Tyr Pro Phe Leu Val Pro Leu Gly
 305 310 315 320
 Leu His Trp Pro Leu Asn Ala Ile Met Ile Gln Asn Ile Asn Thr Leu
 325 330 335
 Gly Tyr Asp Phe Ile Gln Gly Pro Met Gly Ala Trp Asn Phe Ala Cys
 340 345 350
 Phe Gly Leu Val Thr Gly Val Phe Leu Leu Ser Ile Lys Glu Arg Asn
 355 360 365
 Lys Ala Met Arg Gln Val Ser Leu Gly Gly Met Leu Ala Gly Leu Leu
 370 375 380
 Gly Gly Ile Ser Glu Pro Ser Leu Tyr Gly Val Leu Leu Arg Phe Lys

00000001 00000000

385					390					395					400
Lys	Thr	Tyr	Phe	Arg	Leu	Leu	Pro	Gly	Cys	Leu	Ala	Ala			
				405					410						
<210> 33															
<211> 428															
<212> DNA															
<213> Corynebacterium glutamicum															
<220>															
<221> CDS															
<222> (1)..(405)															
<223> FRXA01943															
<400> 33															
cct	gac	cca	atc	ttt	gca	gca	ggc	aag	ctt	gga	cca	ggc	att	gca	atc
Pro	Asp	Pro	Ile	Phe	Ala	Ala	Gly	Lys	Leu	Gly	Pro	Gly	Ile	Ala	Ile
1				5					10				15		48
caa	cca	act	gga	aac	acc	gtt	gtt	gct	cca	gca	gac	gct	act	gtc	atc
Gln	Pro	Thr	Gly	Asn	Thr	Val	Val	Ala	Pro	Ala	Asp	Ala	Thr	Val	Ile
			20					25					30		96
ctt	gtc	cag	aaa	tct	gga	cac	gca	gtg	gca	ttg	cgc	tta	gat	agc	gga
Leu	Val	Gln	Lys	Ser	Gly	His	Ala	Val	Ala	Leu	Arg	Leu	Asp	Ser	Gly
		35					40					45			144
gtt	gaa	atc	ctt	gtc	cac	gtt	gga	ttg	gac	acc	gtg	caa	ttg	ggc	ggc
Val	Glu	Ile	Leu	Val	His	Val	Gly	Leu	Asp	Thr	Val	Gln	Leu	Gly	Gly
	50					55					60				192
gaa	ggc	ttc	acc	gtt	cac	gtt	gag	cgc	agg	cag	caa	gtc	aag	gcg	ggg
Glu	Gly	Phe	Thr	Val	His	Val	Glu	Arg	Arg	Gln	Gln	Val	Lys	Ala	Gly
	65				70				75				80		240
gat	cca	ctg	atc	act	ttt	gac	gct	gac	ttc	att	cga	tcc	aag	gat	cta
Asp	Pro	Leu	Ile	Thr	Phe	Asp	Ala	Asp	Phe	Ile	Arg	Ser	Lys	Asp	Leu
				85					90					95	288
cct	ttg	atc	acc	cca	gtt	gtg	gtg	tct	aac	gcc	gcg	aaa	ttc	ggg	gaa
Pro	Leu	Ile	Thr	Pro	Val	Val	Val	Ser	Asn	Ala	Ala	Lys	Phe	Gly	Glu
			100				105						110		336
att	gaa	ggg	att	cct	gca	gat	cag	gca	aat	tct	tcc	acg	act	gtg	atc
Ile	Glu	Gly	Ile	Pro	Ala	Asp	Gln	Ala	Asn	Ser	Ser	Thr	Thr	Val	Ile
		115					120					125			384
aag	gtc	aac	ggc	aag	aac	gag	taacctggga	tccatggttc	gca						428
Lys	Val	Asn	Gly	Lys	Asn	Glu									
	130				135										
<210> 34															
<211> 135															
<212> PRT															
<213> Corynebacterium glutamicum															
<400> 34															
Pro	Asp	Pro	Ile	Phe	Ala	Ala	Gly	Lys	Leu	Gly	Pro	Gly	Ile	Ala	Ile

1					5					10					15				
Gln	Pro	Thr	Gly	Asn	Thr	Val	Val	Ala	Pro	Ala	Asp	Ala	Thr	Val	Ile				
			20							25				30					
Leu	Val	Gln	Lys	Ser	Gly	His	Ala	Val	Ala	Leu	Arg	Leu	Asp	Ser	Gly				
			35							40				45					
Val	Glu	Ile	Leu	Val	His	Val	Gly	Leu	Asp	Thr	Val	Gln	Leu	Gly	Gly				
			50							55				60					
Glu	Gly	Phe	Thr	Val	His	Val	Glu	Arg	Arg	Gln	Gln	Val	Lys	Ala	Gly				
			65							70				75					
Asp	Pro	Leu	Ile	Thr	Phe	Asp	Ala	Asp	Phe	Ile	Arg	Ser	Lys	Asp	Leu				
			85							90				95					
Pro	Leu	Ile	Thr	Pro	Val	Val	Val	Ser	Asn	Ala	Ala	Lys	Phe	Gly	Glu				
			100							105				110					
Ile	Glu	Gly	Ile	Pro	Ala	Asp	Gln	Ala	Asn	Ser	Ser	Thr	Thr	Val	Ile				
			115							120				125					
Lys	Val	Asn	Gly	Lys	Asn	Glu													
			130							135									

APPENDIX A: DNA SEQUENCES

>RXA00315

TATGATTTTCGGCGGTCCAGTCGGCGGTCTGCTCTTCGGTCTGGTCTACTCACCAATCGTC
 ATCATCGGTCTGCACCAAGCTCTTCCGCGCAATTGAGCTGGAGCTGTTTAAACAGGGTGGGA
 TCCTTCATCTTCGCAACGGCATCTATGGCTAATATCGCCAGGGTGCAGGATGTTTGGCA
 GTGTTCTTCTGGCGAAGAGTGAAGAGCTCAAGGGCTTCAGGTGCTTCAGGTGTCTCC
 GCTGTCTTGGTATTACGGAGCCTCGCATCTTCGGTGTGAACCTTCGCCCTGCGCTGGCG
 TCTTCATCGGTATCGGTACCGCAGCTATCGGTGGCGCTTTGATTGCACCTCTTAAATATC
 AAGGCAGTTGCGTTGGCGCTGCAGGTTCTTGGGTGTTGTTTCTATTGATGCTCCAGAT
 AAGGTCATGTTCTTGGTGTGTCAGTTGTTACCTTCTTCATCGCAATTCGGCGCAGCGATT
 GCTTATGGCCTTTACTTGGTTCGCCGCAACGGCAGCATGATCCAGATGCAACCGCTGCT
 CCAGTGCTCGCAGGAACGACCAAGCCGAAGCAGAAGCACCGCAGAAATTTCAAACGAT
 TCCACCATCATCCAGGCACCTTTGACCGGTGAAGCTATTGCACTGAGCAGCGTCAGCGAT
 GCCATGTTTGCACGCGAAAGCTTGGCTCGGGCGTTGCCATCGTCCCAACCAAGGGCGAG
 TTAGTTTCTCCGGTGAGTGGAAGATTGTGGTGGCATTCACATCTGGCCATGCTTTCGCA
 GTTCGCACCAAGGCTGAGGATGTTTCCAAATGTGGATATCTTGATGCACATTTGGTTTCGAC
 ACAGTAAACCTCAACGGCAGCGACTTTAACCCGCTGAAGAAGCAGGGCGATGAAGTCAAA
 GCAGGGGAGCTGCTGTGTAATTCGATATTGATGCCATTAAAGGCTGCAGGTATGAGGTA
 ACCAGCGCGATTGTTGTTTCGAATTACAAGAAAACCGGACCTGTAAACACTTACGGTTTG
 GCGCAATTAAGCGGGAGCCAACTGCTCAACGTCGCAAAAGAAAGCGGTGCCAGCA
 ACACCA

>RXA00315-downstream
 TAAGTTGAACCTTGAGTGTTCG

>RXA00951

CTCAAGCAATCTTAGAGAAGGCAGCAGCGCGGCGAAGCAGAAGGCTCCTGTCTGTGGCT
 ATGCTGCTTAACACCCACTGCAGCTCCTGCAGCCTCAGTCCCAATCCAAAACCCAGCACAG
 ATCCTCACCGCTCTGTGGCAACGGCTTGGGTACCTCCCTCTTCTCAAACACCCCTTGAG
 CAAGTTTTCGACACCTGGGGTTGGGGTCCATACATGACGCTGGAGGCAACCGACACTATC
 TCGGCCAAGGGCAAGGCCAAGGAGCTGATCTCATATGACCTCTGGTGAATTCGCCGCG
 ACGTTGGGTGATGTTGAATCCCGGTTCAAGTGATCAATGACTTCACGAGCACCGATGAA
 ATCGATGCTCGGCTTCTGTAACGCTACGACATC

>RXA00951-downstream
 TAACTACTTTAAAGGACGAAAA

>RXA01244-upstream

AGATGTCGATTTCTCGAGGAAGAAGTTAAGCCCGAAGAAAACCGTGAATCAGAGCAGGAG
 CGCTTCGACCGCGCTGCAGCCACAGTCTCTTCTTCGT

>RXA01244

TGTGTTGAGCGCTCCGAAGCTGCTGAAGGACCAGCAGCTGAGGTGCTTAAAGCTACTGCT
 GGCATGGTCAATGACCGTGGCTGGCGTAAGCGTGCATCAAGGTTGTCAAGGTGGTGCAC
 CCGTGGGAATACCGCGTGGTTGCAGCAACAACCAAGTTCATCTCCATGTTTCGAAGCCGCA
 GCGGCGCTGATCGCGAGCGCACACAGACTTCGCGGACATCCGCGACCGCGTCATCGCA
 GAACTTCGTGGCGATGAAGAGCCAGGTTCGCGAGCTGTTCCGGACAGGTCATTCTCTTT
 GCAACTGACCTCTCCCAACGACACACCGCGGCACTAGACACAGATCTCTTGTGGGACTT
 GTCAGTGAGCTGGTGGCCCAACGAGCCACACCGCATCATCGCACGCCAGCTCAACGTG
 CCTTGCACTGTCGATCCGGCGCGCGCATCAAGGACATCAAGTCCGCGCAAGAGGTGCT
 ATCGACGCGACGCTTCGGACCAATTGACCGCAACGCGGACCAAGCTGAAGCAACCAAGCTC
 GTCCTCGAGTCCCCTGAGCGCGCTGCTCGCATCGCGGAGTGAAGGGTCTGACACAAAC
 AAGGACGCGTACCAGCTTCAGCTGTTGGCCAACTGCAAGACGCGCACTCTGCACAGCAG
 GCTGCACAGACCGAAGCAGAAGGCATCGGCTGTTCCGCACCGCAACTGTCTTCCTTCC
 GCCACCGAAGGCCAAGCGTTGATGAGCAGGCTGCGGTCTACTCAAAGTGTGCTTGAAGCA
 TTCCAGAGATCCAAGTCTGTTGTCGCTCCCTCGACGAGGTTCTGACAAAGCAGTCCCA
 TTCGATCAGATTGGCTGATGAGATGAACCAAGCACTGGGTGTTCTGGGCTGCGATTCGCA
 CGTGAACAGGTTGATCTGCTGACTCGCCAGCTCGACGCAATTGCGAAGGCCAGCGAAGAA
 CTCGGCGGTGGCGACGACGCCCAACTGGGTATGGCTCCAATGGTGGCTACCGCTTAT

GAAGCAAAGTGGTTTGGCTGACATGTGCCGTGAGCGTGGCCTAATGCCGGCGCCATGATC
GAAGTTCCAGCAGCATCCCTGATGGCAGACAAGATCATGCCCTCACCTGGACTTTGTTTCC
ATCGGTACCAACGACCTGACCCAGTACACCATGGCAGCGGACCGCATGTCTCCTGAGCTT
GCCCTACCTGACCGATCCTTGGCAGCCAGCAGTCTTGCCTGATCAAGCACACCTGTGAC
GAAGGTGCTCGCTTTAACACCCCGGTGGTGTGTTGGTGAAGCAGCAGCAGACCCACTG
TTGGCAACTGTCCTACCCGGTCTTGGCGTGAACCTCCCTGTCCGAGCATCCACTGCTCTC
GCAGCAGTCGGTGCAAGACTGTGAGAGTCAACCTGGAAACCTGTAAGAAGCAGCAGAA
GCAGCACTTGACGCTGAAGGTGCAACTGAAGCACCGCATGCTGTACGCGCAGTGATCGAC
CGACGAGTC

>RXA01244-downstream
TAAACCACCTGTTGAGCTAAAAAG

>RXA01299
ATGGAAATCATGGCCGCGATCATGGCAGCTGGCATGGTCCCACCAATCGCGTTGTCCATT
GCTACCCCTGCTGCGCAAGAAGCTGTTACCCAGCAGCAGCAAGAAAACGGCAAGTCTTCC
TGGCTGCTTGGCCTGGCATTGCTCTCGAAGTGCCATCCCATTCGCCGCGAGCTGACCCCA
TTCCGTGTGATCCAGCAATGATGGCTGGCGGTGCAACCACTGGTGCAATCTCCATGGCA
CTGGGCGTGGCTCTCGGGCTCCACACGGCGGTATCTTGGTGGTCTGGGCAATCGAAGCA
TGGTGGGGTGGCTCATCGCACTTGCAGCAGGCACCATCGTGTCCACCATCGTTGTGATC
GCATGAACGAGTCTTGGCCAAACAAGCGCTCGCTGCAGAGTCGGAAGCAAGAAGCA
CAACAAGCAGCTGTAAACGCA

>RXA01299-downstream
TAATCGGACCTTGACCCGATGTC

>RXA01300-upstream
GATCGACATTAAATCCCTCCCTTGGGGGGTTTAACTAAACAAATCGCTGCGCCCTAATCC
GTTCCGATTACGCGGTAGCAACACGAAGGACACTTTCC

>RXA01300
ATGGCTTCCAAGACTGTAACCGTCGGTTCTCCGTTGGCCTGCAGCAGCTCCAGCATCC
ATCATCGCTGAAGCGGCTGCTGAGTACGACGACGAAATCTTGTGACCCCTGGTTGGCTCC
GATGATGACGAAGAGACCGACGCGTCCCTTCCCTCATGATCATGGCGCTGGGCGCAGAG
CACGGCAACGAAGTTACCGTCACCTCCGACAAACGCTGAAGCTGTTGAGAAGATCGCTGCG
CTTATCGCACAGGACCTTGACGCTGAG

>RXA01300-downstream
TAAACAACGCTCTGCTGTGTTAA

>RXA01503-upstream
GTATCTCTAAAGGCGCTCTAGCTGTTGCAGCTGCAGCGCACTCGGTGGATACGACATCCA
CGACCTATCAAATTCTTTATGCTGCAGGCGATGCTTTTC

>RXA01503
ATGTTCTTGGCAGTCATTTTGGCGATTACTCGCGCTCGTAAATTCGGTGCCAATGTCTTT
ACATCAGTCGCACTCGCTGGTGATTGCTGCACACACAGCTTCAGGCGAGTAACCGTGTG
GTTGAGCGTGAACCTCCAGTCGATGACTCTGTGGCTTCCAAAAGGCTGGTAAATGACGTC
ACCTTCTGGGCAATCCAGTGTGTGCTGCAGTTGGCGTTGCATGTAGCGAGTTGATGAAG
TTGTGCGCA

>RXA01503-downstream
TAAGAGGAGGGCGGTGTCGGTCT

>RXA01883-upstream
CGACTGCGCGCTCTTCTTGGCACTACCATTCCTGCTCTGACCAACTGCCACAGCTG
GTGCAACGGTCACCCAAGTCAAAGGATTGAAAGAAATCAGC

>RXA01883
ATGAATAGCGTAAATAATCTCGCTTGTCGGGTGGATGTCGATTTCCGGGACTCCACC

ACGGATGTCAATCAACAACCTTGGCACTGTTATTTTCGACGCTGGCCGAGCTTCCTCCGCC
 GACGCCCTTGCCAAAGACGCGCTGGATCGTGAAGCAAGTCCGGCACCGGCTTCCTGGT
 CAAGTTGCTATCCCCCACTGCGCTTCGGAAGCCGCTATCTGCCCTACCTTGGCGTTTGCT
 CGCCTGAGCAAGGGTGTGGACTTCAGCGGACCTGATGGCGATGCCAACTTGGTGTTCTCT
 ATTGCAGCACCTGCTGGCGCGGCAAGAGCACCTGAAGATCCTGTCCAAGCTTGCTCGC
 TCCTTGGTGAAGAGGATTTCAATCAAGGCTGTCGAGGAAGCCACCACCGAGCAGGAAATC
 GTGCAGCTTGTGTCGATGCCGTGCTCAACCCAGCACCAAAACCAACCGGACGAGCTGCAGC

>RXA01889-upstream

ACCGAGCCAGCTGCAGCTCCGGCTGCGGCGCGCGTGTGTTAAGAGTGGGCGGCGCTCGAC
 AAGCGTTACTCGTATC

>RXA01889

GTGGCAATCACCGCATGCCAACCGGTATCGCACACACCTACATGGCTGGCGGATTCCTG
 ACGCAAAACGCGGAAGGCCGCGATGATGTGGAACCTGTTGTGGAGACTCAGGGCTCTTCC
 GCTGTACCCCACTCGATCCGAAGATCATCGAAGCTGCCAGCGCGCTCATCTCTGCCACC
 GACGTGGGAGTTAAGACCCGCGAGCGTTTCGCTGGCAAGCCAGTCATTGAATCCGGCGTC
 AAGCGCGCGATCAATGAGCCAGCCAAAGATGATCGACGAGGCCATCGCAGCCTCCAAGAAC
 CCAAAAGCCCGCAAGGTTTCCGGTTCGGTGTGCGCGCATCTGCTGAACCCACCGGCGAG
 AAGCTCGGCTGGGCAAGCGCATCCAGCAGGCAGTCATGACCGCGCTGCTCATATGGTT
 CCAATTCGTAGCTGCCGCGGCGCTCTCTGTGGCTCTCGGCTTCGCATTCTGGTGGATACGAC
 ATGGCGAAGCGCTGGCAAGCAATCGCCACCCAGTTCTCTGACCAACCTGCCAGGCAAC
 ACCGTGATGTTGAC

>RXA01943

CCTGACCCCAATCTTTGCAGCAGGCAAGCTTGGACCAGGCATTGCAATCCAACCAACTGGA
 AACACCGTTGTGTGCTCCAGCAGACGCTACTGTCACTCCTTGTCCAGAAATTCGACACGCA
 GTGGCATTTGCGCTTAGATAGCGGAGTTGAAATCCTTGTCCACGTTGGATTGACACCGTG
 CAATTTGGCGGCGAAGGCTTCACCGTTACGTTGAGCGCAGGCAAGTCAGGCGGGG
 GATCCACTGATCACTTTTGAAGCTGACTTCATTTCGATCCAGGATCTACCTTTGATACCC
 CCACTTGTGGTGTCTAACGCCGCAAAATTCGTTGAAATTTGAAGTATTCTCTGAGATCAG
 GCAATTTCTTCCAGACTGTGATCAAGGTCAACGGCAAGAACGAG

>RXA01943-downstream

TAACCTGGGATCCATGTTGCGCA

>RXA02191-upstream

CCGATTCTTTTTCGCCCAATTTCGTAACGGCGATCCTCTTAAGTGGACAAGAAAGTCTCT
 TGCCCGCGGGAGACAGACCTTACGTTTAGAAAGGTTTGAC

>RXA02191

ATTGCGTCCAAACTGACGACGACATCGCAACATATTTGGAACCTTGGTGGACCAGAC
 AATATTACTTCGATGACTCACTGTGCGACTCGCCTTCGTTCCAAGTGAAGGATCAATCC
 ATTTGTGATCAACAAGAAATGATCTCCGACCCATCAGTTCTTGGCGTAGTACCCCAAGGA
 TCCACCGGTATGCAAGGTGGTATGGGTGGATCTGTTGCAAACTATTACCAAGAAATCCTC
 AAATCTGTATGGAATGAAGCACTTCGCCGACCGTGAAGCTACAGAGAGTTTCATCCAAGAG
 GAATACGGCGGAGTCCGTGGCAAGTACTCGTGGATTGACTACGCGCTTCGAGTTCTTGTCT
 GATACCTTCCGACCAATCTGTGGGCCCTGCTTGGTGCTCACTGATTATTACCTTGTGT
 GTTCTTGGCGGATACTTTGCGTTTGAAGACTTCCGCGCTCCAATGGATGAGCAGCCTGAT
 ACTTATGTATTCCTGCACTCCATGTGGCGCTCGGTCTTCTACTTCTGCCAATATTGGTT
 GGTGCCACCGCAGCTCGAAAGCTCGGCGCAAAAGAGTGGATTGGTGAGCTATTCCAGCC
 GCCTCTTCTACTCCAGAATTTCTTGGCACTGGGTTCGCCGCGGATACCGTCACAGCTCTT
 GGCTGCCAATGGTTCTGAATGACTACTCCGACAGGTATTCACCGCTGATTGACGACA
 ATTGGTCTGTACTGGTGGAAAAGGAGTGAAGAGATCATCCCTGAAGCAGTCCAAGTG
 GTGTTGCTGCCATTCTCTTCCCTGCTGATTATGATCCAGCGACCGCATTCCTGCTTGGGA
 CCTTTCGGCATCGGTGTTGGTAACGGAATTTCCAACTGCTTGAAGCGATTAAACAACTTC
 AGGCCAATTTATCTTTCCATCGTTATCCCATTTGCTCTACCCATTCTTGGTTCACCTTGA
 TTGCATTCGGCACTAAACGCCATCATGATCCAGAACATCAACACCTGGGTTACGACTTC
 ATTCAGGACCAATGGGTGCCTGGAATTCGCGCTGCTTCGGCCTGGTCACCGGCGGTGTC
 TTGCTCTCCATTAAAGAACGAAACAAGGCCATGCGTCAGGTTTCCCTGGGTGGCATGTTG

CTCGGCTGGGGCAAGCGCATCCAGCAGGCAGTCATGACCGCGGTGTCTACATGGTTCCA
 TTCTAGCTGCGCGCGGCTCTCTGTTGGCTCTCGGCTTGCATTCGGTGGATACGACATG
 GCCAACGGCTGGCAAGCAATCGCCACCCAGTTCTCTCTGACCAACCTGCCAGGCAACACC
 CTCGATGTTGACGGCTGGCCATGACCTTCGAGCGTTGAGGCTTCTGTTGTACTTGGCG
 GCAGTCTCTGTTGCCACCGGGCAAGCAGCCATGGGCTTCATCGTGGCAGCCGTGTCTGGC
 TACACCGCATACGCACCTTGCTGCGAGCCAGGCATCGCGCGGGCTTCGTCTGGTGGCGCC
 ATCTCCGTCACCATCGGCGCTGGCTTCATTGGTGGTCTGGTTACCGGTATCTTGGCTGGT
 CTGATTGGCTTGGGATTGGCTCTCGAAGGTGCCAGCGTGGTGCAGTCACTGATGCTCT
 GTGGTCATCATCCGCTACTTACCTCAGTGGTTGTGGTCTGCTCATGTACCTCTCTGCTG
 GGTGCCCATTCGCATCCATCATGACTGGTTTGCAGGACTGGCTATCGTCAATGTCCGGA
 AGCTCCGCCATCTTGTGGGTATCATCTTGGGCTCATGATGTGTTTCGACCTCGCGGGA
 CCGATTAACAGGCGAGCTTACCTCTTGTACCGCGAGGCTGTCTACCGCGGACCAAGCT
 TCCATGGAAATCATGGCGGATCATGGCAGCTGGCATGGTCCCACCAATCGGTTGTTC
 ATTGCTACCTGCTGCGCAAGAAGCTGTTCAACCCAGCAGAGCAAGAAAGCGCAAGTCT
 TCCGTGGCTGCTGGCGTGGCATTCGCTCCGAAGTGCCATCCCATTCGCGGAGCTGAC
 CCATTCCGTGTGATCCAGCAATGATGGCTGGCGGTGCAACCACTGGTGCATCTCCATG
 GCACCTGGCGCTCGGCTCTCGGGCTCCACACGGCGGTATCTTCGTGGTCTGGGCAATCGAA
 CCATGGTGGGCTGGCTCATCGCACTTCGAGCAGCAACCATCGTGTCCACCATCGTTGTC
 ATCGCACTGAAGCAGTTCTTGCCAAACAAGGCCGCTCGCTGCGAAGTCGCGAAGCAAGAA
 GCACAACAGCAGCTGTAACGCA

>RXN01299-downstream
 TAATCGGACCTTGACCGATGTC

>RXN01943-upstream
 CGGATCTTTTTTCGGCCCAATTCGTAACGGCGATCCTCTTAAGTGGACAAGAAAGTCTCT
 TGCCCGCGGAGACAGACCTACGTTTAGAAAGGTTTGAC

>RXN01943
 ATGGCGTCCAACTGACGACGACATCGCAACATATTCTGAAACCTTGGTGGACAGAC
 AATATTACTTCGATGACTCACTGTGCGACTCGCTTCTGCTTCCAAGTGAAGGATCAATCC
 ATTGTGATCAACAAGAAATGACTCCGACCCATCAGTTCCTGGCGTAGTACCCCAAGGA
 TCCACCGGTATGCAAGTGGTGTGATGGGTGGATCTGTGCAAACTATTACCAAGAAATCTTC
 AAATCTGATGGAATGAAGCACTTCGCGCAGCGTGAAGCTACAGAGAGTTTCAATCCAAGAAG
 GAATACGGCGGAGTCCGTGGCAGTACTCGTGGATTGACTACGCTTCGAGTTCTGTGTC
 GATACTTTCGACCAATCTCTGGGCCCTGCTTGGTGCCTCACTGATTATTACCTTGTGTG
 GTTCTTGGGATACTTTCGGTTTGAAGACTTCCGCGCTCCAATGATGAGCAGCCTGAT
 ACTTATGATTCTCGACTCCATGTGGCGCTCGGCTCTCTACTTCTGCCAATATGAGT
 GGTGCCACCGCAGCTCGAAGCTCGGCGCAACAGAGTGGATTGGTGCAGCTATTCCAGCG
 GCACCTCTTACTCCAGAATTCTTGGCACTGGGTCTGCGCGGCGATACCGTACAGTCTTT
 GGCTGCCAATGGTTCTGAATGACTACTCCGACAGGTATTCCACCGCTGATTGCAGCA
 ATTGGTCTGACTACTGGGTGGAAGAAGGACTGAAGAAGATCATCCCTGAAGCAGTCCAAATG
 GTGTTCGTCCCACTTCTTCCCTGCTGATTATGATCCCGAGCGACGCACTTCTGCTTGA
 CCTTTCGGCATCGGTGTTGGTAACGGAATTCCAACCTGCTTGAAGCGATTAAACCACTTC
 AGGCCAATTTACTTTCCATCGTTATCCATTGCTCTACCCATTCTTGGTTCCACTTGA
 TTGCATCTGGCACTAAAGCCCATCATGATCCAGAACATCAACACCTGGGTTACGACTTTC
 ATTCAAGGACCAATGGGTGCCGTGGAATCTCGCGCTGCTTCCGCGTGGTCAACCGCGTGTTC
 TTGCTCTCCATTAAAGGAACGAACAGGCCATCGCTCAGGTTTCCCTGGGTGCGATGTG
 GTCGTGTTTGTCTCGGCGGCACTTTCGAGCCTTCCCTCTACCGTGTCTGCTCCGATTCAAG
 AAGACCTACTTCCGCTCTCTCGCGGTTGTTTGGCAGCGGTATCGTGATGGGATCTTC
 GACATCAAGGCGTACGCTTTCGTGTTCACTCCTTGCTTACCATCCGAGCATGGACCA
 TGGTGGGCTACACCATTTGGTATCGAGTTGCATTCTTGGTTCCATGTTCCCTGTCTTC
 GCACCTGGACTACGTTTCCAAAGAGGCGGATGAGGCAAGTGCAAGGTTGTGCTGAC
 AAGCAGGCAAGAAGATCTGAAGGCAAGAGCTAATGCAACTCCTGCAGCTCCAGTAGCT
 GCTGCAGTTGCGGAGCCGCTGAGGTTGAGGAGCGCTGCTGGCGCTCAACCGCGCTG
 GCAGCTAAGCCGAAGCTGGCGCTGGGGAAGTAGTGGACATTGTTTCCCACCTCGAAGG
 AAGGCAATCCACTTTCTGAAGTACCTGACCAATCTTTCGAGCAGGCACTGTCGACCA
 GGCATTGCAATCCAAACCACTGGAACACCGTTGTTGCTCCAGCAGACGCTACTGTCTC
 GTTCTCCAGAAATCTGGACACGCGAGTGGCATTGCGCTTAGATAGCGGAGTTGAATCTT
 CTCGACCTGGGATTGGACACCGTGCATTTGGGCGCGAAGGCTTACCGTTCAAGTTGAG
 CGCAGGCAAGCTCAAGCGGGGATCCACTGATCACTTTGACGCTGACTTCAATCGA

TCCAAGGATCTACCTTTGATCACCCAGTTGTGGTGTCTAACGCCGCGAAATTCGGTGAA
ATTGAAGGTATTCCTGCAGATCAGGCAAAATCTTCCACGACTGTGATCAAGGTCAACGGC
AAGAACGAG

>RXN01943-downstream
TAACCTGGGATCCATGTTGCGCA

>RXN03002-upstream
GGAACTTCGAGGTGTCTTCGTGGGCGGTACGGAGATCTAGCAAGTGTGGCTTTATGTTT
ACCTATCCGAATCAACATGCAGTGAATTAACATCTACTT

>RXN03002
ATGTTTGTACTCAAAGATCTGCTAAAGGCAGAACGCATAGAACTCGAACGCCAGGTCAAC
GATTGGCGTGAAGGCATCCGCGCCGACAGGTGTACTCTAGAAAAGACAACAGCATTTGAT
TCCGCTACACCGATGCCATGATGCCAGCGTGGAGAAAGAAAGCCCTACATTTGGTGT
GCTCCAGGTTTCGCTTTCCGCGACGCCGCCCGCCAGAGCAGTCCGCGAGACCGCTATG
TCGTTGGGTGCGCTGGCCTCCCTGTTTCTTCGGTCAAGTAAAGATGATCCCTCAAT
CTCATCGTTGCTCTCGCTGCCAAGATGCCACCGCATACCCAAGCGATGGCGGCATTG
GCTAAGCTTTAGAAAAATACGAAAGGATCTCGACGAGGCACAAAGT

RXS00315 - upstream
CTCATGGCATCTGCGCGCTTCGCGTTCCTTGCCAGTGTGTGGTTGTTTACCGCAACCAAGCGTTTCGGC
GGCAATGAGTTCTCTGGGCGCGCGTATTGGT

RXS00315
ATGGCGATGGTGTCCCGAGCTTGGTGAACGGCTACGACCTGGCCGCCACCATGGTGGCGGGCGAAATG
CCAATGTGGTCCCTGTTGGTTTAGATGTTGCCAAGCCGGTTACAGGGCACCCTGCTTCTGTGCTG
GTGGTTTCTTGGATTCTGGCAACGATCGAGAAGTCTCTGCACAAGCGACTCAAGGGCACTGCAGACTTC
CTGATCACTCCAGTGTCTGACGTGTGCTGCTACCGGATTCCTPACATTCATCGCCATTTGGGCCAGCAATG
CGCTGGGTGGGCGATGTGCTGCGACACGGTCTACAGGGACTTTATGATTTCGGTGGTCCAGTCCGGCGGT
CTGCTCTCGGTCTGGTCTACTACCAATCGTCATCACTGGTCTGCCAGTCCCTCCGCCAATTGAG
CTGGAGCTGTTTAAACGAGGTGGATCCTTCATCTTCGCAACGGCATCTATGGCTAATATCGCCAGGGT
GCGGCATGTTTGGCAGTGTCTTCTCGCGAAGAGTGAAGGCTCAAGGGCTTCGAGGTGCTTCAGGT
GTCTCGCTGTCTTGGTATTACGGAGCCTGCGATCTTCGGTGTGAACCTTCGCCCTCGCTGGCGGTTC
TTCATCGGATTCGCTACCGCAGCTATCGGTGGCGCTTTGATTGCACTCTTAAATCAAGGCAGTTGCG
TTGGGCGCTGCAGTTCCTTGGGTGTTGTTCTATGATGCTCCAGATATGGTCACTCTTGGTGTG
GCAGTTGTTTACCTTCCTCATCGCATTCGGCGCAGCGATTGCTTATGGCCTTTACTTGGTTCCGCCGCAAC
GGCAGCATTGATCCAGATCGAACCGCTGCTCCAGTGCTCGAGAACGCCAAAGCGAAGCAAGAG
CCCGCAAAATTTCAAACGATTCCACCATCATCCAGCGACCTTTGACCGTGAAGCTTATGCACTGAC
AGCGTCAGCGATGCTATGTTGCCAGCGGAAGGCTTGGCTCGGGCGTTGCCATCTGCCAACCAAGGGG
CAGTTAGTTTCTCCGGTGAGTGGAAGATTGTTGGGCAATCCCATCTGGCGCATGCTTTCGCGATTCGC
ACCAAGGCTGAGGATGGTCCAATGTGGATATCTTGATGCAATTGGTTTCGACACAGTAAACCTCAAC
GGCACCGCACTTTAACCCGCTGAAGAAGCAGGCGGATGAAGTCAAGCAGGGGAGCTGCTGTGTGAATTC
GATATTGATGCCATTAAAGCTCGAGGTTATGAGTAAACCGCGATTGTTGTTTCGAATTACAAGAAA
ACCGGACCTGTAACACTTACGTTTGGGCGAAATGAAGCGGAGGCCAACCTGCTCAACGTGCAAGAG
AAGAAGCGGTTCGACGCAACCA

RXS00315 - downstream
TAAGTTGAACCTTGAGTGTTCG

RXC00953 - upstream
CTTGCAATCCCCA

RXC00953 -
ATGGCGCCACCAACGGTAGGCAACTACATCATGCAGTCTTCACTCAAGGTCTGCAGTTTCGGCGTTGCA
GTTCCCGTGATTCTCTTTGGTGTCCGACCACTTCTTGGTGAACGTGTCGCCGCAATCCAAGGTATTGCT
GCGAAGGTTGTTCCCGGAGCTATCCCGCATTTGGATGACCCGATCGTGTCCCTACGCGCAGAACCGC
GTTCTCATTTGGTTCTGTGCTTCTTCGTCGGTGGCTTGGTGGCTGACTGTTCTTGATCGTGGCTG
AACCCAGCTTTTGGTGTCCGGTTGATTCTGCTGGTTGGTCCCGCACTTCTTCACTGTTGGCGCGCGG
GGCGTTACGGTAAATGCCAGGGTGGTCTGTCAGGAGCAGTATTTGGCGCTTTGCCAACGGTCTTCTG

ATTACCTTCCTCCCTGCTTTCCTGCTTGGTGTGCTTGGTTCCTTCGGGTCAGAGAACACCCACTTTCGGT
GATGCGGACTTTGGTTGGTTCGGAATCGTTGTTGGTTCTGCAGCCAAGGTGGAAGTGTGGCGGGCTC
ATCTTGTGCTCATCATCGCAGCGGTTCTTCTGGGTGGCGCGATGGTCTTCAGAAAGCGCTCGTGAAT
GGGCACTGGGATCCAGCTCCCAACCGTGAGCGCGTGGAAGGCGGAAGCTGATGCCACTCCAACGGCT
GGGGCTCGGACCTACCTTAAGATTGCTCCTCCGGCGGGCGCTCCTACCCACCGGCTCGAAGC

RXC00953 - downstream
TAAGATCTCCAAAACCTGAGAT

RXC03001 - upstream
CCCGGTTACGTGATCAATGACTTCACGAGCACCGATGAAATCGATGCTGCGCTTCGTGAACGCTACGA
CATCTAACTACTTTAAAGGACGNAATATT

RXC03001 -
ATGGACTGGTTAACCATTCCTCTTTTCCTCGTTAATGAAATCCTTGCGGTTCCGGCTTTCCTCATCGGT
ATCATCACCGCCGTTGGGATTGGGTGCCATGGGGCGTTCCGTCGGTCAGGTTATCGGTGGAGCAATCAAA
GCAACGTTGGGCTTTTTGCTCATTTGGTGCGGGTGCCACGTTGGTCACTGCCTCCCTGGAGCCAATGGGT
GCGATGATCATGGGTGCCACAGGCATGCGTGGTGTGTGCCAACGAATGAAGCCATCGCCGGAATCGCA
CAGGCTGAATACGGCGCGCAGGTGGCGTGGCTGATGATTCTGGGCTTCGCCATCTCTTTGGTGTGGGT
CGTTTACCAACCTGCGTTATGTCTTGCTCAACGGACACCAAGTGTGTTGATGTGCACCATGCTCACC
ATGGTCTTGGCCACCGGAAGATTGATGCGTGGATCTTC

00004271.0022700

APPENDIX B: AMINO ACID SEQUENCES

> RXA00315 (1-1086, translated) 362 residues
YDFGGPVVGL LFGLVYSPIV ITGLHQSFPP IELELFNQGG SFIFATASMA NIAQGAACLA
VFFLAKSEKL KGLAGASGV AVLGITEPAI FGVNLRRLWP FFIGIGTAAI GGALIALFNI
KAVALGAAGF LGVVSIDAPD MVMFLVCAVV TFFIAFGAAI AYGLYLVRNRN GSAIDPATAA
FVPAGTTKAE AEAPAEFSND STIIQAPLTG EAIALSSVSD AMFASGKLGS GSAIVPTKGQ
LVSPVSGKIV VAFPSGHAFV VRTKAEDGSN VDILMHIGFD TVNLTGTHFN PLKQKGDEVK
AGELLCEFDI DAIKAAGYEV TPIVVSNYK KTGVPVNTYGL GEIEAGANLL NVAKKEAVPA
TP

> RXA00951 (1-393, translated) 131 residues
IQAILLEKAA PAKQKAPAVA PAVTPTDAPA ASVQSKTHDK ILTVCGNGLG TSLFLKNTLE
QVFDTWGWF YMTVEATDTI SAKGKAKEAD LIMTSGEIAI TLGDVGIPVH VINDFTSTDE
IDALRLERYD I

> RXA01244 (1-1509, translated) 503 residues
LLERSEAAEG PAAEVLKATA GMVNDRGWRK AVIKGVKGGH PAEYAVVAAT TKFISMFEAA
GGIIAERTTD LRDIRDRVIA ELRGDEEPGL PAVSGQVILF ADDLSPADTA ALDITDLFVGL
VTGLGGPTSH TAIIRQLNV PCIVASGAGI KDIKSGEVL IDGSLGTIDR NADEAEATKL
VSESIERAAE IAENKGPQAT KGYRVQLLA NVQDGNSSAQ AAQTEAEIGL LFRTELCFLS
ATEEPSVDEE AAVYSKYLEA FPESKVVVRS LDAGSDKFPV FASMADEMNP ALGVRLRIA
RGQVDLLTRQ LDAIAKASEE LGRGDDAPTW VMADPMVATAY EAKWFADMCER ERGLIAGAMI
EVPASLMAD KIMPHLDVFS IGTNDLTQYT MAADRMSPFL AYLTDPWQPA VLRLIKHTCD
EGARNTFPG VCGEAAADPL LATVLTGLGV NSLSAAATAL AAVGAKLSEV TLETCKKAAE
AALDAEGATE ARDAVRAVID AAV

> RXA01299 (1-441, translated) 147 residues
MEIMAAIMAA GMVPPIALSI ATLLRKKLFT PAEQENGKSS WLLGLAFVSE GAIPFAAADP
FRVIPAMMAG GATTGAISMA LGVGSRAPHG GIFVVAWIEP WWGWLIALAA GTIVSTIVVI
ALKQFENPKA VAREVAKQEA QQAAVNA

> RXA01300 (1-267, translated) 89 residues
MASKTIVTVGS SVGLHARPAS IIAEAAAEYD DEILLTLVGS DDDEETDASS SLIMINALGAE
HGNEVTVTSD NAEAVEKIEA LIAQDLDAE

> RXA01503 (1-249, translated) 83 residues
MFLAVILAIT AARKFGANVF TSVALAGALL HTQLQAVTVL VDGLQSMTL VAFQKAGNDV
TFLGIPVVLQ LALHVASLMK LSR

> RXA01883 (1-480, translated) 160 residues
MNSVNNSSLV RLDVDFGDST TDVINNLATV IFDAGRASSA DALAKDALDR EAKSGTGVFG
QVAIPHCRSE AVSVPTLGFA RLSKGVDFSG PDGDANLVFL IAAAPAGGGKE HLKILSKLAR
SLVKKDFIKA LQEATTEQEI VDVSDAVLNP APKNHRASCS

> RXA01889 (1-555, translated) 185 residues
VAITACPTGI AHTYMAADSL TQNAEGRDDV ELVVETQGGG AVTPVDPKII EAADAVIFAT
DVGVKDRERF AGKVPVIESGV KRAINEPAKM IDEAIAASKN PNARKVSGSG VASAATETGE
KLGWGKRIQ AVMTGVSVMV PFVAAGQLL ALGFAFGGYD MANGWQAIT QFSLNLFNGN
TVDVD

> RXA01943 (1-405, translated) 135 residues
PDPIFAAGLK GPGIAIQPTG NTVVAPADAT VILVQKSGHA VALRLDSGVE ILVHVGLDVT
QLGGEFTVH VERRQVKAG DPLITFDADF IRSKDLPLIT PVVVSNAAKF GEIEGIPADQ
ANSSTTVIKV NGKNE

> RXA02191 (1-1239, translated) 413 residues
MASKLTTTQS HILENLGGPD NITSMTHCAT RLRFQVKDQS IVDQOEIDSD PSVLGVVPQG
STGMQVMMG SVANYQEIL KLDGMKHFD GEATESSSKK EYGGVRGKYS WIDYAFEPFS
DTFEPILWAL LGASLIITLL VLADTFGLQD FRAMPDEQPD TYVFLHSMWR SVYFELPMV
GATAARKLGA NEWIGAAIPA ALLTPEFLAL GSAGDTVTVF GLPMVLNDYS QGVFPLPIAA

BGI-122CP-APPENDIX B

IGLYWVEKGL KKIPEAVQM VFVFFSLLI MIPATAFLLG PFGIGVNGNI SNLLEAINNF
SPFILSIVIP LLYPFLVPLG LHWPLNAIMI QNINTLGYDF IQGPMGAWN ACFGVLTVGF
LLSIKERNKA MRQVSLGGL AGLLGGISEP SLYGVLLRFK KTYFRLLPGC LAA

>RXN01244 TRANSLATE of: rxn01244.seq check: 8583 from: 1 to: 1704
VATVADVNDQTVLKGTGVVGVRYASAVNITPRPELPQAGEVVAEENREAEQERFDAAA
TVSSRLLERSEAAEGPAAEVLKATAGMVNDRGRKAVIKGVKGHPEAYVAAATKFIS
MFEAAAGLIAERTTDLRIDRIVIAELRGDEEGLPAVSGQVILFADDLSPADTAAALD
LFVGLVTELGGPSTHTAIARQLNVPCIVASGAGIKDIKSGEVLIDGSLGTTDRNDEA
EATKLVSSESLERAAIAEWKGPQATKDGVRVQLLANVDQNSAQQAQTEAGIGLFRTE
LCFLMATEEPSVDEQAAYVSKVLEAFPEKSVVRSLDAGSKDPVFPASMADEMNPLGVR
GLRIARGQVOLLTRQLDAIAKASEELRGDDAPTWVMAMPVATAYEAKWFADMCERGLI
AGAMIEYPAASLMADKIMPHLDEFSIGTNDLTQYMAADRMSPELAYLTPWQPAVLRLI
KHTCDEGARENTPVGVCGEAAADPLLATVLTGLGVNSLSAASSTALAAVGAKESEVTELETC
KKAAEALDAEGATEARDAVRAVIDAAV

>RXN01299 TRANSLATE of: rxn01299.seq check: 4359 from: 1 to: 2064
MNSVNNSSLVRLVDVFGDSTTDVINNLATVIFDAGRASSADALAKDALDREAKSGTGVPG
QVAIPHCRSEAVSVPTLGFARLSKGVDFSGPDGDANLVFLIAAPAGGGKHLKILSKLAR
SLVKDFIKALQEATTEQEIVDVVDAVLNPAKKTTEFAAPAAAAVAESGAASTSVTRIV
ATACPTGIAHTYMAADSLTQNAEGRDDVELVVTQGSASVTPVDPKII EAADAVIFATD
VGKDRERFAGKPVIESGVKRAINEPAKMIDEAIAASKNPNARKVSGSGVAASAEETGEK
LGWKKRIQQAVMTGVSYMVPEVAAGLLALLGAFAGGYDMANGWQAIATQFSLTNLPGNT
VDVDGVMATFERSGFLLYFGAVLFATQAAAGFIVAALSGYTAVALAGRPGLAPFVGGA
ISVTIAGGFIIGGLVTGILAGLIALWIGSKNKVPRVQSLMPVVI I PLLTSVVGLVMYLL
GREPLASIMTGLQDWLSSMSGSSAILLGIILGMMCFDLGPGVYNKAAAYLFGTAGLSTGQQA
SMEIMAAIMAAAGMVPPIALSIATLLRKKLFTPAEQENGKSSWLLGLAFVSEGAIFPAAD
PFRVIPAMMAGGATTGAISMALGVGSRAPHGGI FVVWAIEPWWGWLIALAAGTIVSTIVV
IALKQFWNPAAVAEVAKEQAAQAAVNA

>RXN01943 TRANSLATE of: rxn01943.seq check: 1650 from: 1 to: 2049
MASKLTTTSQHILENLGGPDNITSMTHCATRLRFQVQDQSIQVDDQCEIDSDPSVLGVVPG
STGMQVVMGGSVANYQEILLKLDGMKHFADGEATESSSKKEYGGVRGKYSWIDYAFELFS
DTRFPLNALGASLIITLLVLADTFGLQDFRAPMDEQDPTVYVFLHSMWRSVVFYFLPIMV
GATAARKLGANEWIGAAIPALLTPEFLALGSAGDTVTVFGLPMVNLNDYSGQVFPFLIAA
IGLYWVEKGLKKIIPAEQVMVFVFFSLLIMIPATAFLGPGFIVGVNGNISNLEAINNF
SPFILSIVIP LLYPFLVPLGLHWPNAIMI QNINTLGYDFIQGPMGAWNACFGVLTVGF
LLSIKERNKAMRQVSLGGLAGLLGGISEPSLYGVLLRFKTYFRLLPGCLAGGIVMGIF
DIKAYAFVFTSLTIPAMPDPLGYTIGIAVAFVFSMFLVLDYRSNEERDEARAKVAAD
KQAEEDLKAENATPAAPVAAAGAGAGAGAGAAGAATAVAAPKLAAGEVVDIVVPSLEG
KAIPLSEVPDPIFAAGKLGPGIATQPTGNTVVAADATVILVQKSGHAAVALRLDSGVEIL
VHVGDLTVQLGGEFTVHVERRQVQKAGDELITFDADFIRSKDLPLITPVVVSNAAKFGE
IEGIPADQANSTTVIKVNGKNE

>RXN03002 TRANSLATE of: rxn03002.seq check: 5800 from: 1 to: 408
MFVLKDLLKAERIELDRTVTDREGIRAAGVLEKTNISIDSAYTDAMIASVEEKGPYIVV
APGFAFAHARPSRAVERAMSWRLASPSVFGHKSNDPLNLIVALAAKDATAHTQMAAAL
AKALGKYRKDLDEAQS

>RXS00315 TRANSLATE of: RXS00315.seq check: 1474 from: 1 to: 1404
MAMVFPSSLVNGYDVAATMAAGEMPMSWFLGLDVAQAGYQGTVLPVLVVSWILATIEKFLHKLKRGTADE
LITPVLTLTGFLTPTAIGPAMRWGVDVLAHGLQGLYDFGPGVGLLFLGLVYSPVITGLHQSFPPTE
LELFNGGSGFTFATASMANIAQGAACLAVFFLAKSEKLKGLAGASGVSAVLGITEPAIFGNLRLRWPF
FIGITAAIGGALIALENIKAVAGLGAAGFLGVVSIDAPDMVMEFLVCVVTFFIAFGAAIAYGLYLVRN
GSIDPDATAAPVPAGTTKAEAEAPAEFSNDSTIIQAPITGEATLSSVS DAMFASGKLGSGVAIVPTKG
QLVSPVSGKIVVAFPSGHAFVARTKAEDGSNDILMHIGDFTVNLNGTHFNPLKQGGDEVKAGELLCEFD
DIDAIKAGYEVTPPIVVSNIYKKTGPVNTYGLGEIEAGANLLNVARKEAVPATP

>RXC00953 TRANSLATE of: RXC00953.seq check: 8687 from: 1 to: 753

MAPPTVGNIMQSFTQGLQFGVAVAVILFGVRTILGELVPAFQGIAAKVVPGAIPALDAPIVFFPYAQNA
VLIGFLSSFFVGGLVGLTVLASWLNPAFGVALILPGLVPHFFTGGAAGVYGNATGRRGAVFGAFANGLL
ITFLPAFLLGVLGSFGSENTTFGDADFGWFGIVVGSAAKVEGAGGLILLIIAAVLLGGAMVPQKRNVN
GHWDPAPNRERVEKAEADATPTAGARTYPKIAPPAGAPTPPARS

>RXC03001 TRANSLATE of: RXC03001 .seq check: 9853 from: 1 to: 453
MDWLTIPFLVNEILAVPAFLIGIITAVGLGAMGRSVGQVIGGAIKATLGFLLIGAGATLVTASLEPLG
AMIMGATGMRGVVPTNEAIAGIAQAEYGAQVAWLMILGFAISLVLARFTNLRYVLLNGHHVLLMCTMLT
MVLATGRVDANIF

09604231.062700